

A note on selective line-graphs and partition colorings

Denis Cornaz^{a,*}, Fabio Furini^a, Enrico Malaguti^b, Alberto Santini^{c,d}

^a University Paris-Dauphine, LAMSADE, Paris, France

^b Università di Bologna, Dipartimento di Ingegneria dell'Energia Elettrica e dell'Informazione "Guglielmo Marconi", Bologna, Italy

^c Pompeu Fabra University, Barcelona, Spain

^d Barcelona Graduate School of Mathematics, Spain

A B S T R A C T

We extend a one-to-one correspondence between the set of all colorings of any graph and the set of all stable sets of an auxiliary graph, from graphs to partitioned graphs. This correspondence has an application to Selective Coloring and to Selective Max-Coloring.

Keywords:

Selective/partition coloring

Line graph

Stable set

Max-coloring

1. Introduction

Throughout, G is a simple undirected graph with vertex set $V(G)$ and edge set $E(G)$, with no loops nor parallel edges. A subset of $V(G)$ is *stable* if it contains no edges and it is a *clique* if it has all possible edges. A *coloring* of G is a partition of $V(G)$ into stable sets. The *chromatic number* $\chi(G)$ is the minimum number of stable sets in a coloring of G . The *coloring problem*, that is determining $\chi(G)$, is one of the most studied in graph theory and optimization, and one of the most important class is probably that of perfect graphs, those the induced subgraphs of which have a coloring with no more colors than the biggest clique, see e.g. [19].

A generalization of the coloring problem is the *selective coloring problem* (SCP, also known as *the partition coloring problem*). In this version, the vertices of G are partitioned into disjoint clusters, which can be assumed stable sets, and only one vertex needs to be colored in each cluster. Throughout the paper, we let $\mathcal{P} = \{V_1, \dots, V_{|\mathcal{P}|}\}$ denote a partition of $V(G)$ into non-empty stable sets, and we will refer to the pair $(G, \mathcal{P})$ as a *partitioned graph*. The members V_j of $\mathcal{P}$ are called *clusters*.

A *selective coloring* of $(G, \mathcal{P})$ is a collection of pairwise disjoint stable sets $S_1, \dots, S_k$ of G so that a S_i hits a cluster at most once and every cluster is touched by exactly one S_i . Formally, $|S \cap V_j| = 1$ for all $V_j \in \mathcal{P}$, where $S = \cup_i S_i$ is the (disjoint) union of the S_i 's. The *selective chromatic number* $\chi(G, \mathcal{P})$ of $(G, \mathcal{P})$ is the

minimum k over all selective colorings $S_1, \dots, S_k$. In other words, select one vertex per cluster to induce a subgraph H of G with $\chi(H)$ minimum. Observe that if every cluster V_j is a singleton, then a selective coloring is a coloring and $\chi(G, \mathcal{P})$ is the usual chromatic number.

The SCP is important for its various applications [8], in particular in telecommunications [14,15], and as a challenging problem to solve in practice [11,13,20]. Its algorithmical complexity has been investigated in several classes of graphs [9] including a worse-case version of the problem [7]. Conform and perfect partitioned graphs have been studied in [1], in particular, the class of i -threshold graphs is introduced to characterize primal-dual integrality of associated polyhedra. An auxiliary graph is introduced in [5,16], called the *sandwich line-graph* in [6] where it is shown to provide a chromatic version of Gallai identities and to strengthen Lovász's sandwich theorem, that is, the semidefinite programming (SDP) relaxation for the classical coloring problem. The sandwich line-graph is used in [3,17] to strengthen the linear programming (LP) approach. In [2] this auxiliary graph is used to find classes of graphs in which the coloring problem can be solved in polynomial time. An extension of the coloring problem, with weights, namely Max-Coloring, has been studied for its applications in batch scheduling [10]. In [4], the approach of [5] to solve Max-Coloring with the auxiliary graph, is shown to be efficient in practice and is extended to other coloring problems.

A first contribution of this paper is a procedure to construct an auxiliary graph of a partitioned graph, to extend some properties of the sandwich line-graph to the SCP. A further generalization of the problem is the *Selective Max-Coloring Problem* (SMCP). In this version each cluster V_j has an associated weight $c(V_j)$. Each

color gets the weight of the heaviest cluster it covers and the objective is to find a selective coloring of minimum weight. A second contribution of this paper is to show that the line-graph method for the SCP can be easily extended to the SMCP and that it outperforms the natural Mixed-Integer model for the SMCP in computational tests.

Previous results. The *clique-partition number* $\bar{\chi}(G)$ is equal to the chromatic number $\chi(\bar{G})$ of the complementary graph $\bar{G}$ of G . An equivalent definition for a perfect graph is that it satisfies $\alpha(G) = \bar{\chi}(G)$ for all subgraphs, where the *stability number* $\alpha(G)$ of G is the maximum cardinality of a stable set of G . A graph parameter $\beta(G)$ is said to be *sandwich* if it satisfies $\alpha(G) \leq \beta(G) \leq \bar{\chi}(G)$ for all graphs G . For instance, the fractional clique-partition number $\bar{\chi}_f(G)$ is sandwich, and the so-called sandwich theorem by Lovász states that the theta function $\vartheta(G)$ is a sandwich graph parameter [19]. The main definition in this section is the following:

Definition 1.1. A pair of arcs $\{e, f\}$ of a digraph is *simplicial* if

1. e and f have the same tail,
2. the heads of e, f are adjacent.

Let $L(G)$ be the *line-graph* of G , that is, it has vertex-set $V(L(G)) = E(G)$ the edge-set of G , with an edge $ef \in E(L(G))$ if e and f are adjacent in G . A *sandwich line-graph* of G is any graph built from $L(G)$ by removing edges corresponding to simplicial pairs of arcs in some orientation $\vec{G}$ of G without 3-dicycles. Given an orientation $\vec{G}$, the corresponding sandwich line-graph is denoted by $S(\vec{G})$. Before extending sandwich line-graphs to partitioned graphs, let us summarize their properties. For every sandwich graph parameter $\beta(\cdot)$, for any graph G , and for any orientation $\vec{G}$ of G without 3-dicycle, the following *chromatic Gallai identities* hold (the right equality has been observed in [5,16], and the left equality in [6]):

$$\alpha(G) + \bar{\chi}(S(\vec{G})) = |V(G)| = \alpha(S(\vec{G})) + \bar{\chi}(G) \quad (1)$$

which implies that the modified parameter $|V(G)| - \beta(S(\vec{G}))$ is still sandwich. Restricted to triangle-free graphs, $S(\vec{G}) = L(G)$ and (1) is equivalent to the classical Gallai identities ($\alpha(G) + \tau(G) = |V(G)| = \nu(G) + \rho(G)$, see e.g. [19]). Sandwich line-graphs improve LP bound for stability ($\bar{\chi}_f$) and SDP bound for coloring (ϑ), since the following inequalities hold [6]:

$$\begin{aligned} \alpha(G) &\leq |V(G)| - \bar{\chi}_f(S(\vec{G})) \leq \bar{\chi}_f(G) \\ \vartheta(G) &\leq |V(G)| - \vartheta(S(\vec{G})) \leq \bar{\chi}(G) \end{aligned} \quad (2)$$

2. A line-graph solution method for the selective coloring problem

The *complementary partitioned graph* $(\bar{G}, \mathcal{P})$ of $(G, \mathcal{P})$ is the partitioned graph with an edge uv if and only if u, v belong to different clusters and uv is not an edge of G . So the clusters remain stable sets in $(\bar{G}, \mathcal{P})$. We will also consider *partitioned digraph* $(\vec{G}, \mathcal{P})$ that is a partitioned graph where the edges of G have been orientated. A *selective clique-partition* of $(G, \mathcal{P})$ is a selective coloring of $(\vec{G}, \mathcal{P})$. The problem of determining the *selective clique-partition number* $\bar{\chi}(G, \mathcal{P}) = \chi(\bar{G}, \mathcal{P})$ is that of determining the selective chromatic number in the complementary partitioned graph. Trivially, any algorithm computing $\bar{\chi}(G, \mathcal{P})$ solves SCP.

Not all properties of sandwich graphs extend to partitioned graphs but in the following we extend the right equality in (1). As for sandwich line-graphs, there is a 1-to-1 correspondence between selective colorings and stable sets of the auxiliary graph.

Definition 2.1. The *partition line-graph* $L(G, \mathcal{P})$ of a partitioned graph $(G, \mathcal{P})$ is the graph with vertex-set $V(L(G, \mathcal{P})) = E(G)$ the

edge-set of G , with an edge $ef \in E(L(G, \mathcal{P}))$ if there is a cluster V_i so that both e and f have a vertex in V_i .

Clearly, if every cluster is a singleton then $L(G, \mathcal{P}) = L(G)$. In fact, $L(G, \mathcal{P}) = L(G/\mathcal{P})$ where $G/\mathcal{P}$ is the graph obtained by contracting each cluster. Observe also that stable sets of $L(G, \mathcal{P})$ are matchings of G so that no cluster is covered more than once by an edge of the matching. In the following, we let $\vec{G}$ denote some orientation of G .

Definition 2.2. The *selective line-graph* $S(\vec{G}, \mathcal{P})$ of $(G, \mathcal{P})$ is the graph obtained from the partition line-graph $L(\vec{G}, \mathcal{P})$ by removing edges ef corresponding to simplicial pairs of $\vec{G}$.

Theorem 2.1. The equality $\bar{\chi}(G, \mathcal{P}) + \alpha(S(\vec{G}, \mathcal{P})) = |\mathcal{P}|$ holds for any partitioned graph $(G, \mathcal{P})$ and any orientation $\vec{G}$ without 3-dicycle.

Proof. First observe that if a selective clique-partition of $(G, \mathcal{P})$ has a clique of size one, then any vertex in the cluster containing this clique can be replaced by any other vertex of the cluster to form another selective clique-partition with as many cliques. In the following we assume that each cluster has a particular vertex which is automatically chosen to be the clique of size one if there is one in the cluster. Let us prove the equality now.

It suffices to show that there is a 1-to-1 correspondence (up to choosing an arbitrary vertex per cluster) between the set of all stable sets of the selective line-graph $S(\vec{G}, \mathcal{P})$ and the set of all selective clique-partitions of $(G, \mathcal{P})$, and that this correspondence is so that $|\mathcal{K}| + |S| = |\mathcal{P}|$, for any pair $(\mathcal{K}, S)$ in correspondence, where $\mathcal{K}$ is the selective clique-partition of $(G, \mathcal{P})$ and S the stable set of $S(\vec{G}, \mathcal{P})$. Indeed, in particular, it would imply the equality for the two optima. The rest of the proof consists in describing the 1-to-1 correspondence.

Let S be a stable set of $S(\vec{G}, \mathcal{P})$, so S is a set of arcs in the digraph $\vec{G}$. Let U be the set of the vertices of G covered by S and let $(U_1, S_1), \dots, (U_p, S_p)$ be the (non-trivial) components of the partial subdigraph (U, S) of $\vec{G}$. Since two adjacent arcs in S must form a simplicial pair of $\vec{G}$, it follows that (U_i, S_i) is a simplicial out-star of $\vec{G}$, that is, a partial subdigraph whose vertex-set $U_i = \{u_i, u_i^1, \dots, u_i^{|S_i|}\}$ is a clique of G and whose arc-set has the form $S_i = \{u_i u_i^1, \dots, u_i u_i^{|S_i|}\}$, where u_i is the center of S_i and $u_i^1, \dots, u_i^{|S_i|}$ are the leaves of S_i . Since every cluster is a stable set, clearly, no cluster of $(G, \mathcal{P})$ contains two vertices of the clique U_i . Since only two simplicial arcs of S may have a vertex in a same cluster, no cluster contains two centers u_i, u_j . Moreover, no cluster contains a center u_i and a leaf u_i^k . Hence every cluster contains at most one vertex of U . We can build a selective clique-partition $\mathcal{K}$ of $(G, \mathcal{P})$ by choosing the p non-trivial cliques (with at least two vertices) corresponding to the components of (U, S) and complete it with 1-cliques by choosing the particular vertex for each cluster not covered by S . The graph obtained from (V, S) by contracting each cluster is a forest with $|S|$ arcs, with $|\mathcal{K}|$ components, and with $|\mathcal{P}|$ vertices, hence $|\mathcal{K}| + |S| = |\mathcal{P}|$.

Let $\mathcal{K} = \{K_1, \dots, K_{|\mathcal{K}|}\}$ be a selective clique-partition of $(G, \mathcal{P})$. Since $\vec{G}$ has no 3-dicycle, every clique K_i induces an acyclic subdigraph of $\vec{G}$, which thus is spanned by exactly one out-star S_i . So the union $S = \bigcup_i S_i$ is a stable set in the selective line-graph $S(\vec{G}, \mathcal{P})$. Moreover, $|S| = \sum_{i=1}^{|\mathcal{K}|} (|K_i| - 1) = |\mathcal{P}| - |\mathcal{K}|$. $\square$

Selective coloring and clique-partition problems are equivalent, Theorem 2.1 shows how to solve SCP in $(G, \mathcal{P})$ by finding a maximum stable set in the selective line-graph of $(\vec{G}, \mathcal{P})$.

2.1. Example of the transformation

Fig. 1 shows a partitioned graph $(G, \mathcal{P})$ and its partition line-graph $L(G, \mathcal{P})$.





Fig. 1. Partitioned graph and its partition line-graph.

Since the graph G in Fig. 1 has no triangle, then any orientation $\vec{G}$ has no simplicial pair, and then $L(G, \mathcal{P}) = S(\vec{G}, \mathcal{P})$. The stable set $S = \{6|9, 2|3\}$ of $L(G, \mathcal{P})$ corresponds to the selective clique-partition $\mathcal{K} = \{\{0\}, \{2, 3\}, \{6, 9\}\}$ of $(G, \mathcal{P})$, which illustrate Theorem 2.1.

For another example, suppose that we add the edge $\{6, 7\}$ to G and that we choose the orientation $\vec{G}$ from vertex i to vertex j if $i > j$, then $\{(9, 6), (9, 7)\}$ becomes a simplicial pair. In $S(\vec{G}, \mathcal{P})$, the vertex $6|7$ is added, adjacent to vertices $0|1, 1|2, 2|3, 5|7, 7|9, 6|8, 6|9$, and the edge between $6|9$ and $7|9$ is deleted. Thus $S = \{6|9, 7|9, 5|8\}$ is a stable set of $S(\vec{G}, \mathcal{P})$. It corresponds to the selective clique-partition $\mathcal{K} = \{\{6, 7, 9\}, \{5, 8\}\}$ of $(G, \mathcal{P})$.

2.2. Extension to the Selective Max-Coloring Problem

Max-Coloring is a weighted extension of the classical coloring problem. Given (G, c) where G is a graph and c is a cost vector in $\mathbb{Z}^{V(G)}$, the weight of a coloring $\mathcal{K} = \{K_1, \dots, K_{|\mathcal{K}|}\}$ of G is $w(\mathcal{K}) = \sum_{i=1}^{|\mathcal{K}|} \max_{v \in K_i} c(v)$, and it has to be minimized.

We consider a more general problem, extending Max-Coloring to partition graphs. Given $(G, \mathcal{P}, c)$ where $(G, \mathcal{P})$ is a partitioned graph and c is a cost vector in $\mathbb{Z}^{\mathcal{P}}$, defined the weight of a selective coloring $\mathcal{K} = \{K_1, \dots, K_{|\mathcal{K}|}\}$ of G as $w(\mathcal{K}) = \sum_{i=1}^{|\mathcal{K}|} \max_{V_j: V_j \cap K_i \neq \emptyset} c(V_j)$. The SMCP is to find a selective coloring with minimum weight, denoted $\chi(G, \mathcal{P}, c)$. Clearly, when clusters are singletons SMCP is exactly Max-Coloring.

We can reduce SMCP to the Maximum Weighted Stable Set problem. Actually, SMCP is equivalent to finding a selective clique-partition with minimum weight, where the weight is defined similarly. We add two new steps in the transformation of Theorem 2.1:

1. Choose the orientation $\vec{G}$ of the edges of G as follows: for each edge $\{u, v\}$ then $u \in V_i$ and $v \in V_j$ for two distinct clusters V_i, V_j , orientate $\{u, v\}$ from u to v if $c(V_i) \geq c(V_j)$, in the other direction otherwise.
2. Attribute the cost $\vec{c}(uv) = c(V_j)$ to each vertex uv of $S(\vec{G}, \mathcal{P})$, where $u \in V_i, v \in V_j$ and $c(V_i) \geq c(V_j)$.

Let $\{\mathcal{K}, S\}$ be a pair in correspondence as in the proof of Theorem 2.1. So S is the disjoint union of out-stars S_i the center u_i of which is in the cluster spanned by S_i with maximum cost. It follows that

$$\sum_{i=1}^{|\mathcal{K}|} \sum_{k=1}^{|S_i|} \vec{c}(u_i u_i^k) = \sum_{j=1}^{|\mathcal{P}|} c(V_j) - \sum_{i=1}^{|\mathcal{K}|} \max_{V_j: V_j \cap S_i \neq \emptyset} c(V_j)$$

Hence, by denoting $c(\mathcal{P}) = \sum_{j=1}^{|\mathcal{P}|} c(V_j)$, by denoting $\alpha(G, c)$ the maximum of $\sum_{v \in S} c(v)$ over all stable sets of G , and by denoting

$\bar{\chi}(G, \mathcal{P}, c)$ the minimum weight of a selective clique-partition of $(G, \mathcal{P}, c)$, we obtain the following corollary of Theorem 2.1:

Corollary 2.2. $\bar{\chi}(G, \mathcal{P}, c) + \alpha(S(\vec{G}, \mathcal{P}), \vec{c}) = c(\mathcal{P})$.

We conclude mentioning that we performed a set of computational experiments the purpose of which is to compare the performance of our method, based on solving a Maximum Weighted Stable Set Problem on the selective line-graph, with the performance of a state-of-the-art solver on a natural MIP formulation for the SMCP. We used Sewell's Maximum Weight Stable Set solver, part of Held's `exactcolors` package [12]. This method is able to solve all the generated instances (with up to 100 vertices, various densities, and different numbers of clusters) very quickly, whereas using the MIP model results in several unsolved instances and larger average computing times. For all details see our on-line report [18].

References

- [1] F. Bonomo, D. Cornaz, T. Ekim, B. Ries, Perfectness of clustered graphs, *Discrete Optim.* 10 (4) (2013) 296–303.
- [2] F. Bonomo, M. Giandomenico, F. Rossi, A note on the Cornaz-Jost transformation to solve the graph coloring problem, *Inf. Process. Lett.* 113 (18) (2013) 649–652.
- [3] V.A. Campos, R.C. Correa, D.D. Donne, J. Marenco, A. Wagler, Polyhedral studies of vertex coloring problems: The asymmetric representatives formulation. *CoRR abs/1509.02485* (2015).
- [4] D. Cornaz, F. Furini, E. Malaguti, Solving vertex coloring problems as maximum weight stable set problems, *Discrete Appl. Math.* 217 (2017) 151–162.
- [5] D. Cornaz, V. Jost, A one-to-one correspondence between colorings and stable sets, *Oper. Res. Lett.* 36 (6) (2008) 673–676.
- [6] D. Cornaz, P. Meurdesoif, Chromatic Gallai identities operating on Lovász number, *Math. Program.* 144 (1–2) (2014) 347–368.
- [7] M. Demange, T. Ekim, B. Ries, On the minimum and maximum selective graph coloring problems in some graph classes, *Discrete Appl. Math.* 204 (2016) 77–89.
- [8] M. Demange, T. Ekim, B. Ries, C. Tanasescu, On some applications of the selective graph coloring problem, *European J. Oper. Res.* 240 (2) (2015) 307–314.
- [9] M. Demange, J. Monnot, P. Pop, B. Ries, On the complexity of the selective graph coloring problem in some special classes of graphs, *Theoret. Comput. Sci.* 540 (2014) 89–102.
- [10] L. Epstein, M.M. Halldórsson, A. Levin, H. Shachnai, Weighted sum coloring in batch scheduling of conflicting jobs, *Algorithmica* 55 (4) (2009) 643–665.
- [11] F. Furini, E. Malaguti, A. Santini, An exact algorithm for the Partition Coloring Problem, *Comput. OR* 92 (2018) 170–181.
- [12] S. Held, W.J. Cook, E.C. Sewell, Maximum-weight stable sets and safe lower bounds for graph coloring, *Math. Program. Comput.* 4 (4) (2012) 363–381.
- [13] E.A. Hoshino, Y.A. Frota, C.C. de Souza, A branch-and-price approach for the partition coloring problem, *Oper. Res. Lett.* 39 (2011) 132–137.
- [14] G. Li, R. Simha, The partition coloring problem and its application to wavelength routing and assignment, *Proceedings of Optical Networks Workshop*, Richardson, Texas, 2000.

- [15] T. Noronha, C. Ribeiro, Routing and wavelength assign by partition coloring, *European J. Oper. Res.* 171 (2006) 797–810.
- [16] G. Palubeckis, On the recursive largest first algorithm for graph colouring, *Int. J. Comput. Math.* 85 (2) (2008) 191–200.
- [17] G. Palubeckis, Facet-inducing web and antiweb inequalities for the graph coloring polytope, *Discrete Appl. Math.* 158 (18) (2010) 2075–2080.
- [18] A. Santini, sgcp-via-clique, <https://github.com/alberto-santini/sgcp-via-cliques>, (2019).
- [19] A. Schrijver, *Combinatorial Optimization: Polyhedra and Efficiency*, Springer-Verlag, Berlin Heidelberg, 2003.
- [20] O. Seker, T. Ekim, Z.C. Taskin, An Exact Cutting Plane Algorithm to Solve the Selective Graph Coloring Problem in Perfect Graphs. *CoRR* abs/1811.12094 (2018).