

Theoretical and computational study of several linearisation techniques for binary quadratic problems

Fabio Furini¹  · Emiliano Traversi²

Abstract

We perform a theoretical and computational study of the classical linearisation techniques (LT) and we propose a new LT for binary quadratic problems (BQPs). We discuss the relations between the linear programming (LP) relaxations of the considered LT for generic BQPs. We prove that for a specific class of BQP all the LTs have the same LP relaxation value. We also compare the LT computational performance for four different BQPs from the literature. We consider the Unconstrained BQP and the maximum cut of edge-weighted graphs and, in order to measure the effects of constraints on the computational performance, we also consider the quadratic extension of two classical combinatorial optimization problems, i.e., the knapsack and stable set problems.

Keywords Linearisation techniques · Binary quadratic problems · Max cut problem · Quadratic knapsack problem · Quadratic stable set problem

1 Introduction

A widely used technique for solving binary quadratic problems (BQPs) is by formulating them as mixed integer linear programs (MILPs). The main advantage of these linear reformulations is the use of generic MILP solvers, which keep improving their computational performance [see for example Lodi (2010)]. Several linearisation techniques (LTs) have been proposed in the literature, the seminal works of this stream of research are Fortet (1960) and Glover and Woolsey (1973), both based on introducing additional binary variables. Other LTs have been proposed based only on introducing additional continuous variables; in this stream of research we cite the following works: Glover and Woolsey (1974), Glover (1975), and Sherali and Smith (2007). All these LTs are presented in detail in Sect. 2 and compared

✉ Fabio Furini
fabio.furini@dauphine.fr

Emiliano Traversi
emiliano.traversi@lipn.univ-paris13.fr

¹ PSL, CNRS, LAMSADE UMR 7243, Université Paris Dauphine, 75775 Paris Cedex

² Laboratoire d'Informatique de Paris Nord, Université de Paris 13, Sorbonne Paris Ci Clement, 93430 Villetaneuse, France

in Sect. 3, which is devoted to computational results. The LT of Chaovalitwongse et al. (2004) is deeply related to the LT of Sherali and Smith and, for this reason, these two techniques will be discussed together. A further LT is presented by Adams and Sherali (1986), this approach is generalized to design the reformulation–linearisation technique (RLT) in Sherali and Adams (1998). Recently, other interesting LTs have been introduced in Gueye and Michelon (2009) and Hansen and Meyer (2009). In Gueye and Michelon (2009), the authors propose a framework for unconstrained BQP based on two steps: in the first step the objective function is decomposed with a clique covering heuristic algorithm, while in the second step, valid inequalities are added to the model to get a better dual bound. In Hansen and Meyer (2009), a new family of LTs with a linear number of additional variables and constraints is presented. These LTs are based on a polyform representation of the objective function. The authors show that the best LT among the ones proposed is as tight as the LT of Glover and Woolsey (1974) and it requires the solution of an auxiliary LP with a quadratic number of additional variables and constraints.

1.1 Paper contribution

A first contribution of this paper is an overview of the strength of the linear programming (LP) relaxations of several LTs. Our analysis points out new interconnections between the classical LTs and an alternative LT, called extended linear formulation (ELF). This new LT has been inspired by the ideas proposed in Jaumard et al. (1998), where a reformulation specifically conceived for the quadratic stable set problem (QSSP) is proposed. For generic BQPs, the LT proposed by Glover and Woolsey (1974) and ELF are stronger than the LTs proposed by Glover (1975) and Sherali and Smith (2007). The first two LTs require a quadratic number of additional variables and constraints, while the third and the forth require only a linear number.

A new theoretical result presented is the equivalence between the LP-relaxation value of all the LTs introduced for a specific class of BQPs. This class is rather generic and it contains the BQP formulation of the max-cut problem.

An additional contribution of this paper is an extensive computational comparison of the LTs for four different classes of problem with and without linear constraints. In particular, we tested the Unconstrained BQP, the max cut problem and the quadratic extensions of the Knapsack and the Stable Set problems.

2 Linearisation techniques for binary quadratic problems

A generic binary quadratic problem (BQP), with n variables and p constraints, can be formulated using the following quadratic formulation:

$$\min \left\{ \sum_{i=1}^n \sum_{j=1}^n Q_{ij} x_i x_j + \sum_{i=1}^n L_i x_i : x \in K, \quad x \in \{0, 1\}^n \right\},$$

where $Q \in \mathbb{R}^{n \times n}$, $L \in \mathbb{R}^n$, $K = \{x \in \mathbb{R}^n : Ax \geq b\}$, $A \in \mathbb{R}^{p \times n}$ and $b \in \mathbb{R}^p$. Q is a generic symmetric matrix, not restricted to being convex. Without loss of generality, we may assume $Q_{ii} = 0$, since we may incorporate those elements in the linear terms L_i (for a given binary variable x we have $x^2 = x$). In the following, we define as the Linear Programming (LP) Relaxation of a formulation, the same formulation where the constraints $x \in \{0, 1\}^n$ are

substituted by $x \in [0, 1]^n$. In the next sections, we describe several alternatives for LTs for BQPs present in the literature and then we introduce the new Extended Linear Formulation (ELF).

2.1 Glover–Woolsey linear formulation

The standard method for linearising the quadratic terms is the one introduced by Glover and Woolsey and described in Glover and Woolsey (1974). This linear formulation, called GW (Glover and Woolsey 1974), reads as follows.

$$\begin{aligned}
 \text{GW(Glover and Woolsey 1974): } \min & \sum_{i=1}^n \sum_{j=i+1}^n 2Q_{ij}y_{ij} + \sum_{i=1}^n L_i x_i \\
 & y_{ij} \leq x_i & i, j = 1, \dots, n, \quad i < j & (1) \\
 & y_{ij} \leq x_j & i, j = 1, \dots, n, \quad i < j & (2) \\
 & y_{ij} \geq x_i + x_j - 1 & i, j = 1, \dots, n, \quad i < j & (3) \\
 & y_{ij} \geq 0 & i, j = 1, \dots, n, \quad i < j & (4) \\
 & x \in K \\
 & x \in \{0, 1\}^n .
 \end{aligned}$$

A new variable y_{ij} takes the place of the product between the original variables x_i and x_j in the objective function. We recall that GW (Glover and Woolsey 1974) increases the size of the problem by adding $n(n-1)/2$ variables and $4n(n-1)/2$ constraints. These constraints enforce $y_{ij} = x_i x_j$ for all couples of binary variables x_i and x_j .

A constraint is redundant for a formulation if its removal does not change the set of optimal solutions of its LP-relaxation. Accordingly, the number of constraints of GW (Glover and Woolsey 1974) can be reduced as follows:

Proposition 1 *Inequalities (1) and (2), corresponding to non-negative entries of Q , and inequalities (3) and (4), corresponding to non-positive entries of Q , are redundant for GW (Glover and Woolsey 1974).*

Proof See for example Forrester and Greenberg (2008). □

GW (Glover and Woolsey 1974) can be strengthened by applying the so-called Reformulation Linearisation Technique (RLT) presented in Sherali and Adams (1998). The RLT is a procedure divided into two steps: the reformulation step creates additional nonlinear constraints by multiplying the constraints in K by product factors of the binary variables x and their complements $(1-x)$, and subsequently enforces the identity $x^2 = x$. The linearisation step then substitutes a continuous variable for each distinct product of variables. Hence, GW (Glover and Woolsey 1974) can be viewed as first level RLT applied only on the bound constraints $0 \leq x \leq 1$. An alternative way to improve the strength of GW (Glover and Woolsey 1974) is to include the so-called triangle and cycle inequalities [see for example Gueye and Michelon (2009) or Padberg (1989)]. In our analysis we do not consider these enhancements since we are interested in the basic version of GW (Glover and Woolsey 1974).

2.2 Glover linear formulation

In this section we introduce the linearisation described in Glover (1975). This linear formulation, called G (Glover 1975), reads as follows.

$$\begin{aligned}
 \text{G(Glover 1975): } \min & \sum_{i=1}^n w_i + \sum_{i=1}^n L_i x_i \\
 & w_i \leq Q_i^+ x_i & i = 1, \dots, n & (5) \\
 & w_i \geq Q_i^- x_i & i = 1, \dots, n & (6) \\
 & w_i \leq \sum_{j=1}^n Q_{ij} x_j - Q_i^- (1 - x_i) & i = 1, \dots, n & (7) \\
 & w_i \geq \sum_{j=1}^n Q_{ij} x_j - Q_i^+ (1 - x_i) & i = 1, \dots, n & (8) \\
 & x \in K \\
 & x \in \{0, 1\}^n,
 \end{aligned}$$

where Q_i^+ and Q_i^- are suitable upper and lower bounds on the expression $\sum_{j=1}^n Q_{ij} x_j$, respectively. The main idea behind this linearisation is the introduction of the additional set of variables w_i representing the quantity $\sum_{j=1}^n Q_{ij} x_j$ if $x_i = 1$, or taking a value of 0 otherwise. Formulation G (Glover 1975) has the big advantage of using less variables and constraints than GW (Glover and Woolsey 1974). On the other hand, it requires the introduction of the so-called “big-M” constraints, leading to a weaker LP-relaxation.

An important point concerns the computation of Q_i^- and Q_i^+ . In Glover (1975), the author suggests to impose:

$$Q_i^- = \sum_{j=1}^n \min\{0, Q_{ij}\}, \quad Q_i^+ = \sum_{j=1}^n \max\{0, Q_{ij}\}. \quad (9)$$

Subsequently, in several works [see for example Adams et al. (2004) and Wang et al. (2012)] this approach has been improved by taking into account the feasible region K and hence computing the values for Q_i^- and Q_i^+ with more accuracy. This can be done as follows:

$$Q_i^- = \min_{x \in K} \sum_{j=1}^n Q_{ij} x_j, \quad Q_i^+ = \max_{x \in K} \sum_{j=1}^n Q_{ij} x_j. \quad (10)$$

It is easy to check that Eqs. (9) and (10) coincide when $K = \emptyset$. From now on, when we refer to G (Glover 1975) we imply the version with Q_i^- and Q_i^+ defined in (9), since computing Eq. (10) may require the solution of an optimization problem which depends on the structure of K . Regardless of the method used for computing Q_i^- and Q_i^+ , G (Glover 1975) increases the size of the problem by adding only n variables and $4n$ constraints.

However, since we are minimizing and the coefficients of the w variables are positive, it is possible to reduce the number of constraints:

Proposition 2 *Inequalities (5) and (7) are redundant for G (Glover 1975).*

Proof See Adams et al. (2004). □

2.3 Sherali–Smith linear formulation

The third method linearises the quadratic terms using the techniques described in Sherali and Smith (2007). This linear formulation, called SS (Sherali and Smith 2007), reads as follows:

$$\begin{aligned}
 \text{SS(Sherali and Smith 2007): } \min & \sum_{i=1}^n s_i + \sum_{i=1}^n (L_i + Q_i^-) x_i \\
 & y_i = \sum_{j=1}^n Q_{ij} x_j - s_i - Q_i^- \quad i = 1, \dots, n \quad (11) \\
 & y_i \leq (Q_i^+ - Q_i^-)(1 - x_i) \quad i = 1, \dots, n \quad (12) \\
 & s_i \leq (Q_i^+ - Q_i^-) x_i \quad i = 1, \dots, n \quad (13) \\
 & y_i \geq 0 \quad i = 1, \dots, n \quad (14) \\
 & s_i \geq 0 \quad i = 1, \dots, n \quad (15) \\
 & x \in K \\
 & x \in \{0, 1\}^n,
 \end{aligned}$$

where Q_i^+ and Q_i^- are defined as in (9) or in (10). Since the latter case requires the computation of an additional optimization problem, we use (9) as for G (Glover 1975). In Sherali and Smith (2007), the authors introduce three formulations, called BP, $\bar{\text{BP}}$ and BP-strong. However, in absence of quadratic constraints, they are all equivalent and they can be written as SS (Sherali and Smith 2007). The idea behind SS (Sherali and Smith 2007) is similar to the one behind G (Glover 1975), i.e., the introduction of additional variables representing the sum $\sum_{j=1}^n Q_{ij} - Q_i^-$ and using big-M constraints. SS (Sherali and Smith 2007) increases the size of the problem by adding $2n$ variables, $4n$ inequalities and n equations. More precisely, variables y can be projected out using Eq. (11), thus resulting in only n additional variables and $4n$ inequalities (12)–(15).

Formulation SS (Sherali and Smith 2007) is a strengthening of a precedent formulation proposed by Chaovalitwongse et al. (2004) called CPP (Chaovalitwongse et al. 2004) which uses instead the following weaker (see Sherali and Smith 2007) definition of Q_i^+ and Q_i^- :

$$Q_i^+ = \max_i \sum_{j=1}^n |Q_{ij}|, \quad Q_i^- = -\max_i \sum_{j=1}^n |Q_{ij}|.$$

Like for G (Glover 1975), also for SS (Sherali and Smith 2007) [and analogously for CPP (Chaovalitwongse et al. 2004)] some constraints can be eliminated:

Proposition 3 *Inequalities (13) and (14), are redundant for SS (Sherali and Smith 2007).*

Proof See Sherali and Smith (2007). □

2.4 Extended linear formulation

We now introduce a new linearisation, called Extended Linear Formulation (ELF).

$$\text{ELF: } \min \sum_{i=1}^n \sum_{j=1}^n Q_{ij} + \sum_{i=1}^n L_i x_i - \sum_{i=1}^n \sum_{j=i+1}^n 2Q_{ij} (z_{ij}^i + z_{ij}^j)$$

$$z_{ij}^i + z_{ij}^j \leq 1 \quad i, j = 1, \dots, n, \quad i < j \quad (16)$$

$$x_i + z_{ij}^i \leq 1 \quad i, j = 1, \dots, n, \quad i < j \quad (17)$$

$$x_j + z_{ij}^j \leq 1 \quad i, j = 1, \dots, n, \quad i < j \quad (18)$$

$$x_i + z_{ij}^i + z_{ij}^j \geq 1 \quad i, j = 1, \dots, n, \quad i < j \quad (19)$$

$$x_j + z_{ij}^i + z_{ij}^j \geq 1 \quad i, j = 1, \dots, n, \quad i < j \quad (20)$$

$$x \in K$$

$$x \in \{0, 1\}^n.$$

This linearisation increases the size of the problem by adding $n(n-1)$ variables and $5n(n-1)$ constraints. A pair of new variables z_{ij}^i and z_{ij}^j is used instead of the product of variables x_i and x_j . These new variables modify the objective function and appear in the new set of constraints. The total sum of the quadratic costs is paid in the objective function ($\sum_{i=1}^n \sum_{j=1}^n Q_{ij}$) and then the correct value is reconstructed with the use of the z variables ($\sum_{i=1}^n \sum_{j=i+1}^n 2Q_{ij}(z_{ij}^i + z_{ij}^j)$). The values of the z variables are set according to the values of the x variables thanks to constraints (16)–(20), i.e., if $x_i = x_j = 1$ then $z_{ij}^i = z_{ij}^j = 0$, on the other hand, if one or both variables x_i and x_j have a value of 0, then we have $z_{ij}^i + z_{ij}^j = 1$, which implies the correctness of the formulation.

Finally, the following Proposition allows the reduction of the number of constraints also for ELF:

Proposition 4 *Inequalities (19) and (20), corresponding to non-negative entries of Q , and inequalities (16), (17) and (18), corresponding to non-positive entries of Q are redundant for ELF.*

Proof Let $(\tilde{x}, \tilde{z}^i, \tilde{z}^j)$ be an optimal solution of the LP-relaxation of ELF satisfying all inequalities except one of the constraints (19) associated to an entry $Q_{kl} \geq 0$. Let $x_k + z_{kl}^i + z_{kl}^j \geq 1$ be the violated inequality. And w.l.o.g, we assume that $\tilde{z}_{kl}^j \geq 0$, since the other constraints impose: $0 \leq \tilde{z}_{kl}^i + \tilde{z}_{kl}^j \leq 1$. We can increase the value of $\tilde{z}_{kl}^i$ to $\tilde{z}_{kl}^i + \Delta$ with $\Delta = \min\{1 - \tilde{z}_{kl}^i - \tilde{z}_{kl}^j, 1 - \tilde{x}_k - \tilde{z}_{kl}^i\}$. The constraint is violated hence $\tilde{x}_k + \tilde{z}_{kl}^i + \tilde{z}_{kl}^j < 1$, this implies: (i) $0 < 1 - \tilde{z}_{kl}^i - \tilde{z}_{kl}^j$, since $0 \leq \tilde{x}_k$ and $\tilde{x}_k < 1 - \tilde{z}_{kl}^i - \tilde{z}_{kl}^j$; (ii) $0 < 1 - \tilde{x}_k - \tilde{z}_{kl}^i$, since $\tilde{x}_k + \tilde{z}_{kl}^i < 1$ and $\tilde{z}_{kl}^j \geq 0$; and therefore, Δ is positive. By observing that Δ corresponds to the minimum between the slacks of constraints (16) and (17) associated to k and l , the new solution is feasible by construction. The objective function value decreases by $-\Delta Q_{kl}$. This contradicts the fact of being optimal and violating one of the constraints (19). The same holds for constraints (20).

Similar considerations can be done for inequalities (16), (17) and (18) when $Q_{kl} \leq 0$. Hence, eliminating from ELF the inequalities (19) and (20), corresponding to a couple of indices k and l with non-negative Q_{kl} or the inequalities (16), (17) and (18), corresponding to a couple of indices k and l with non-positive Q_{kl} , does not change the set of optimal solutions of the LP-relaxation of ELF (i.e., they are redundant). $\square$

Constraints (16), (17) and (18) provide upper bounds on the variables z , while constraints (19) and (20) provide lower bounds. Hence, according to the sign of the coefficients of the matrix Q and the sense of the objective function, some of them are redundant.

Table 1 Number of additional variables and constraints used in each LT

Formulations	Variables	Constraints	
		Original	Reduced
GW (Glover and Woolsey 1974)	$n(n-1)/2$	$2n(n-1)$	$n(n-1)$
G (Glover 1975), CPP (Chaovalitwongse et al. 2004) and SS (Sherali and Smith 2007)	n	$4n$	$2n$
ELF	$n(n-1)$	$5n(n-1)$	$\approx \frac{5}{2}n(n-1)$

2.5 Formulation summary

In Table 1, we report the number of additional variables and constraints needed by each LT. The columns concerning the constraints are subdivided in two parts: the table column—Original—report the number of inequalities needed by each formulation while the table column—Reduced—report the number of non redundant constraints needed by each formulation (see Propositions 1, 2, 3 and 4). In the following, this reduction of the formulation dimensions will be referred to as Constraint-Redundancy elimination. As far as GW (Glover and Woolsey 1974), G (Glover 1975), CPP (Chaovalitwongse et al. 2004) and SS (Sherali and Smith 2007) are concerned, regardless of the sign of the entries of Q , the Constraint - Redundancy elimination allows to reduce by one half the number of additional constraints. For ELF the situation is slightly different because the number of non redundant constraints depends on the sign of Q and it is hence included in the interval $[2n(n-1), 3n(n-1)]$, for this reason we reported the average value of $\frac{5}{2}n(n-1)$. As explained in Adams et al. (2004), the number of constraints of G (Glover 1975) can be further reduced to n via a variable transformation.

2.5.1 Formulation strength

In this section, we present a comparison of the strength of the LT discussed in the previous sections.

The strong connection between G (Glover 1975) and SS (Sherali and Smith 2007) is given by the following proposition:

Proposition 5 *The polyhedra defined by the LP-relaxations of G (Glover 1975) and SS (Sherali and Smith 2007) are isomorphic under a linear transformation that keeps the values of the x variables unchanged and the relation between the s and the w variables is the following:*

$$s_i = w_i - Q_i^- x_i \quad i = 1, \dots, n.$$

Proof As first step, it is possible to apply to SS (Sherali and Smith 2007) the variable substitution $s_i = w_i - Q_i^- x_i$ and subsequently equations (11) can be used to substitute variables y into the remaining constraints, thus obtaining exactly G (Glover 1975). By applying to G (Glover 1975) the variable substitution $w_i = s_i + Q_i^- x_i$, we obtain SS (Sherali and Smith 2007). $\square$

A direct consequence of Proposition 5 is that the optimal LP-relaxation values of G (Glover 1975) and SS (Sherali and Smith 2007) are identical.

The following proposition shows that ELF is an extended formulation of GW (Glover and Woolsey 1974):

Proposition 6 *The polyhedra defined by the LP-relaxations of GW (Glover and Woolsey 1974) and ELF are isomorphic under a linear transformation that keeps the values of the x variables unchanged and the relations between the z and the y variables are the following:*

$$z_{ij}^i = 1 - x_i, \quad z_{ij}^j = x_i - y_{ij} \quad i, j = 1, \dots, n, i < j.$$

Proof Let $(\bar{x}, \bar{y})$ be a feasible solution to the LP-relaxation of GW (Glover and Woolsey 1974). We consider the following solution $(\tilde{x}, \tilde{z}^i, \tilde{z}^j)$ of the LP-relaxation of ELF: $\tilde{x}_i = \bar{x}_i$, $\tilde{z}_{ij}^i = 1 - \bar{x}_i$, $\tilde{z}_{ij}^j = \bar{x}_i - \bar{y}_{ij}$. It is easy to check that it is feasible and it has the same value. Now let $(\tilde{x}, \tilde{z}^i, \tilde{z}^j)$ be a feasible solution to the LP-relaxation of ELF. We consider the following solution $(\bar{x}, \bar{y})$ of the LP-relaxation of GW (Glover and Woolsey 1974): $\bar{x}_i = \tilde{x}_i$, $\bar{y}_{ij} = 1 - \tilde{z}_{ij}^i - \tilde{z}_{ij}^j$. Again, it is easy to check that it is feasible and it has the same value. This implies that for every feasible solution of the LP-relaxation of GW (Glover and Woolsey 1974) there exists one feasible solution of the LP-relaxation of ELF of the same value and vice versa, proving the statement. $\square$

Also in this case, a direct consequence of Proposition 6 is that the LP-relaxation values of GW (Glover and Woolsey 1974) and ELF are identical. The new linearisation ELF can be seen as an extended formulation (see e.g., Conforti et al. 2010) of GW in the sense that the projection on the (x, y) -space of the polytope defined by (16)–(20), and constraints $y_{ij} = 1 - z_{ij}^i - z_{ij}^j$ ($i, j = 1, \dots, n, i < j$), is equal to the polytope defined by (1)–(4) and by $x \in [0, 1]^n$. Moreover, Proposition 6 also implies that every valid inequality for GW (Glover and Woolsey 1974) is also valid for ELF.

We recall that in the following analysis we use formulas (9), which imply:

$$\sum_{j=1}^n Q_{ij} = Q_i^+ + Q_i^-.$$

To the best of our knowledge, no formal proof concerning the relation between the LP relaxations of GW (Glover and Woolsey 1974) and G (Glover 1975) has been given in the literature. For sake of completeness, we present the following Proposition, that will be useful in the rest of the section:

Proposition 7 *The LP-relaxation of GW (Glover and Woolsey 1974) is stronger than the LP relaxation of G (Glover 1975).*

Proof Let $(\tilde{x}, \tilde{y})$ be a feasible solution of the LP-relaxation of GW (Glover and Woolsey 1974), we consider the solution $(\bar{x}, \bar{w})$ obtained from $(\tilde{x}, \tilde{y})$ as follows:

$$\bar{x}_i = \tilde{x}_i \quad \bar{w}_i = \sum_{j=1}^n Q_{ij} \tilde{y}_{ij} \quad i = 1, \dots, n;$$

both solutions have by construction the same objective function value. We now show that the solution $(\bar{x}, \bar{w})$ is feasible for the LP-relaxation of G (Glover 1975). We notice that, for each $i = 1, \dots, n$, we have:

$$\begin{aligned}
\bar{w}_i &= \sum_{j=1}^n Q_{ij} \tilde{y}_{ij} = \sum_{\substack{j=1, \dots, n : \\ Q_{ij} < 0}} Q_{ij} \tilde{y}_{ij} + \sum_{\substack{j=1, \dots, n : \\ Q_{ij} \geq 0}} Q_{ij} \tilde{y}_{ij} \geq \sum_{\substack{j=1, \dots, n : \\ Q_{ij} < 0}} Q_{ij} \tilde{y}_{ij} \\
&\geq \sum_{\substack{j=1, \dots, n : \\ Q_{ij} < 0}} Q_{ij} \tilde{x}_i = Q_i^- \tilde{x}_i
\end{aligned}$$

where the last inequality derives from $\tilde{y}_{ij} \leq \tilde{x}_i$. Analogously, for each $i = 1, \dots, n$, we have:

$$\begin{aligned}
\bar{w}_i &= \sum_{j=1}^n Q_{ij} \tilde{y}_{ij} = \sum_{\substack{j=1, \dots, n : \\ Q_{ij} < 0}} Q_{ij} \tilde{y}_{ij} + \sum_{\substack{j=1, \dots, n : \\ Q_{ij} \geq 0}} Q_{ij} \tilde{y}_{ij} \\
&\geq \sum_{\substack{j=1, \dots, n : \\ Q_{ij} < 0}} Q_{ij} \tilde{x}_j + \sum_{\substack{j=1, \dots, n : \\ Q_{ij} \geq 0}} Q_{ij} \tilde{x}_i + \sum_{\substack{j=1, \dots, n : \\ Q_{ij} \geq 0}} Q_{ij} \tilde{x}_j - \sum_{\substack{j=1, \dots, n : \\ Q_{ij} \geq 0}} Q_{ij} \\
&= \sum_{j=1}^n Q_{ij} \tilde{x}_j - Q_i^+(1 - \tilde{x}_i)
\end{aligned}$$

where the first inequalities derive from $\tilde{y}_{ij} \leq \tilde{x}_i$ and $\tilde{y}_{ij} \geq \tilde{x}_i + \tilde{x}_j - 1$. The two sequences of inequalities show that the solution $(\bar{x}, \bar{y})$ respects inequalities (6) and (8). Finally, we provide an instance where the optimal value of the LP-relaxation of GW (Glover and Woolsey 1974) is greater than the optimal value of the LP-relaxation of G (Glover 1975). Consider the

following instance with $K = \emptyset$, $n = 3$ and $L = [1 \ -1 \ 8]$; $Q = \begin{bmatrix} 0 & -1 & 1 \\ -1 & 0 & -5 \\ 1 & -5 & 0 \end{bmatrix}$. The

optimal solution of the LP-relaxation of GW (Glover and Woolsey 1974) is $\tilde{x}_1 = 0$, $\tilde{x}_2 = 1$, $\tilde{x}_3 = 1$, $\tilde{y}_{12} = 0$, $\tilde{y}_{13} = 0$, $\tilde{y}_{23} = 1$ and the optimal solution value is -3 . On the other hand, the optimal solution of the LP-relaxation of G (Glover 1975) is $\bar{x}_1 = 0.54$, $\bar{x}_2 = 1$, $\bar{x}_3 = 0.90$, $\bar{w}_1 = -0.54$, $\bar{w}_2 = -5.09$, $\bar{w}_3 = -4.54$ and the optimal solution value is -3.36 . $\square$

We now introduce two lemmas that are used to show a new property of the LP-relaxation of GW (Glover and Woolsey 1974) and G (Glover 1975). Both lemmas use the following Proposition:

Proposition 8 *In case of $K = \emptyset$, given a feasible vector $\bar{x}$ for GW (Glover and Woolsey 1974) (resp. G (Glover 1975)), the optimal values of the $\bar{y}$ (resp. $\bar{w}$) are:*

$$\begin{aligned}
\bar{y}_{ij} &= \begin{cases} \max\{0, \bar{x}_i + \bar{x}_j - 1\} & \text{if } Q_{ij} \geq 0 \\ \min\{\bar{x}_i, \bar{x}_j\} & \text{if } Q_{ij} < 0 \end{cases} \quad i, j = 1, \dots, n, \\
\bar{w}_i &= \max \left\{ Q_i^- \bar{x}_i, \sum_{j=1}^n Q_{ij} \bar{x}_j - Q_i^+(1 - \bar{x}_i) \right\} \quad i = 1, \dots, n.
\end{aligned}$$

Proof The statement is a direct consequence of the sign of the y and w variables in the respective objective functions and of the fact that there are no additional constraints, except for the ones linking the x variables to the y and w variables respectively. $\square$

Lemma 1 If $K = \emptyset$, let $(\tilde{x}, \tilde{y})$ and $(\bar{x}, \bar{w})$ be two optimal solutions of the LP-relaxations of GW (Glover and Woolsey 1974) and G (Glover 1975) respectively. If $\tilde{x}_i = \bar{x}_i = \frac{1}{2}$ for all $i = 1, \dots, n$, both solutions have the same objective function value equal to $\frac{1}{2} \sum_{i=1}^n (Q_i^- + L_i)$.

Proof It is sufficient to compute the objective function value of both solutions. If $Q_{ij} \geq 0$, we have $\tilde{y}_{ij} = \max\{0, \tilde{x}_i + \tilde{x}_j - 1\} = \max\{0, 0\} = 0$. If $Q_{ij} < 0$, we have $\tilde{y}_{ij} = \min\{\tilde{x}_i, \tilde{x}_j\} = \min\left\{\frac{1}{2}, \frac{1}{2}\right\} = \frac{1}{2}$. The objective function value of GW (Glover and Woolsey 1974) is hence the following:

$$\sum_{i=1}^n \sum_{j=1}^n Q_{ij} \tilde{y}_{ij} + \sum_{i=1}^n L_i \tilde{x}_i = \frac{1}{2} \sum_{\substack{i,j=1,\dots,n : \\ Q_{ij} < 0}} Q_{ij} + \frac{1}{2} \sum_{i=1}^n L_i = \frac{1}{2} \sum_{i=1}^n (Q_i^- + L_i);$$

Moreover, we have $\bar{w}_i = \max\{Q_i^- x_i, \sum_{j=1}^n Q_{ij} \bar{x}_j - Q_i^+(1 - \bar{x}_i)\}$, leading to the following objective function value of G (Glover 1975):

$$\begin{aligned} \sum_{i=1}^n \bar{w}_i + \sum_{i=1}^n L_i \bar{x}_i &= \sum_{i=1}^n \max \left\{ Q_i^- \bar{x}_i, \sum_{j=1}^n Q_{ij} \bar{x}_j - Q_i^+(1 - \bar{x}_i) \right\} + \sum_{i=1}^n L_i \bar{x}_i \\ &= \sum_{i=1}^n \max \left\{ \frac{1}{2} Q_i^-, \frac{1}{2} \sum_{j=1}^n (Q_{ij} - Q_i^+) \right\} + \frac{1}{2} \sum_{i=1}^n L_i \\ &= \sum_{i=1}^n \max \left\{ \frac{1}{2} Q_i^-, \frac{1}{2} Q_i^- \right\} + \frac{1}{2} \sum_{i=1}^n L_i = \frac{1}{2} \sum_{i=1}^n (Q_i^- + L_i). \end{aligned}$$

□

Lemma 2 If $K = \emptyset$ and $L_i = -\sum_{j=1}^n Q_{ij}$ ($i = 1, 2, \dots, n$), an optimal solution of the LP-relaxation of GW (Glover and Woolsey 1974) is the following:

$$\begin{aligned} \bar{x}_i &= \frac{1}{2}, \quad i = 1, \dots, n, \\ \bar{y}_{ij} &= \begin{cases} 0 & \text{if } Q_{ij} \geq 0; \\ \frac{1}{2} & \text{if } Q_{ij} < 0 \end{cases} \quad i, j = 1, \dots, n. \end{aligned}$$

Proof The objective function is separable, hence, we measure the impact of changing one single variable at a time. We consider the solutions $\hat{x} = \bar{x} \pm \epsilon e_{\hat{i}}$ with $\hat{i} = 1, \dots, n$ and $\epsilon \geq 0$, where $e_{\hat{i}}$ is an n -dimensional vector with value 1 in the $\hat{i}$ -th component and 0 otherwise. We consider first the case $\hat{x} = \bar{x} + \epsilon e_{\hat{i}}$. Proposition 8 implies that the $\hat{y}$ variable corresponding to $\hat{x} = \bar{x} + \epsilon e_{\hat{i}}$ is equal to $\bar{y}$, except for $i = \hat{i}$, where we have:

$$\hat{y}_{\hat{i}j} = \begin{cases} \epsilon & \text{if } Q_{\hat{i}j} \geq 0 \\ \frac{1}{2} & \text{if } Q_{\hat{i}j} < 0 \end{cases} \quad j = 1, \dots, n.$$

If we consider the objective function value of the LP-relaxation of GW (Glover and Woolsey 1974) we obtain, for a given $\hat{i}$:

$$\begin{aligned}
& \sum_{i=1}^n \sum_{j=1}^n Q_{ij} \hat{y}_{ij} + \sum_{i=1}^n L_i \hat{x}_i = \sum_{i=1}^n \sum_{j=1}^n Q_{ij} \hat{y}_{ij} - \sum_{i=1}^n \left(\sum_{j=1}^n Q_{ij} \right) \hat{x}_i \\
& = \sum_{i=1}^n \sum_{j=1}^n Q_{ij} \bar{y}_{ij} + \epsilon \sum_{\substack{j=1, \dots, n : \\ Q_{ij} \geq 0}} Q_{ij} - \sum_{i=1}^n \left(\sum_{j=1}^n Q_{ij} \right) \bar{x}_i - \epsilon \sum_{\substack{j=1, \dots, n : \\ Q_{ij} \geq 0}} Q_{ij} - \epsilon \sum_{\substack{j=1, \dots, n : \\ Q_{ij} < 0}} Q_{ij} \\
& = \sum_{i=1}^n \sum_{j=1}^n Q_{ij} \bar{y}_{ij} - \sum_{i=1}^n \left(\sum_{j=1}^n Q_{ij} \right) \bar{x}_i - \epsilon \sum_{\substack{j=1, \dots, n : \\ Q_{ij} < 0}} Q_{ij} \geq \sum_{i=1}^n \sum_{j=1}^n Q_{ij} \bar{y}_{ij} + \sum_{i=1}^n L_i \bar{x}_i.
\end{aligned}$$

Secondly, if we consider the solution $\hat{x} = x - \epsilon e_{\hat{i}}$, it also differs from $\bar{x}$ when $i = \hat{i}$, where we have:

$$\bar{y}_{ij} = \begin{cases} 0 & \text{if } Q_{ij} \geq 0 \\ \frac{1}{2} - \epsilon & \text{if } Q_{ij} < 0 \end{cases} \quad i, j = 1, \dots, n.$$

This leads to

$$\begin{aligned}
& \sum_{i=1}^n \sum_{j=1}^n Q_{ij} \hat{y}_{ij} + \sum_{i=1}^n L_i \hat{x}_i = \sum_{i=1}^n \sum_{j=1}^n Q_{ij} \hat{y}_{ij} - \sum_{i=1}^n \left(\sum_{j=1}^n Q_{ij} \right) \hat{x}_i \\
& = \sum_{i=1}^n \sum_{j=1}^n Q_{ij} \bar{y}_{ij} - \epsilon \sum_{\substack{j=1, \dots, n : \\ Q_{ij} < 0}} Q_{ij} - \sum_{i=1}^n \left(\sum_{j=1}^n Q_{ij} \right) \bar{x}_i + \epsilon \sum_{\substack{j=1, \dots, n : \\ Q_{ij} \geq 0}} Q_{ij} + \epsilon \sum_{\substack{j=1, \dots, n : \\ Q_{ij} < 0}} Q_{ij} \\
& = \sum_{i=1}^n \sum_{j=1}^n Q_{ij} \bar{y}_{ij} - \sum_{i=1}^n \left(\sum_{j=1}^n Q_{ij} \right) \bar{x}_i + \epsilon \sum_{\substack{j=1, \dots, n : \\ Q_{ij} \geq 0}} Q_{ij} \geq \sum_{i=1}^n \sum_{j=1}^n Q_{ij} \bar{y}_{ij} + \sum_{i=1}^n L_i \bar{x}_i.
\end{aligned}$$

In both cases, the objective function value of $(\hat{x}, \hat{y})$ is greater than the one of $(\bar{x}, \bar{y})$, proving that $(\bar{x}, \bar{y})$ is optimal. $\square$

The following theorem introduces an interesting class of BQP instances, where all the LT considered in this paper are equivalent in terms of LP-relaxation:

Theorem 1 *If $K = \emptyset$ and $L_i = -\sum_{j=1}^n Q_{ij}$ ($i = 1, 2, \dots, n$), the optimal solution values of the LP-relaxation of GW (Glover and Woolsey 1974) and G (Glover 1975) coincide and they are both equal to $\frac{1}{2} \sum_{i=1}^n (Q_i^- + L_i)$. Moreover, let $(\bar{x}, \bar{w})$ and $(\tilde{x}, \tilde{y})$ be the optimal solutions of their LP-relaxations, we have $\bar{x}_i = \tilde{x}_i = \frac{1}{2}$ ($i = 1, \dots, n$) and $\tilde{y}$ and $\bar{w}$ are defined accordingly to Proposition 8.*

Proof If $L_i = -\sum_{j=1}^n Q_{ij}$, we can apply Lemma 2 to identify the optimal solution of the LP-relaxation of GW (Glover and Woolsey 1974) and its value. We can hence use Lemma 1 to show that there exists an optimal solution to G (Glover 1975) with $\bar{x} = \tilde{x}$ and with the same objective function value. Finally, we notice that, G (Glover 1975) can not be stronger than

Table 2 Summary of the relations between the strength of the LP-relaxations

LP of GW (Glover and Woolsey 1974)	$\Rightarrow$ (Prop. 7)	LP of G (Glover 1975)
$\Downarrow$ (Prop. 6)		$\Downarrow$ (Prop. 5)
LP of ELF		LP of SS (Sherali and Smith 2007)
	$\Rightarrow$ (see Sherali and Smith 2007)	LP of CPP (Chaovalitwongse et al. 2004)

GW (Glover and Woolsey 1974) (Proposition 7), for this reason, we have that the solution with $\bar{x} = \tilde{x} = \frac{1}{2}$ is also an optimal solution to G (Glover 1975). $\square$

An important class of instances, satisfying Theorem 1, corresponds to BQP reformulation of the Max Cut problem (see Sect. 3.1).

Finally, in Table 2, we summarize the relations between the different LP-relaxations of the formulations studied. The relation “ $A \Leftrightarrow B$ ” stands for “A and B have the same LP-relaxation value” and “ $A \Rightarrow B$ ” stands for “A has a stronger LP-relaxation value than B”. For each relation, we report the reference where it is proved.

3 Computational experiments

In this section, we assess the computational performances of the LTs discussed in this paper, i.e., GW (Glover and Woolsey 1974), G (Glover 1975), SS (Sherali and Smith 2007) and the new ELF. Preliminary experiments showed that the LP-relaxation of CPP (Chaovalitwongse et al. 2004) is extremely weak on all the test-beds, this is due to the fact of having large big-M values compared to the ones of SS (Sherali and Smith 2007). Since this formulation is computationally dominated by SS (Sherali and Smith 2007), it is dropped from the comparison tests.

The first part of our tests is based on the Unconstrained BQP and on the Maximum Cut Problem. The second part of our tests is based on quadratic instances with linear constraints, more precisely we consider the Quadratic Knapsack Problem and the Quadratic Stable Set Problem. In our computational comparison we decided not to use any problem dependent valid inequalities, since our goal is to compare the formulations in their basic versions.

All the experiments have been performed on a computer with a 3.40 Ghz 8-core Intel Core i7-3770 processor and 16Gb RAM, running a 64 bits Linux operating system. We use CPLEX 12.7.0 (ILOG 2017) as MILP solver ran single-threaded with default parameter settings. The tests have been performed with the Constraint-Redundancy elimination activated. It is worth mentioning that, thanks to preliminary experiments, we observed a substantial worsening (up to several orders of magnitude) of the overall computing time of CPLEX when this intuitive adjustment of the LTs is not applied.

MILP solvers implement different branch-and-cut algorithms and they all heuristically separate generic cuts in order to improve the quality of the dual bounds. In addition, the solvers perform several variable-and-constraint reduction techniques which are based on the specific formulation to solve. These aspects must be taken into account when analyzing the computational performance of different LTs. For example a more aggressive strategy of a solver applied to a specific LT (rather than to another one) may influence the computational behavior. Unfortunately these distortions generated by the specific implementations of the

solvers are impossible to eliminate. In other words, the computational analysis of the next section is dependent on the choice of CPLEX 12.7.0 and the conclusions may be slightly different in case another solver is used (or simply other settings of parameters). Nevertheless our experiments give an overall assessment of the relative computational performance of the different LTs for four different classes of problems and using one of the state-of-the-art MILP solvers.

3.1 Unconstrained problems

3.1.1 Problem description

Our first aim is to investigate the strength of the different formulations without the influence of additional constraints ($K = \emptyset$). For this reason, we adopt the Biq Mac library (see Wiecele 2007) as a case study. The Biq Mac library is a collection of instances widely used in the literature, see for example Rendl et al. (2010) or, more recently, Krislock et al. (2014). It is composed of two families of problems: the first one is the *Unconstrained BQP* (UBQP):

$$\min \left\{ \sum_{i=1}^n \sum_{j=1}^n Q_{ij} x_i x_j + \sum_{i=1}^n L_i x_i : x \in \{0, 1\}^n \right\},$$

where Q is a symmetric matrix of order n . The second family is the *Max Cut* (MC) problem. The MC consists of finding a maximum weighted bipartition of a graph G of n vertices [see for example Rendl et al. (2010)] and it can be formulated as follows:

$$\max \left\{ \sum_{i=1}^n \sum_{j=1}^n \tilde{Q}_{ij} x_i x_j : x \in \{-1, 1\}^n \right\},$$

where $\tilde{Q}$ is the Laplacian matrix of the graph G . It is well known that the UBQP and the MC are equivalent [see Caprara (2008) or Krislock et al. (2014)].

The Biq Mac library used for the experiments is divided into five classes: `beasley`, `gka`, and `be` are UBQP instances, while `rudy`, `ising` are MC instances. The classes altogether form a test bed of 343 instances. Some are randomly generated instances and others come from a statistical physics application. In each class, the instances differ in terms of size n , density of the matrix Q and L [for further details on the instance features, we refer the reader to Wiecele (2007)].

In our experiments we used the UBQP formulation of the MC problem obtained with the following transformation. Let w_{ij} ($i, j = 1, \dots, n$, $w_{ij} = w_{ji}$) be the set of weights associated to a given graph G , the UBQP for the MC on G has the following objective function (see Caprara 2008):

$$\begin{aligned} \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n w_{ij} [x_i(1-x_j) + x_j(1-x_i)] &= - \sum_{i=1}^n \sum_{j=1}^n w_{ij} x_i x_j + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n w_{ij} (x_i + x_j) = \\ &- \sum_{i=1}^n \sum_{j=1}^n w_{ij} x_i x_j + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n w_{ij} x_i + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n w_{ij} x_j = - \sum_{i=1}^n \sum_{j=1}^n w_{ij} x_i x_j \\ &+ \sum_{i=1}^n \sum_{j=1}^n w_{ij} x_i. \end{aligned}$$

The last term of the chain of equations shows that all UBQP instances arising from MC instances have $L_i = -\sum_{j=1}^n Q_{ij}$ and hence belong to the class of instances of Theorem 1.

3.1.2 Tables description

In Tables 3 and 4 we report the results concerning the whole Biq Mac Library. Both tables are divided into 5 horizontal blocks, one for each class of instances (`be`, `beasley`, `gka`, `ising` and `rudy`). In each line we group together the results relative to instances of the same size (same values of n). The first two columns of both tables report the size of the subclasses of instances and their cardinality (i.e., the number of instances of a specific subclass).

In Table 3 we report the results concerning the computational behavior for solving the LP-relaxation of the LTs considered. The table provides the following information:

- *LP times* The average time required by CPLEX to solve the LP-relaxation to proven optimality.
- *LP iterations* The average number of simplex iterations need by CPLEX to solve the LP-relaxation.
- *LP gaps* The average gap (in percentage) between the optimal value of the LP-relaxation and the optimal solution value (or best know incumbent solution).

In Table 4 we report the results concerning the computational behavior for solving the test problems with the different LTs to proven optimality. The table provides the following information:

- *MILP times* The average computational time required by CPLEX to solve the problem to proved optimality and the number of instances solved within the time-limit of 1800 seconds (we report `tl` and `0` in case all the instances of a specific subclass and formulation reach the time limit). In each line, the most efficient LT is reported in bold text, i.e., the one with shorter computing time or larger number of instances solved respectively.
- *MILP branching nodes* The average number of nodes explored by CPLEX in the Branch-and-Bound tree.
- *MILP exit gaps* The average percentage exit gap between the upper and the lower bounds computed by CPLEX in case the time limit is reached (`0.0` is reported otherwise). The LT with the lowest open gap is reported in bold text.

Tables discussion Table 3 shows, as expected, that the four LTs considered present the same LP-relaxation values for the MC instances. On the other hand, the situation for the Biq instances is different: GW (Glover and Woolsey 1974) and ELF are characterized by stronger LP-relaxation values (the difference is less clear for the `be` and `beasley` class of instances). As far as LP times and iterations are concerned, GW (Glover and Woolsey 1974) and ELF take a higher number of iterations and longer computing times than G (Glover 1975) and SS (Sherali and Smith 2007). This figure can be explained by the higher number of additional variables and constraints. Since for the MC instances all the LP-relaxation bounds are equivalent, SS (Sherali and Smith 2007) is the more effective option to compute it.

Concerning Table 4, the formulations with the best computational behavior for the Biq Mac instances are GW (Glover and Woolsey 1974) and ELF. These formulations clearly outperform G (Glover 1975) and SS (Sherali and Smith 2007) concerning both the average computing time and the total number of instances solved. For the Biq instances, this is due

Table 3 Computational comparison of the LP-relaxations of the LTs for Biq Mac instances

<i>n</i>	#	LP times		LP iterations				LP gaps					
		GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)				
<i>be</i>	100	10	0.07	0.31	0.01	0.01	4292.1	8831.0	210.5	100.0	73.6	73.6	73.6
	120	20	0.05	0.22	0.01	0.01	3451.2	7033.0	259.7	122.4	63.6	63.7	63.7
	150	20	0.27	0.48	0.02	0.01	4899.6	10,985.4	328.2	151.5	69.1	69.1	69.1
	200	20	0.82	1.50	0.04	0.02	8669.5	19,494.1	437.4	200.6	79.1	79.1	79.1
	250	10	0.07	0.11	0.04	0.03	2798.4	5856.1	569.6	291.5	42.6	43.7	43.7
<i>beasley</i>	50	10	0.00	0.00	0.00	0.00	120.9	234.2	82.0	63.4	0.1	10.2	10.2
	100	10	0.01	0.01	0.00	0.00	676.1	1034.4	218.0	136.5	12.9	22.9	22.9
	250	10	0.07	0.12	0.05	0.03	2822.2	5943.2	598.8	278.8	44.3	44.9	44.9
	500	10	0.64	1.11	0.32	0.19	10,543.1	22,787.4	1184.8	504.1	87.4	87.4	87.4
<i>gka</i>	20	1	0.00	0.00	0.00	0.00	28.0	408.0	20.0	21.0	66.0	67.3	67.3
	30	3	0.00	0.00	0.00	0.00	171.7	561.3	54.3	34.7	40.3	43.1	43.1
	40	2	0.00	0.02	0.00	0.00	337.5	1382.5	62.5	41.0	60.4	61.2	61.2
	50	4	0.00	0.01	0.00	0.00	312.3	1143.0	84.0	58.5	34.3	39.4	39.4
	60	3	0.00	0.01	0.00	0.00	328.7	1738.3	91.3	71.3	38.3	41.7	41.7
	70	3	0.01	0.02	0.00	0.00	390.3	2252.0	122.7	81.7	40.1	45.8	45.8
	80	3	0.00	0.02	0.00	0.00	465.7	2779.0	133.0	98.0	35.1	43.1	43.1
	90	2	0.01	0.04	0.00	0.00	341.0	4443.0	131.5	108.5	45.7	52.7	52.7

Table 3 continued

<i>n</i>	#	LP times				LP iterations				LP gaps			
		GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)
<i>ising</i>													
100	13	0.02	0.10	0.00	0.00	1925.1	4654.2	203.5	109.3	50.7	50.7	53.3	53.3
125	1	0.03	0.15	0.01	0.01	182.0	15,366.0	118.0	121.0	92.2	92.2	92.7	92.7
200	5	0.27	0.60	0.03	0.02	4898.2	10,759.0	431.6	209.6	58.9	58.9	59.7	59.7
500	5	32.52	64.78	0.55	0.35	50,312.4	114,504.4	1153.4	500.4	92.8	92.8	92.8	92.8
<i>ising</i>													
100	9	0.03	0.07	0.01	0.00	3437.1	6506.9	203.7	101.1	20.4	20.4	20.4	20.4
125	3	0.01	0.01	0.00	0.00	510.0	842.7	261.7	127.3	27.2	27.2	27.2	27.2
150	6	0.13	0.29	0.02	0.01	10,253.0	19,422.7	306.8	150.0	19.2	19.2	19.2	19.2
200	6	0.24	0.57	0.03	0.02	15,686.3	29,183.7	409.2	200.0	18.8	18.8	18.8	18.8
216	3	0.01	0.02	0.01	0.01	777.0	1438.3	456.3	217.7	29.1	29.1	29.1	29.1
225	3	0.00	0.02	0.01	0.01	633.0	1082.7	462.0	234.7	16.9	16.9	16.9	16.9
250	6	0.32	0.95	0.05	0.04	20,947.0	39,318.8	512.5	250.0	19.9	19.9	19.9	19.9
300	6	0.56	1.15	0.08	0.07	26,308.3	49,666.8	616.3	300.0	19.5	19.5	19.5	19.5
343	3	0.02	0.04	0.03	0.02	1233.3	2283.3	722.7	346.3	29.7	29.7	29.7	29.7
400	3	0.02	0.04	0.02	0.02	1117.0	1936.7	817.0	415.7	17.9	17.9	17.9	17.9
<i>rudy</i>													
60	10	0.00	0.02	0.00	0.00	618.9	2334.2	63.2	63.2	40.1	40.1	40.1	40.1
80	30	0.01	0.06	0.00	0.00	1372.1	3478.5	136.2	81.0	59.0	59.0	59.0	59.0
100	90	0.02	0.10	0.00	0.00	1916.5	5402.2	161.7	102.6	56.6	56.6	56.6	56.6

Table 4 Computational comparison of the different formulations for solving Biq Mac instances to proven optimality

<i>n</i>	#	MILP times				MILP branching nodes				exit gaps			
		GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)
<i>be</i>													
100	10	tl	0	tl	0	903	159,877	489,570	294,827	33.8	64.7	50.0	51.1
120	20	1730.6	1	tl	0	4400	368,818	286,643	218,557	20.5	50.4	40.0	40.8
150	20	tl	0	tl	0	1667	234,478	229,528	162,299	33.2	60.2	47.3	50.5
200	20	tl	0	tl	0	266	117,224	129,028	207,618	57.2	69.0	54.7	62.1
250	10	tl	0	tl	0	4032	433,728	41,637	42,396	8.5	32.7	25.5	25.3
<i>beasley</i>													
50	10	0.0	10	0.0	10	0	0	74	1	0.0	0.0	0.0	0.0
100	10	0.3	10	11.2	10	5	142	1637	1440	0.0	0.0	0.0	0.0
250	10	tl	0	tl	0	3642	484,278	41,674	47,243	10.1	35.4	27.4	27.5
500	10	tl	0	tl	0	0	75,038	6188	86,809	81.7	58.9	54.4	59.2
<i>gka</i>													
20	1	0.0	1	0.0	1	0	21	44	43	0.0	0.0	0.0	0.0
30	3	0.0	3	0.1	3	0	33	247	109	0.0	0.0	0.0	0.0
40	2	0.7	2	1.8	2	0	363	2222	1324	0.0	0.0	0.0	0.0
50	4	0.6	4	10.2	4	2	771	4607	3771	0.0	0.0	0.0	0.0
60	3	0.5	3	5.2	3	9	375	1153	1460	0.0	0.0	0.0	0.0
70	3	0.8	3	7.1	3	12	266	1927	1731	0.0	0.0	0.0	0.0
80	3	0.5	3	4.9	3	12	160	1466	887	0.0	0.0	0.0	0.0
90	2	0.5	2	1.6	2	30	141	784	532	0.0	0.0	0.0	0.0

Table 4 continued

<i>n</i>	#	MLP times			MLP branching nodes					exit gaps			
		GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)
100	13	987.5	6	1241.4	4	2848	332,199	264,265	208,236	9.9	30.1	24.6	25.4
125	1	3.8	1	6.9	1	176	529	3170	2827	0.0	0.0	0.0	0.0
200	5	1465.3	1	tl	0	1079	211,712	118,335	114,323	24.9	50.1	39.9	44.9
500	5	tl	0	tl	tl	0	19242	5329	130,956	90.1	86.3	75.3	78.0
<i>ising</i>													
100	9	74.2	9	65.6	9	1644.5	1	1699.4	1	5679	10,918	1,943,960	3,449,968
125	3	1.6	3	17.9	3	tl	0	tl	0	38	27,326	3,487,383	4,886,222
150	6	902.3	4	1045.3	3	tl	0	tl	0	16,394	55,880	370,536	1,059,850
200	6	1697.4	1	tl	0	tl	0	tl	0	30,829	98,811	265,229	647,454
216	3	13.6	3	tl	0	tl	0	tl	0	595	1,896,661	3,025,926	2,521,665
225	3	0.8	3	4.9	3	tl	0	tl	0	57	6695	3,973,997	3,392,715
250	6	tl	0	tl	0	tl	0	tl	0	28,752	66,611	202,984	413,173
300	6	tl	0	tl	0	tl	0	tl	0	27,969	58,156	161,895	282,489
343	3	140.4	3	tl	0	tl	0	tl	0	2496	1,124,983	1645263	1550100
400	3	2.4	3	900.7	3	tl	0	tl	0	162	594,238	1,580,827	1,505,487
<i>rudy</i>													
60	10	tl	0	tl	0	tl	0	tl	0	44,880	1,897,854	3,508,202	1,294,517
80	30	1197.4	10	1209.3	10	tl	0	tl	0	4691	503,775	1,005,699	717,697
100	90	1223.8	30	1561.2	20	tl	0	tl	0	3897	676,430	784,450	533,997
											37.5	35.3	33.7

to the fact of having stronger LP-relaxation values which allows a better computational convergence. For the Mac instances, where the LP-relaxation values coincides, the different behavior can be explained by the better performance of GW (Glover and Woolsey 1974) and ELF during the branching scheme and by the efficacy of the generic cuts generated by CPLEX. In case the time limit is reached, ELF explores a higher number of branching nodes than GW (Glover and Woolsey 1974) but the average exit gaps are higher. This figure is due to the different policy of CPLEX in separating cuts at the root node, i.e., for GW (Glover and Woolsey 1974), CPLEX spends a large amount of time in improving the root node bound, while for ELF CPLEX starts immediately to branch.

3.2 Constrained problems

In the remaining part of this computational section we introduce and discuss the other two test problems considered, i.e., the Quadratic Stable Set Problem and the Quadratic Knapsack Problem. These two problems are used to measure the behaviour of the different LTs discussed in this paper in presence of constraints.

3.2.1 Quadratic stable set problem

Problem description We recall that a stable set is a subset of fully disconnected vertices. Following the notation of Sect. 2, we present the problem in minimization form. The formal definition of the Quadratic Stable Set Problem (QSSP) is the following: given an undirected graph $G = (V, E)$, with $V = \{v_1, \dots, v_n\}$ the set of vertices and E the set of edges, a vector of linear cost $L \in \mathbb{R}^n$ on the vertices and a symmetric matrix of quadratic costs $Q \in \mathbb{R}^{n \times n}$ on couples of vertices, the QSSP searches for a stable set of G with minimum cost. In other words, if vertices i and j are in the solution, not only the linear costs are collected but also an additional quadratic cost equal to Q_{ij} is considered.

The mathematical formulation of the QSSP reads as follows:

$$\min \left\{ \sum_{i=1}^n \sum_{j=i+1}^n 2Q_{ij}x_i x_j + \sum_{i=1}^n L_i x_i : x_i + x_j \leq 1, \forall (v_i, v_j) \in E, x \in \{0, 1\}^n \right\}$$

This quadratic counterpart of the Linear Stable Set Problem has not received much attention in the literature [for further details on the QSSP we refer the interested reader to Furini and Traversi (2013)]. As test-bed we used 16 instances introduced in Furini and Traversi (2013), based on random graphs a number of vertices $n = 150$, density $\mu \in \{50\%, 75\%\}$, and a percentage of negative quadratic costs $\nu \in \{25\%, 50\%, 75\%\}$. We used 2 instances for each class using different random seeds.

In the following paragraph we discuss the special case of QSSP with $\nu = 0\%$

A relevant special case As observed in Jaumard et al. (1998), the QSSP with only non-negative costs, i.e., $Q_{ij} \geq 0, v_i, v_j \in V$, can be viewed as a weighted Stable Set Problem (SSP) on an extended graph $\tilde{G} = (\tilde{V}, \tilde{E})$, where $\tilde{V} = V \cup V_1 \cup V_2, \tilde{E} = E \cup E_1 \cup E_2 \cup E_{12}$ and

$$V_1 = \{v_{ij}^i : v_i, v_j \in V, Q_{ij} > 0\},$$

$$V_2 = \{v_{ij}^j : v_i, v_j \in V, Q_{ij} > 0\},$$




Fig. 1 Example: a graph G (left) and the corresponding extended graph $\tilde{G}$ (right)

$$\begin{aligned}
 E_1 &= \{(v_i, v_{ij}^i), v_i \in V, v_{ij}^i \in V_1 : Q_{ij} > 0\}, \\
 E_2 &= \{(v_{ij}^j, v_j), v_j \in V, v_{ij}^j \in V_2 : Q_{ij} > 0\}, \\
 E_{12} &= \{(v_{ij}^i, v_{ij}^j), v_{ij}^i \in V_1, v_{ij}^j \in V_2 : Q_{ij} > 0\}.
 \end{aligned}$$

The cost of nodes $v_{ij}^i \in V_1$ and $v_{ij}^j \in V_2$ are equal to $-Q_{ij}$, replacing in this way the initial quadratic costs Q . The optimal solution value of QSSP on G is then equal to the optimal solution value of the weighted SSP on $\tilde{G}$ plus the constant $\sum_{i=1}^n \sum_{j=1}^n Q_{ij}$.

An example of the extended graph is given in Fig. 1. On the left part of the figure we report a graph $G = (V, E)$ with 6 vertices and 9 edges and the following costs:

$$Q = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 0 & 1 & 2 \\ 0 & 3 & 0 & 0 & 4 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 4 & 0 & 0 & 1 \\ 0 & 2 & 1 & 0 & 1 & 0 \end{bmatrix}; L = [-15 \ -15 \ -2 \ -10 \ -8 \ -8].$$

The dashed lines represent a positive entry Q_{ij} . On the right part of the figure we report the corresponding extended graph $\tilde{G} = (\tilde{V}, \tilde{E})$ with 18 vertices and 25 edges. In both graph, the optimal solutions are represented by the vertices in red. The cost of the stable set $\{v_2, v_5, v_6\}$ in G is equal to -23 , while the cost of the stable set $\{v_2, v_5, v_6, v_{23}^3, v_{35}^3, v_{36}^3\}$ in $\tilde{G}$ is equal to -47 . Finally, the constant part is equal to 24.

In other words, for the QSSP with $Q_{ij} \geq 0$, ELF introduces only the sets of constraints (16), (17) and (18) and, hence, it is equivalent to the standard ILP formulation with edge constraints for the Weighted Stable Set Problem on the extended graph $\tilde{G}$.

Table description In Table 5 we report the results concerning the computational behaviour of the four LTs studied on the QSSP benchmark. In each line, we group together the average results relative to instances with same values of μ and ν . Each entry of the table provides the

Table 5 Computational comparison of the different formulations for solving QSSP instances

n	μ	ν	#	MILP times				MILP branching nodes			
				GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)
150	50	25	2	139.68	340.94	216.82	334.01	1613.50	3143.00	198,666.50	126,996.00
		50	2	1216.65	540.30	291.52	366.40	3915.50	10,166.50	337,011.00	171,489.00
	75	2	1490.78	626.17	314.91	362.90	14,382.00	15,674.50	394,338.50	169,204.00	
		25	2	37.59	52.26	19.29	44.41	149.50	139.00	6277.00	7910.00
	75	50	2	85.14	84.61	20.70	44.03	260.50	264.00	9651.00	7731.50
		75	2	98.99	119.84	23.99	43.58	832.50	835.50	13,440.50	10,856.00
n	μ	ν	#	LP times				LP gaps			
				GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)
150	50	25	2	0.05	0.49	0.07	0.08	98.14	98.14	99.08	99.08
		50	2	0.06	0.86	0.02	0.06	98.06	98.06	99.01	99.01
	75	2	0.07	1.28	0.01	0.02	98.00	98.00	98.98	98.98	
		25	2	0.01	0.41	0.01	0.12	97.47	97.47	99.41	99.41
	75	50	2	0.01	0.59	0.01	0.08	97.68	97.68	99.42	99.42
		75	2	0.01	0.05	0.01	0.03	98.41	98.41	99.59	99.59

aggregated results on 2 instances ($\# = 2$ on the table). We provide, for each LT, the following information:

- *MILP times* The average computational time required by CPLEX to solve the problem to proved optimality.
- *MILP branching nodes* The average number of nodes in the Branch-and-Bound tree explored by CPLEX.
- *LP times* The average time required by CPLEX to solve the LP-relaxation to proven optimality.
- *LP gaps* The average percentage gap between the optimal value of the LP-relaxation and the optimal solution.

Table discussion The formulations with the best computational behavior for the QSSP instances are G (Glover 1975) and SS (Sherali and Smith 2007) on average. ELF presents on average a better computing time of GW (Glover and Woolsey 1974). The number of MILP branching nodes of ELF is slightly greater than GW (Glover and Woolsey 1974). G (Glover 1975) and SS (Sherali and Smith 2007) are characterized by a higher number of nodes. This fact is not surprising since those formulations have a weaker LP-relaxation compared to GW (Glover and Woolsey 1974) and ELF. However, the weakness of G (Glover 1975) and SS (Sherali and Smith 2007) is counterbalanced by their smaller size in terms of variables and constraints (See Table 1) which leads to a faster node solution time. Finally, it is interesting to notice that the more difficult instances are the one with density $\mu = 50\%$ and percentage of negative costs $\nu \in \{50\%, 75\%\}$.

3.2.2 Quadratic knapsack problem

Problem description The Quadratic Knapsack Problem (QKP) consists in maximizing a quadratic function with positive coefficients subject to a linear capacity constraint. The mathematical formulation reads as follows:

$$\max \left\{ \sum_{i=1}^n \sum_{j=i+1}^n 2Q_{ij}x_i x_j + \sum_{i=1}^n L_i x_i : \sum_{i=1}^n c_i x_i \leq C, x \in \{0, 1\}^n \right\},$$

where c_i ($i = 1, \dots, n$) are integer non negative coefficients representing the size of the items and C is the capacity of the knapsack. In addition to the classical knapsack, where only linear profits L_i are considered, here we maximize also the profits associated to couples of objects (their profit is represented by Q_{ij}). For further information on the problem, see Caprara et al. (1999) and Pisinger (2007). In our computational tests we used randomly generated instances taken from Billionnet and Soutif (2004) with $n = 100$ and density $\delta = 25\%, 50\%, 75\%, 100\%$, i.e., δ is the number of non-zero coefficients of the objective function divided by $n(n-1)/2$.

Table description In Table 6 we report the results concerning the computational behaviour for solving the test problems to proven optimality. Table 6 presents the same information of Table 5. Each line presents the aggregated results concerning 10 instances with same size and δ .

Table 6 Computational comparison of the different formulations for solving QKP instances

n	δ	#	MILP times		MILP branching nodes					
			GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)
100	25	10	0.54	0.51	0.98	0.86	188.50	188.10	1004.80	717.60
	50	10	4.40	4.06	2.76	2.78	438.50	405.10	2303.50	1978.10
	75	10	39.39	42.23	19.96	59.65	3122.20	4212.70	9549.00	18980.50
	100	10	113.82	145.09	5.87	6.73	5098.44	4918.67	7570.89	5816.89
n	δ	#	LP times		LP gaps					
			GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)	GW (Glover and Woolsey 1974)	ELF	G (Glover 1975)	SS (Sherali and Smith 2007)
100	25	10	0.03	0.03	0.00	0.00	1.71	1.71	12.41	12.41
	50	10	0.08	0.11	0.00	0.00	1.51	1.51	11.15	11.15
	75	10	0.15	0.27	0.00	0.01	4.48	4.48	13.53	13.53
	100	10	0.37	0.50	0.01	0.00	1.19	1.19	11.93	11.93

Table discussion It is interesting to notice that, when the density δ is high, the number of MILP branching nodes is comparable for all the formulations considered. Apart from this feature, the computational behaviour for the QKP of the formulations studied is in line with the one obtained for the QSSP, showing that G (Glover 1975) and SS (Sherali and Smith 2007) are the formulations with the better computational behaviour.

4 Conclusions

In this paper we compared several LTs for binary quadratic problems present in the literature, called here GW (Glover and Woolsey 1974), G (Glover 1975) and SS (Sherali and Smith 2007) with a new formulation called ELF. We showed the equivalence of the LP-relaxation values of G (Glover 1975), SS (Sherali and Smith 2007), GW (Glover and Woolsey 1974) and ELF when dealing with a specific class of Unconstrained binary quadratic problems. Among the formulations studied, GW (Glover and Woolsey 1974) and ELF provide the better performances in practice when applied to unconstrained problems while G (Glover 1975) and, to a lesser extent, SS (Sherali and Smith 2007) perform better on the special classes of constrained problems that have been tested in the paper. Whether this observation generalizes to other types of constrained problems is an interesting question that may be worth further investigations. The new proposed formulation ELF performs relatively well in all classes of instances. Even if GW (Glover and Woolsey 1974) and ELF present the same LP-relaxation, their behaviour in practice can be different according to the problem considered: GW (Glover and Woolsey 1974) is slightly preferable over ELF when dealing with Biq Mac instances, while the contrary happens when solving QSSP instances. For the two considered problems with constraints, the trade off between the strength of the LP-relaxation and the speed in the solution of the MILP branching nodes plays in favour of G (Glover 1975) and SS (Sherali and Smith 2007). Finally, we recall that the selection of the best LT depends also on the current performance and engineering of the MILP solvers.

References

- Adams, W. P., Forrester, R. J., & Glover, F. W. (2004). Comparisons and enhancement strategies for linearizing mixed 0–1 quadratic programs. *Discrete Optimization*, 1(2), 99–120.
- Adams, W. P., & Sherali, H. D. (1986). A tight linearization and an algorithm for zero-one quadratic programming problems. *Management Science*, 32(10), 1274–1290.
- Billionnet, A., & Soutif, E. (2004). An exact method based on lagrangian decomposition for the 01 quadratic knapsack problem. *European Journal of Operational Research*, 157(3), 565–575.
- Caprara, A. (2008). Constrained 0–1 quadratic programming: Basic approaches and extensions. *European Journal of Operational Research*, 187, 1494–1503.
- Caprara, A., Pisinger, D., & Toth, P. (1999). Exact solution of the quadratic knapsack problem. *INFORMS Journal on Computing*, 11(2), 125–137.
- Chaovalitwongse, W., Pardalos, P. M., & Prokopyev, O. A. (2004). A new linearization technique for multi-quadratic 01 programming problems. *Operations Research Letters*, 32(6), 517–522.
- Conforti, M., Cornuéjols, G., & Zambelli, G. (2010). Extended formulations in combinatorial optimization. *4OR*, 8(1), 1–48.
- Forrester, R., & Greenberg, H. (2008). Quadratic binary programming models in computational biology. *Algorithmic Operations Research*, 3(2), 110–129.
- Fortet, R. (1960). L'algebre de boole et ses applications en recherche operationnelle. *Trabajos de Estadística*, 11(2), 111–118.

-
- Furini, F., & Traversi, E. (2013). Hybrid SDP bounding procedure. *Lecture Notes in Computer Science*, 7933, 248–259.
- Glover, F. (1975). Improved linear integer programming formulations of nonlinear integer programs. *Management Science*, 22(4), 455–460.
- Glover, F., & Woolsey, E. (1973). Further reduction of zero-one polynomial programming problems to zero-one linear programming problems. *Operations Research*, 21(1), 156–161.
- Glover, F., & Woolsey, E. (1974). Converting the 0–1 polynomial programming problem to a 0–1 linear program. *Operations Research*, 22(1), 180–182.
- Gueye, S., & Michelon, P. (2009). A linearization framework for unconstrained quadratic (0–1) problems. *Discrete Applied Mathematics*, 157(6), 1255–1266.
- Hansen, P., & Meyer, C. (2009). Improved compact linearizations for the unconstrained quadratic 01 minimization problem. *Discrete Applied Mathematics*, 157(6), 1267–1290.
- ILOG IBM. (2017). Cplex optimizer.
- Jaumard, B., Marcotte, O., & Meyer, C. (1998). *Estimation of the Quality of Cellular Networks Using Column Generation Techniques*. Cahiers du GÉRAD. Groupe d'études et de recherche en analyse des décisions.
- Krislock, N., Malick, J., & Roupin, F. (2014). Improved semidefinite bounding procedure for solving Max-Cut problems to optimality. *Mathematical Programming*, 143(1), 61–86.
- Lodi, A. (2010). Mixed integer programming computation. In M. Jünger, T. M. Liebling, D. Naddef, G. L. Nemhauser, W. R. Pulleyblank, G. Reinelt, G. Rinaldi, & L. A. Wolsey (Eds.), *50 Years of integer programming 1958–2008* (pp. 619–645). Berlin: Springer.
- Padberg, M. (1989). The boolean quadric polytope: Some characteristics, facets and relatives. *Mathematical Programming*, 45(1), 139–172.
- Pisinger, D. (2007). The quadratic knapsack problem: A survey. *Discrete Applied Mathematics*, 155(5), 623–648.
- Rendl, F., Rinaldi, G., & Wiegele, A. (2010). Solving max-cut to optimality by intersecting semidefinite and polyhedral relaxations. *Mathematical Programming*, 121(2), 307–335.
- Sherali, H. D., & Adams, W. P. (1998). *A reformulation-linearization technique for solving discrete and continuous nonconvex problems*. Berlin: Springer.
- Sherali, H. D., & Smith, J. C. (2007). An improved linearization strategy for zero-one quadratic programming problems. *Optimization Letters*, 1(1), 33–47.
- Wang, H., Kochenberger, G., & Glover, F. (2012). A computational study on the quadratic knapsack problem with multiple constraints. *Computers & Operations Research*, 39(1), 3–11.
- Wiegele, A. (2007). Biq mac library—A collection of max-cut and quadratic 01 programming instances of medium size. *Technical report, Alpen-Adria-Universität Klagenfurt, Austria*.