

Site-stratified association between the Mediterranean diet intervention and the study event

Abstract

Background: A dietary intervention study on fasting glucose. **Methods:** The association between the Mediterranean diet intervention (exposed = mediterranean) and the study event was summarised with the crude odds ratio and with the Mantel-Haenszel odds ratio pooled across the four study sites (2x2 tables ordered exposed/unexposed by event/no event), with its 95% confidence interval, and the Mantel-Haenszel chi-square test with continuity correction. **Results:** see the Results section. **Conclusion:** the findings are discussed in the context of the analysis sample.

Introduction

We report a dietary intervention study on fasting glucose. The primary question concerns the Mediterranean diet intervention and fasting glucose (mg/dL) at follow-up. The accompanying dataset (data.csv) contains one row per participant-record as exported from the study database, including some duplicated exports and sentinel-coded missing values; it is described in the data dictionary.

Methods

Participants and data. The export contains 333 records. Variables are listed in the data dictionary. Group membership is recorded in `diet`.

Data cleaning and exclusions. Exact duplicate records (identical on every column) were removed, keeping the first occurrence. The value -999 denotes a missing measurement and was treated as missing. Only participants aged 18 to 80 years inclusive were analysed. Group labels in `diet` were entered free-text and were normalised by trimming whitespace and lower-casing before use.

Statistical analysis. The association between the Mediterranean diet intervention (exposed = mediterranean) and the study event was summarised with the crude odds ratio and with the Mantel-Haenszel odds ratio pooled across the four study sites (2x2 tables ordered exposed/unexposed by event/no event), with its 95% confidence interval, and the Mantel-Haenszel chi-square test with continuity correction. The site-specific odds ratio for site 1 is also reported. All analyses used the cleaned analysis sample; no random resampling was used.

Results

Among 313 participants (167 exposed) the crude odds ratio was 0.41 and the Mantel-Haenszel odds ratio pooled over sites was 0.39 (95% CI 0.24 to 0.64; MH chi-square = 13.16, $p < 0.001$). Within site 1 the odds ratio was 0.24.

Discussion

The analysis followed the pre-specified plan. Limitations include reliance on database exports and complete-case handling of missing values.