

Time to the study event by caffeine administration: a survival analysis

Abstract

Background: A laboratory study of caffeine and attention. Methods: Follow-up time in days was computed from the enrolment date to the date of event or last contact. Results: see the Results section. Conclusion: the findings are discussed in the context of the analysis sample.

Introduction

We report a laboratory study of caffeine and attention. The primary question concerns caffeine administration and reaction time (ms) on the Stroop task. The accompanying dataset (data.csv) contains one row per participant-record as exported from the study database, including some duplicated exports and sentinel-coded missing values; it is described in the data dictionary.

Methods

Participants and data. The export contains 225 records. Variables are listed in the data dictionary. Group membership is recorded in `condition`.

Data cleaning and exclusions. Exact duplicate records (identical on every column) were removed, keeping the first occurrence. The value -999 denotes a missing measurement and was treated as missing. Only participants aged 18 to 80 years inclusive were analysed. Group labels in `condition` were entered free-text and were normalised by trimming whitespace and lower-casing before use.

Statistical analysis. Follow-up time in days was computed from the enrolment date to the date of event or last contact. Follow-up was administratively censored at 730 days: participants with longer follow-up were censored at 730 days and counted as event-free, and records with zero follow-up were excluded. The hazard of the study event was modelled with a Cox proportional-hazards model with caffeine administration (1 = caffeine, 0 = placebo) and age (years) as covariates, using the Breslow method for ties. Hazard ratios with Wald 95% confidence intervals ($z = 1.96$) are reported. All analyses used the cleaned analysis sample; no random resampling was used.

Results

Of 207 participants (median follow-up 484.00 days), 88 had an event. The exposure hazard ratio was 0.47 (95% CI 0.30 to 0.72, $p < 0.001$); the hazard ratio per year of age was 1.03.

Discussion

The analysis followed the pre-specified plan. Limitations include reliance on database exports and complete-case handling of missing values.