

Fasting glucose across baseline tertiles

Abstract

Background: A dietary intervention study on fasting glucose. Methods: Participants were divided into three groups (low, middle, high) by tertiles of the baseline value, using the first and second tertile cut-points of the analysis sample (values equal to a cut-point were assigned to the lower group). Results: see the Results section. Conclusion: the findings are discussed in the context of the analysis sample.

Introduction

We report a dietary intervention study on fasting glucose. The primary question was whether the Mediterranean diet intervention is associated with fasting glucose (mg/dL) at follow-up. The dataset accompanying this report (data.csv) contains one row per participant and is described in the data dictionary.

Methods

Participants and data. Data were collected for 258 participants. Each record contains the variables listed in the data dictionary. Group membership is recorded in the column `diet`.

Exclusion criteria and preprocessing. Only participants aged 18 to 80 years inclusive were included in the analysis. Participants with missing values for glucose_fu were excluded (complete-case analysis). After the exclusions above, observations whose fasting glucose (mg/dL) at follow-up lay more than three sample standard deviations from the overall mean were treated as outliers and removed.

Statistical analysis. Participants were divided into three groups (low, middle, high) by tertiles of the baseline value, using the first and second tertile cut-points of the analysis sample (values equal to a cut-point were assigned to the lower group). Mean fasting glucose (mg/dL) at follow-up was compared across the three groups with a one-way analysis of variance; eta squared was computed as the between-group sum of squares divided by the total sum of squares. All analyses were performed on the analysis sample defined above. Statistics were computed with standard scientific software; no random resampling was used.

Results

The low, middle and high tertile groups comprised 75, 76 and 73 participants, with mean fasting glucose (mg/dL) at follow-up of 97.11, 99.62 and 108.95 respectively ($F = 18.84$, $p < 0.001$, $\eta^2 = 0.146$).

Discussion

The analysis followed the pre-specified plan described in the Methods. Limitations include the observational nature of some comparisons and the exclusion of records with missing values.