

# Type Ia Supernova Remnants, Circumstellar Interaction, and Supernova Progenitors

**Carles Badenes, Travis Court**  
(University of Pittsburgh/PITT PACC)  
**Herman Lee** (U. Kyoto)  
**Dan Patnaude.**(SAO/CfA)



University of  
**Pittsburgh**



# This Talk in a Nutshell

Carles Badenes  
EAS 06/28/22

## X-ray spectra of SNRs constrain SN Ia explosion physics & progenitor properties:

- $^{56}\text{Ni}$  mass  $\Leftrightarrow$  brightness [Badenes+ 06,08; Krause+ 08; Rest+ 08].
- WD mass  $\Leftrightarrow$  Ch/sub-Ch explosions [Yamaguchi+ 15].
- Pre-SN mass loss rate  $\Leftrightarrow$  WD accretion mode [Badenes+ 07, Williams+ 11, Yamaguchi+ 14].


**We are working on a new model grid to map the parameter space for CSM interaction in Type Ia SNRs**



## X-ray spectra of SNRs constrain SN Ia explosion physics & progenitor properties:

- $^{56}\text{Ni}$  mass  $\Leftrightarrow$  brightness [Badenes+ 06,08; Krause+ 08; Rest+ 08].
- WD mass  $\Leftrightarrow$  Ch/sub-Ch explosions [Yamaguchi+ 15].
- Pre-SN mass loss rate  $\Leftrightarrow$  WD accretion mode [Badenes+ 07, Williams+ 11, Yamaguchi+ 14].

**We are working on a new model grid to map the parameter space for CSM interaction in Type Ia SNRs**




# This Talk in a Nutshell

Carles Badenes  
EAS 06/28/22

## X-ray spectra of SNRs constrain SN Ia explosion physics & progenitor properties:

- $^{56}\text{Ni}$  mass  $\Leftrightarrow$  brightness [Badenes+ 06,08; Krause+ 08; Rest+ 08].
- WD mass  $\Leftrightarrow$  Ch/sub-Ch explosions [Yamaguchi+ 15].
- Pre-SN mass loss rate  $\Leftrightarrow$  WD accretion mode [Badenes+ 07, Williams+ 11, Yamaguchi+ 14].

**We are working on a new model grid to map the parameter space for CSM interaction in Type Ia SNRs**




Yamaguchi+ 14

# Circumstellar Interaction in SNRs

Carles Badenes  
EAS 06/28/22

- **AM structure**  $\Rightarrow$  progenitor mass-loss. Modified circumstellar medium (CSM) vs. undisturbed interstellar medium (ISM)
- **SNe**  $\Rightarrow$  Follow-up (radio/X-ray) probes to  $\sim 100$  AU.
- **SNRs**  $\Rightarrow$  **spatial (and temporal) scales relevant for stellar evolution** of SN progenitors ( $t \lesssim \tau_{KH}$ ). **Probe dynamical interaction!**







# Circumstellar Interaction in SNRs

Carles Badenes  
EAS 06/28/22



- **X-ray spectra  $\Rightarrow$  constrain AM structure.** NEI plasma: ionization timescale ( $n_e t$ ) [Badenes+ 07].
- **High  $n_e t \Rightarrow$  high centroid energy and line flux.**



# CSM Interaction in SNRs: Fe K


Carles Badenes  
EAS 06/28/22

- Use **Fe K $\alpha$  line blend** at  $\sim 6.5$  keV as an AM density diagnostic.
- All SNe (Ia and CC) eject some Fe  $\Rightarrow$  innermost layers.
- Large  $n_e t$  required to fully ionize Fe  $\Rightarrow$  **large dynamic range in  $\rho_{AM}$** .
- **Bulk properties** ( $n_e t \Leftrightarrow$  line centroids, ages, radii)  $\Leftrightarrow$  progenitor mass loss.



# SNRs in Bulk


- Require Fe K $\alpha$  centroid: **25 SNRs** (*Suzaku*, *Chandra*, *XMM*) [Yamaguchi+ 14, Borkowski+ 13, Maggi+ 16, M-R+ 18].
- **Bulk properties:** Fe K $\alpha$  centroid, Fe K $\alpha$  luminosity, radius, age.
- A pattern emerges: **Ia SNRs below  $\sim 6.55$  keV (Fe<sup>+21</sup>), CC SNRs above** [Yamaguchi+ 14, but see Maggi & Acero 17].
- **CC/Ia SNRs have similar ages/radii, but very different  $n_e t \Leftrightarrow$  different AM densities!**









# Models

- **HD+NEI Models required** to draw quantitative conclusions on progenitor mass loss!



# Models: Uniform AM


- **HD+NEI Models required** to draw quantitative conclusions on progenitor mass loss!
- **Uniform AM models work remarkably well for most SN Ia** [Badenes+07, Yamaguchi+ 14, M-R+ 18].



# Models: Uniform AM





- **HD+NEI Models required** to draw quantitative conclusions on progenitor mass loss!
- **Uniform AM models work remarkably well for most SN Ia** [Badenes+07, Yamaguchi+ 14, M-R+ 18].
- Exceptions: N103B, RCW 86, Kepler, ... [Badenes+ 07, Patnaude+ 12, Williams+ 11,14, Broersen+ 14].

**Most Type Ia SNRs show no evidence for strongly modified AM on ~pc scales**



# Models: Outflows



- **Fast outflows:** large cavities, consistent with RCW 86 [Badenes+ 07, Williams+ 11, Broersen+ 14].
- **Slow outflows:** dense CSM, ruled out [Badenes+ 07, Patnaude & Badenes 17].
- Intermediate cases?





# Models: CE cocoons

- **Post-CE PN models** [García-Segura+ 18].
- Models with short delays (SN  $\lesssim 1000$  yr after CE) do not work. Longer delays might be OK.






# Looking Forward

Carles Badenes  
EAS 06/28/22

- **Explore parameter space for AM interaction in Type Ia SNRs** [Travis Court PhD, in progress].
- **Part of a large effort:** CR-modified dynamics [Lee+ 14], CC SNR models. [Patnaude+15,17, Jacovich+ 21].
- **XRISM:** velocity vs.  $n_e t$  for Tycho and SN1006  $\Leftrightarrow$  discriminate Ch and sub-Ch explosions.




# Looking Forward

Carles Badenes  
EAS 06/28/22


- **Explore parameter space for AM interaction in Type Ia SNRs** [Travis Court PhD, in progress].

- **Part of a large effort:** CR-modified dynamics [Lee+ 14], CC SNR models. [Patnaude+15,17, Jacovich+ 21].

- **XRISM:** velocity vs.  $n_e t$  for Tycho and SN1006  $\Leftrightarrow$  discriminate Ch and sub-Ch explosions.





Age: 1000 yr  
Element: Si



# Extra Plots


Carles Badenes  
EAS 06/28/22






# Extra Plots

Carles Badenes  
EAS 06/28/22



# Extra Plots


Carles Badenes  
EAS 06/28/22








# Extra Plots

Carles Badenes  
EAS 06/28/22



# Extra Plots


Carles Badenes  
EAS 06/28/22



# A Step Back

Carles Badenes  
EAS 06/28/22

- **SN Ia AM density estimates** from radio/X-ray SNe ( $\sim 10$  d,  $\sim 0.01$  pc) and SNRs ( $\sim 500$  yr,  $\sim$  several pc) **are consistent with the warm phase of the ISM** [Badenes+ 07, Chomiuk+ 12, 16, Perez-Torres+ 14, Raymond+ 07, Slane+ 14, Borkowski+ 14].  $\Rightarrow$  **'clean' mergers (DD)?**
- **Mild CSM interaction allowed**, maybe small ( $\sim 0.5$  pc) cavities [Patnaude+ 12, Slane+ 14], but not large ones (except for RCW86!).



- **SN Ia AM density estimates** from radio/X-ray SNe ( $\sim 10$  d,  $\sim 0.01$  pc) and SNRs ( $\sim 500$  yr,  $\sim$ several pc) **are consistent with the warm phase of the ISM** [Badenes+ 07, Chomiuk+ 12, 16, Perez-Torres+ 14, Raymond+ 07, Slane+ 14, Borkowski+ 14].  $\Rightarrow$  **'clean' mergers (DD)?**
- **Mild CSM interaction allowed**, maybe small ( $\sim 0.5$  pc) cavities [Patnaude+ 12, Slane+ 14], but not large ones (except for RCW86!).

