




MODELING THE PATH FROM MASSIVE STARS TO SNe AND SNRs: RECENT ADVANCES AND FUTURE PROSPECTS

Salvatore Orlando



INAF - Osservatorio Astronomico di Palermo, Italy



F. Bocchino, B. Olmi, O. Petruk, S. Ustamujic, A. Tutone	(INAF – OAPA, Italy)
M. Miceli, G. Peres, R. Giuffrida, V. Sapienza	(Palermo University, Italy)
H.-T. Janka, A. Wongwathanarat	(Max-Planck-Institut für Astrophysik, Germany)
S. Nagataki, M. Ono, A. Dohi, G. Ferrand	(ABBL/RIKEN, Japan)
E. Greco, J. Vink	(Anton Pannekoek Inst. for Astronomy, Univ. of Amsterdam, NL)
H. Umeda	(Tokyo University, Japan)
M.A. Aloy, P. Mimica	(Valencia University, Spain)
D. Milisavljevic	(Dep. Physics and Astronomy, Purdue University, USA)
D. Patnaude	(Smithsonian Astrophysical Observatory, USA)
D. Burrows	(Penn State University, USA)
S. Park, A. Pazhayath Ravi	(Stanford University, USA)





Supernova Remnants




*Information about progenitor - SN - CR
encoded in the observations*

- probe the physics of SN engines
- investigate the final stages of stellar evolution
- study the process of particle acceleration




- pristine structures and features of progenitor SN




Wongwathanarat+ 2017

- interaction of SN blast with the CSM




Orlando+ 2015

- cosmic rays acceleration



Caprioli 2015



**SNRs are sources of information on the parent SNe
and the progenitor stars**

the progenitor – SN – SNR connection




**an essential step to open new exploring windows on
the physics of SNe and SNRs**


How to link progenitor – SN – SNR ?



- **Multi-physics**




- **Multi-scale**



- **Multi-dimensions**



How to link progenitor – SN – SNR ? The strategy



A unique laboratory to study the star-SN-SNR connection



Cassiopeia A


Observations suggest that its morphology and expansion rate are consistent with a remnant expanding through the wind of the progenitor star
 (e.g. Lee+ 2014)


The bulk of asymmetries observed in Cas A is intrinsic to the explosion

This remnant is one of the best studied and its 3D structure has been characterized in good detail

(e.g. DeLaney+ 2010,
 Milisavljevic & Fesen 2013, 2015,
 Holland-Ashford+ 2019,



- 3 Fe-rich regions
- 2 Si-rich jets
- Rings circling Fe-rich regions







Effects of the SN anisotropies

Cas A: Spatial Distribution of the ejecta



(Orlando+ 2016)



(Milisavljevic & Fesen 2013)

link the main asymmetries and geometry of Cas A's bulk ejecta to the physical characteristics of anisotropies developed soon after the SN explosion








average physical characteristics of post-explosion anisotropies that are able to reproduce the observed Fe-rich regions and Si-rich jets

Constrain energy and masses of post-explosion asymmetries

Cas A: effects of anisotropies in the SN



3D simulations of a neutrino-driven SN explosion reproducing basic properties of Cas A
 (Wongwathanarat+ 2017)



Initial condition:
 after core bounce

Final time:
 ~ 1 day after explosion

Three pronounced iron-rich fingers that may correspond to the extended iron-rich regions observed in Cas A

Major asymmetries observed in Cas A explained by a neutrino-driven explosion
 No need to invoke rapid rotation or jet-driven explosion


The Ni-bubble effect in the transition SN - SNR



3D simulations of a neutrino-driven SN explosion

Asymmetrically expanding ejecta from shock breakout to homologous expansion phase (roughly one year)

(Gabler et al. 2021)



Ni-bubble effect

inflation of the ejecta caused by the heating due to the radioactive decay chain $^{56}\text{Ni} \rightarrow ^{56}\text{Co} \rightarrow ^{56}\text{Fe}$


- acceleration of bulk of Ni
- inflation of initially overdense Ni-rich clumps
- underdense, extended fingers, enveloped by overdense skins of compressed surrounding matter

The remnant of a neutrino-driven CC SN

Self-consistent description of the whole 3D evolution of a neutrino-driven SN explosion, from the CC to the SNR at the age of 2000 years


identify the geometric and physical properties of the post-explosion anisotropies responsible for the morphology of Cas A

main asymmetries and features explained by the interaction of the reverse shock with the initial large-scale asymmetries






(Orlando et al. 2021)

stochastic processes (e.g., convective overturn and the standing accretion shock instability; SASI) that originate during the first seconds of the SN blast




Effects of the inhomogeneous CSM

The forward and reverse shock dynamics of Cas A






(Gotthelf et al. 2001)




(Delaney 2004)

most of the X-ray synchrotron emission originates from the western part of the reverse shock (Vink 2020)



Cas A reveals past interaction with circumstellar shell



Hints on the mass-loss history of the progenitor star and its nature



Cas A reveals past interaction with circumstellar shell



Interaction with a dense asymmetric circumstellar shell

FS interaction: 185 yr (a.d. 1850-1860)

Charles Darwin publishes The Origin of Species (1859)


RS interaction: 296 yr (a.d. 1940-1946)

World War II (1940-45)

Most likely massive eruption from the progenitor star that occurred about 10^4 - 10^5 years prior to core-collapse

Total mass of the shell M_{sh} estimated between 2 and $3 M_{\odot}$


Reverse shock velocity in the ejecta reference frame





geometric center of the
from the geometric cent

of the X-ray synchrotron
es from the western part
ock

SN explosions in the CSM of massive stars




(Meyer et al. 2021)



- Runaway RSG
- Episodic mass eruption
- Wolf-Rayet star

Take away points

- **SNRs morphology and properties reflect**
 - Interaction with the inhomogeneous ambient environment
 - Asymmetries and anisotropies inherited from the parent SNe
 - Structure of the progenitor star at collapse
- **Deciphering multi- λ observations of SNRs crucial to extract information about**
 - CSM; mass loss history of the progenitor star
 - complex phases after the core-collapse; SN engine
 - nature of the progenitor stellar system



HOW


3D HD/MHD models can help in linking SNRs to their parent SNe and progenitor stars

the progenitor – SN – SNR connection
has breakthrough potential to open new exploring windows on the physics of massive stars, SNe and SNRs

THE CHALLENGE

Deciphering observations might critically depend on the models

- They should connect ***self-consistently*** stellar progenitor → SN → SNR
 - multi-physics, multi-scale, multi-dimension (progenitor, SN, SNR)
- They should be based on ***solid observational facts***
 - account for dynamics, energetics, and spectral properties of SNe and SNRs



Model - Observations