

## Forum Review

# Peroxidases of Trypanosomatids

Helena Castro<sup>1</sup> and Ana M. Tomás<sup>1,2</sup>

### Abstract

This article provides an overview about the recent advances in the dissection of the peroxide metabolism of Trypanosomatidae. This family of protozoan organisms comprises the medically relevant parasites *Trypanosoma brucei*, *Trypanosoma cruzi*, and *Leishmania* spp. Over the past 10 years, three major families of peroxidases have been identified in these organisms: (a) 2-cysteine peroxiredoxins, (b) nonselenium glutathione peroxidases, and (c) ascorbate peroxidases. In trypanosomatids, these enzymes display the unique feature of using reducing equivalents derived from trypanothione, a dithiol found exclusively in these protozoa. The electron transfer between trypanothione and the peroxidases is mediated by a redox shuttle, which can either be tryparedoxin, ascorbate, or even glutathione. The preference for the intermediate molecule differs among each peroxidase and so does the specificity for the peroxide substrate. These observations, added to the fact that these peroxidases are distributed throughout different subcellular compartments, point to the existence of an elaborate peroxide metabolism in trypanosomatids. With the completion of the trypanosomatids genome, other molecules displaying peroxidase activity might be added to this list in the future. *Antioxid. Redox Signal.* 10, 000–000.

### Trypanosomatids

KINETOPLASTIDA ARE PRIMITIVE EUKARYOTIC ORGANISMS that parasitize animals and plants, some of which are relevant due to the diseases they cause and the economic losses thereof associated. Among the Order Kinetoplastida, the Trypanosomatidae family acquires particular significance because it includes the human parasites *Trypanosoma brucei*, *Trypanosoma cruzi*, and *Leishmania* spp. These are the causative agents of sleeping sickness, Chagas' disease, and the three manifestations of leishmaniasis, which together cause 112,000 deaths per year (<http://www.who.int/tdr/>, last accessed May 2008). Apart from infecting humans, *T. brucei brucei*, *T. congolense*, and *T. vivax* also cause Nagana in cattle in Africa, and some species of *Leishmania* lead to viscerocutaneous leishmaniasis in dogs. The medically important trypanosomatids have digenic life cycles that alternate between a mammalian host and an insect vector, usually the responsible for disease transmission (Fig. 1). Another member of Trypanosomatidae is *Crithidia fasciculata*, a parasite of mosquitoes, noninfectious to animals and, for this reason, frequently used as model for the disease-causing species. Being phylogenetically close, trypanosomatids share many biochemical traits, among which


a peroxide metabolism exclusively dependent on a unique dithiol, trypanothione. Throughout this review, the various trypanothione-dependent peroxidases of trypanosomatids will be presented from a comparative point of view, and their importance to the parasites analyzed.

### Trypanothione

In nearly all living organisms, maintenance of an intracellular reducing milieu is possible by means of high concentrations of the sulfur-containing tripeptide, glutathione (GSH). In members of the order Kinetoplastida, however, most of their glutathione content is found in the form of a unique thiol, *N*<sup>1</sup>,*N*<sup>8</sup>-bis(glutathionyl)spermidine, also known as trypanothione (34).

Trypanothione is a conjugate of two glutathione molecules with one spermidine. In trypanosomatids, as in other organisms, glutathione is synthesized by the consecutive activity of the enzymes  $\alpha$ -glutamylcysteine synthetase and glutathione synthetase (41, 51). This is not the case of spermidine, for which the biosynthetic pathway diverges according to the trypanosomatid species. While in *T. brucei*, *C. fasciculata*, and *Leishmania* spp. this polyamine is synthesized from or-

<sup>1</sup>Instituto de Biologia Molecular e Celular (IBMC) and <sup>2</sup>Instituto de Ciências Biomédicas Abel Salazar (ICBAS), Universidade do Porto, Porto, Portugal.



**FIG. 1. Life cycles of *Trypanosoma brucei*, *T. cruzi*, and *Leishmania* spp.** *T. brucei* is transmitted to man and to other mammals when *Glossina* spp. (tsetse fly) has a bloodmeal. This parasite is extracellular and flagellated during the complete life cycle. In the mammalian hosts, *T. brucei* is found in the circulatory and lymphatic systems. Both man and cattle are susceptible to disease. *T. cruzi* is transmitted to a wide range of mammals through the feces of infected triatomines, but only man is afflicted by the disease. In mammals, *T. cruzi* alternates between an extracellular flagellated infective stage and an intracellular aflagellated replicative form, the so-called amastigote. Almost any cell, phagocytic or nonphagocytic, can be infected, although the parasite has a tropism for muscle and nervous cells. *Leishmania* spp. are a complex of over 20 species. Infective flagellated forms, inoculated into different mammals by the bite of phlebotomine insects, are internalized by phagocytic cells, mainly macrophages. In phagolysosomes, those transform into amastigotes that replicate and infect new macrophages. Both man and dogs can develop leishmaniasis.


nithine and methionine via a ubiquitous pathway consisting of four enzymatic steps, in *T. cruzi* one of the enzymes, ornithine decarboxylase, is absent, and these organisms depend on efficient polyamine uptake by high-affinity transporters (46).

The synthesis of trypanothione starts with formation of glutathionylspermidine, followed by insertion of a second glutathione molecule to yield trypanothione. These ATP-dependent reactions are catalyzed by one single enzyme, trypanothione synthetase (TRYS) (25, 26, 78, 79). Trypanothione biosynthesis shows, nevertheless, some divergence among the individual trypanosomatids. Indeed, contrary to what was observed in *T. brucei* (25) and in some species of *Leishmania* (79), *C. fasciculata* harbors an additional enzyme capable of driving the first step of synthesis, glutathionylspermidine synthetase (GSPS) (26). Genes encoding for putative GSPSs are also found in the genomes of *T. cruzi* and *L. infantum* (<http://www.genedb.org/>, last accessed May 2008). *T. cruzi* TRYs is additionally capable of conjugating glutathione with other polyamines (5), a feature that may be advantageous during a potential shortage of polyamines in the surrounding environment.

Since its discovery in 1985, a myriad of biological functions have been directly or indirectly attributed to trypan-

othione, namely ascorbate homeostasis, synthesis of deoxyribonucleotides by ribonucleotide reductase, conjugation and export of metals and drugs, glyoxal removal, and peroxide metabolism (reviewed in ref. 60). These reactions consume reducing equivalents derived from dihydrotrypanothione [ $T(SH)_2$ ], the reduced form of trypanothione, and generate trypanothione disulfide ( $TS_2$ ). Regeneration of the dithiol is made possible by the activity of trypanothione reductase (TR) (Fig. 2), an enzyme present in all Kinetoplastida representatives (reviewed in ref. 36). TR belongs to the protein family of FAD disulfide oxidoreductases, that also comprises glutathione reductase, lipoamide dehydrogenase, thioredoxin reductase, and the flavoprotein AhpF of the alkylhydroperoxide reductase system of *S. typhimurium*. The catalytic mechanism of TR resembles that of glutathione reductase, the main difference being the high specificity of each enzyme towards their respective disulfide substrates (reviewed in ref. 59).

Trypanothione participation in numerous physiological pathways renders this thiol crucial for Kinetoplastida survival. Illustrating this, TRYs was found essential for *T. brucei* survival (6, 27) and TR was shown to be critical for survival and/or infectivity of *L. donovani* and *T. brucei* (reviewed in ref. 75).



**FIG. 2. Peroxidases of trypanosomatids.** Hydroperoxide and peroxynitrite ( $\text{ONOO}^-$ ) elimination in trypanosomatids is mainly carried out by peroxidases (dark gray boxes) belonging to three distinct families of enzymes: 2-Cys peroxidoredoxins (2-Cys PRX), nonselenium glutathione peroxidase-like enzymes (nsGPX), and ascorbate peroxidases (APX). Trypanosomatid nsGPXs can be further segregated into groups A and B, according to their amino acid sequences and their preference for the reductant. In these organisms, peroxidases are fueled by reducing equivalents derived from dihydrotrypanothione  $[\text{T}(\text{SH})_2]$ , which itself is kept

reduced by the NADPH-dependent flavoenzyme trypanothione reductase (TR). Electron transfer between  $\text{T}(\text{SH})_2$  and the various peroxidases occurs via an intermediate molecule, either tryparedoxin (TXN), glutathione (GSH), or ascorbate (ASC). Molecules, such as thioredoxin, can also act as redox shuttles between trypanothione and the peroxidases, although less efficiently than TXN. The broken arrow was chosen to point out the high  $K_M$  of nsGPX-B for GSH. ROOH, small organic chain hydroperoxides (e.g., *t*-butyl hydroperoxide and cumene hydroperoxide); LipOOH, lipid hydroperoxides (e.g., linoleic acid hydroperoxide and phosphatidylcholine hydroperoxide).

### Trypanothione-dependent Peroxidases

It has long been known that trypanosomatids lack catalase and classical selenium-containing glutathione peroxidases, two major hydroperoxide-eliminating enzymes generally present in eukaryotes (reviewed in ref. 36). Instead, these organisms' hydroperoxide metabolism was found to depend on trypanothione (16, 47, 82, 83), even though no trypanothione peroxidase molecule could ever be isolated. The enzymatic pathway linking trypanothione to hydroperoxide reduction was eventually solved in 1997 (76), as consisting of the concerted action of two proteins, a tryparedoxin and a tryparedoxin peroxidase belonging to the peroxidoredoxin family of enzymes. This unique trypanothione-dependent antioxidant system, first described in *C. fasciculata* (76), is common to all pathogenic species of the Trypanosomatidae family, including *T. brucei rhodesiense* (32), *L. infantum* (18), *T. brucei brucei*, *T. cruzi*, *L. major*, and *L. donovani* (reviewed in ref. 60). Since the dissection of this enzymatic pathway, other trypanothione-dependent systems culminating in hydroperoxide reduction have been described, as detailed later in this review.

#### Tryparedoxin/tryparedoxin peroxidase pathways

**Tryparedoxins.** Tryparedoxin (TXN) is the term coined to designate members of the thioredoxin superfamily that are found exclusively in Kinetoplastida (76). TXNs are trypanothione-dependent oxidoreductases that differ from typical thioredoxins in several aspects: (a) they share only 15% homology with thioredoxins, (b) are ~5 kDa larger than thioredoxins, and (c) the TXNs' active site motif is Trp-Cys-Pro-Cys-Arg, instead of Trp-Cys-(Gly/Ala)-Pro-Cys-Lys for most thioredoxins. In addition, unlike typical thioredoxins, TXNs are not directly reduced by a NADPH-dependent flavoprotein. Instead, the electron transfer between the flavoprotein trypanothione reductase (TR) and TXN is mediated by trypanothione. TXN can also be reduced by glutathione,

although with a very low efficacy (19, 40, 65), unlikely to be physiologically relevant. Kinetically, TXN follows an enzyme substitution or ping-pong mechanism, whereby the reducing  $[\text{T}(\text{SH})_2]$  and the oxidizing (peroxidase) substrates react with the enzyme in two independent steps. The rate limiting step of the catalysis is the reaction with trypanothione (19, 40, 109). TXNs are highly abundant proteins (28, 76, 109), representing 3–5% of the total soluble cell protein content of *C. fasciculata* (76) and *T. cruzi* (109). They are found in the cytosol and, at least in *Leishmania* (19), also in the parasite's single mitochondrion. Depending on their subcellular compartmentalization, TXNs react with different tryparedoxin peroxidases (76, 109–111) and also with other substrates, namely ribonucleotide reductase (31) and possibly the Universal Minicircle Sequence Binding Protein (UMSBP) (77), in this way being linked to diverse cell functions such as peroxide metabolism, synthesis of deoxynucleotides, and regulation of mitochondrial DNA replication, respectively.

**Tryparedoxin peroxidases.** The term "tryparedoxin peroxidase" designates all peroxidases that use TXN as source of reducing electrons during removal of peroxides. These include members of the peroxidoredoxin and of the non-selenium glutathione peroxidase families (Fig. 2).

**2-Cysteine peroxidoredoxins (2-Cys PRXs).** Peroxidoredoxins (PRXs) represent a ubiquitous family of antioxidant enzymes with hydroperoxide and peroxynitrite reducing activity. Recently, these enzymes have also been attributed the function of chaperones (22, 52, 53, 71). Peroxidoredoxins lack prosthetic groups or tightly bound metal ions and, instead, use redox active cysteines (Cys) to reduce their substrates. Depending on the number of Cys residues directly involved in catalysis and on the mechanism of reaction, peroxidoredoxins fall into three categories: (a) typical 2-Cys peroxidoredoxins, (b) atypi-


TABLE 1. COMPARATIVE ANALYSIS OF THE PEROXIDASES OF THE HUMAN TRYPANOSOMATIDS *Trypanosoma brucei*, *T. cruzi*, AND *Leishmania* spp.

Organism	Peroxidase	Subcellular compartment	Reductant	Peroxide preference	$k'$ ( $M^{-1} s^{-1}$ )	References	Author's designation
<i>T. brucei</i>	2-Cys PRX	Cytosol	TXN	H <sub>2</sub> O <sub>2</sub> /ROOH ONOO <sup>-</sup>	$1.7 \times 10^7$ $9 \times 10^5$	14, 99 100	TRYP1 or TacTXNPx
	nsGPX	Mitochondrion	TXN	H <sub>2</sub> O <sub>2</sub> /ROOH	ND	99	TRYP2
	nsGPX-A1/2	Cytosol	TXN	H <sub>2</sub> O <sub>2</sub> /ROOH/ThyOOH <sup>§</sup>	ND	49, 93	Px I and Px II
	nsGPX-A3	Mitochondrion and glycosomes	TXN	H <sub>2</sub> O <sub>2</sub> /ROOH/ThyOOH	$1 \times 10^5$	24, 49, 93	Px III
<i>T. cruzi</i>	APX	ND	ND	ND	ND	—	—
	2-Cys PRX	—	—	—	—	—	—
	PRX1*	Cytosol	TXN	H <sub>2</sub> O <sub>2</sub> /ROOH ONOO <sup>-</sup>	$3.2 \times 10^4$ $7.2 \times 10^5$	42, 108 100	TcCPX or TcTXNPx
	PRX2	Mitochondrion	TXN	H <sub>2</sub> O <sub>2</sub> /ROOH	ND	108	TcMPX
<i>Leishmania</i> spp.	nsGPX	Cytosol and glycosomes	TXN	ROOH/LipOOH	$2.4 \times 10^5$	107, 109	TcGPXI
	nsGPX-A1	Cytosol†	ND	ND	ND	—	—
	nsGPX-A2/3	ER	GSH ( $K_M = 5$ mm)	LipOOH	$1.8 \times 10^4$	111	TcGPXII
	nsGPX-B	ER	Ascorbate	H <sub>2</sub> O <sub>2</sub>	$3.5 \times 10^6$	110	TcAPX
	APX	ER	TXN	H <sub>2</sub> O <sub>2</sub> /ROOH	$2.4 \times 10^5$	8, 9, 18, 37, 54, 61	LicTXNPx1/2, LcPxn2/1,
	2-Cys PRX	Cytosol	TXN	ONOO <sup>-</sup>	$1 \times 10^6$	a	LaPxn2/1, Lmf30 or LdTXNPx
	PRX3	Glycosomes†	ND	ND	ND	8	LcPxn3
				H <sub>2</sub> O <sub>2</sub> /ROOH	$3.8 \times 10^6$	17, 18	LimTXNPx
				ONOO <sup>-</sup>	$1.6 \times 10^6$	a	
	nsGPX	Mitochondrion and/or glycosomes†	TXN	H <sub>2</sub> O <sub>2</sub> /ROOH	ND	—	—
	nsGPX-A1	Mitochondrion and/or glycosomes†	ND	ND	ND	—	—
	nsGPX-A2†	Mitochondrion†	ND	ND	ND	—	—
	nsGPX-A3	Cytosol†	ND	ND	ND	—	—
	nsGPX-B	ND	ND	ND	ND	—	—
	APX	ND	Ascorbate	H <sub>2</sub> O <sub>2</sub>	ND	2	LmAPX

\*In the *T. cruzi* genome, two additional open reading frames are annotated, which code for protein sequences carrying one amino substitution relative to PRX1; the substitutions are unlikely to be functionally relevant and for that reason the molecules are not listed here; †not annotated in the *L. infantum* genome project; ‡based on the presence or absence of an obvious organelle targeting signal and on bioinformatics analysis, but not experimentally demonstrated; §not determined, but possibly the same as the *T. brucei* nsGPX-A3; ¶S. Romao and Ana M. Tomás, unpublished.

ND, not determined; ROOH, small chain organic hydroperoxides (e.g., *t*-butyl and cumene hydroperoxides); ThyOOH, thymine hydroperoxide; LipOOH, lipid hydroperoxides (e.g., linoleic acid and phosphatidylcholine hydroperoxides); Lm, *L. major*; Ld, *L. donovani*; Li, *L. infantum*; Lc, *L. chagasi*; La, *L. aethiopica*.





**FIG. 3.** Alignment of trypanosomatid 2-Cys peroxiredoxins. Sequence accession numbers are the following: Tb\_PRX1, AAG45225; Tc\_PRX1, CAA09922; Li\_PRX1, AY058210; Li\_PRX2, AAG40074 (*Leishmania chagasi*); Li\_PRX3, AAK69587 (*L. chagasi*); Tb\_PRX2, Q9GU47; Tc\_PRX2, CAA06923; Li\_PRX4, AY058209. In the *T. cruzi* genome, two additional open reading frames are annotated, which code for protein sequences carrying one amino acid substitution relative to Tc\_PRX1. These substitutions are unlikely to be functionally relevant and, for that reason, the sequences are not shown here. The VCP and IPC motifs directly involved in catalysis are highlighted in *white letters on black*. Residues implicated in 2-Cys PRX sensitivity to overoxidation by  $H_2O_2$  are shown in *white letters on gray*. Amino-terminus mitochondrial and C-terminus glycosomal targeting signals are marked on *light gray background*. A sequence homologous to CDKs-mediated phosphorylation motifs is depicted in **bold on dark gray**. Tb, *T. brucei*; Tc, *T. cruzi*; Li, *L. infantum* (same as *L. chagasi*).

cal 2-Cys peroxiredoxins, and (c) 1-Cys peroxiredoxins (reviewed in ref. 86).

Peroxiredoxins were the first enzymes reported to display trypanredoxin peroxidase (TXNPx) activity (76). Peroxiredoxins of trypanosomatids are typical 2-Cys peroxiredoxins and can be grouped according to their cytosolic or mitochondrial compartmentalization (Table 1). Mitochondrial peroxiredoxins are unusual in having the second redox active Cys embedded in an Ile-Pro-Cys motif, instead of Val-Cys-Pro often found in 2-Cys peroxiredoxins (Fig. 3). Also, mitochondrial peroxiredoxins are encoded by single-copy genes, while the cytosolic enzymes may include closely related genes clustered within the same chromosomal locus (see Fig. 3. for accession numbers). In *Leishmania* spp., one of these genes carries a glycosomal targeting signal, suggesting that the enzyme may localize to this peroxisome-like organelle. Trypanosomatid peroxiredoxins are broad-spectrum peroxidases acting on  $H_2O_2$  and organic hydroperoxides. The crithidial enzyme also decomposes complex lipid hydroperoxides (76); however, the peroxiredoxins of *T. brucei*

and *Leishmania* react poorly and are rapidly inactivated by these substrates (in particular, linoleic acid and phosphatidylcholin hydroperoxides) (14, 17, 37, 61). In addition, the Kinetoplastida enzymes act catalytically on peroxynitrite decomposition at considerably high rates (9, 100, Table 1). Evidences about the function of peroxiredoxins within the parasites was obtained from *Leishmania* and *T. cruzi* overexpression mutants, which are more resistant to hydroperoxides and peroxynitrite added exogenously (9, 18, 62, 108), and also from double-stranded RNA interference in *T. brucei* (112).

Typical 2-Cys peroxiredoxins are obligate homodimers, containing subunits of ~22 kDa and two identical reaction centres. According to the generally accepted mechanism for these enzymes, the proximal Cys of one enzyme subunit reduces the peroxide substrate being oxidized to a cysteine sulfenic acid (Cys-SOH). This residue is subsequently attacked by the C-terminal Cys of an inverted subunit to form a stable intersubunit disulfide bond, which is then resolved by a cell-specific disulfide oxidoreductase (reviewed in ref.

86). In the case of trypanosomatids, this oxidoreductase is TXN. Such sequence of independent catalytic steps meets the definition of an enzyme substitution mechanism, typical for peroxidases in general. Surprisingly, however, deviations from this mechanism were observed upon kinetic analysis of the peroxidoredoxin enzymes of *L. infantum* and *T. brucei* (14, 17), and also upon review of the original kinetic data of *C. fasciculata* and *T. cruzi* (14). The results revealed that the enzymatic activity slowed down as the hydroperoxide concentration increased. Although the reasons for this behavior remain undisclosed, Budde *et al.* (14) provided an interpretation based on the Changeux–Monod model for oligomeric proteins (70). The authors could apply this model to 2-Cys peroxidoredoxins because they exist as decamers built up of five dimers arranged in a ring-like structure, the transition between the dimeric and the decameric states being redox-dependent (3, 14, 17, 116). According to the proposed model, oxidation of one reaction center within the decamer would induce conformational changes in remote dimers, ultimately impairing their reactivity with the hydroperoxide substrate (14).

In the majority of eukaryotic cells, peroxidoredoxins are considered to act as regulators of  $H_2O_2$ -mediated intracellular signaling processes, such as cell differentiation, proliferation, and apoptosis, and not as general antioxidant devices. Such fine-tuning role is possible because these enzymes are sensitive to inactivation by relatively low amounts of  $H_2O_2$ , a phenomenon attributed to overoxidation of the proximal Cys by the hydroperoxide substrate (reviewed in refs. 91, 92, 115). The inactive sulfenic acid residue ( $-SO_2H$ ) resulting from overoxidation can be reduced back to the catalytically active thiol form by a sulfiredoxin or a sestrin molecule (reviewed in ref. 55). In the case of trypanosomatids, it is, nevertheless, unclear whether peroxidoredoxins are involved in regulation. Certainly, these parasites possess additional peroxidases (presented later in this review) capable of general antioxidant functions. Also, in trypanosomatids peroxidoredoxins are likely present in large amounts (76), a feature that has been related to their regulatory role in other organisms (91, 115). Furthermore, the two motifs required for  $H_2O_2$ -sensitivity [*i.e.*, the Gly–Gly–Leu–Gly motif and the C-terminal arm containing a Tyr–Phe sequence (115)], are conserved in almost all trypanosomatid peroxidoredoxin sequences (Fig. 3). Dissection of the crystal structure of the *T. cruzi* cytosolic enzyme strongly suggests that these motifs may modulate activity (85) and the kinetic behavior of these enzymes complies with sensitivity to hydroperoxide inactivation (14, 17). So far, however, the only evidence that trypanosomatid peroxidoredoxins can be overoxidized came from a study performed on ONOO<sup>−</sup>-challenged *T. cruzi* (84). In addition, molecules capable of reverting the overoxidized state of peroxidoredoxins are apparently missing in these parasites. Indeed, sulfiredoxin is not present in the trypanosomatid genome and, although all three medically relevant organisms possess a putative open reading frame containing a sestrin signature (accession numbers: LmjF18.0650, LinJ18\_V3.0660, Tb10.05.0190, Tc00.1047053511277.130), the homology to sestrins of higher eukaryotes is too low to infer a function.

One consequence of inactivation by  $H_2O_2$  is the aggregation of peroxidoredoxins into complexes with molecular weight higher than the decamer, which is usually associated with loss of peroxidase activity and appearance of chaperone ac-

tivity (22, 52). More recently, aggregation and functional switching of peroxidoredoxins was described to also take place as a result of phosphorylation by cyclin-dependent kinases (CDKs) (53). Phosphorylation occurs at the threonine residue located 38 residues downstream of the proximal redox-active cysteine, sitting in a (Ser/Thr)–Pro–Xaa–(Lys/Arg) consensus sequence (20). Interestingly, in all mitochondrial peroxidoredoxins of trypanosomatids this sequence is conserved (Fig. 3). However, their subcellular compartmentalization renders phosphorylation by CDKs unlikely.

**Non-selenium glutathione peroxidase enzymes (nsGPX-As).** Glutathione peroxidases (GPXs) comprise a group of enzymes that act as pivotal antioxidants by reducing  $H_2O_2$  or organic hydroperoxides with high catalytic efficiency ( $\sim 10^8 M^{-1} s^{-1}$ ) and in different cellular locations (reviewed in ref. 13). GPXs share a common catalytic core formed by selenocysteine (SeCys), tryptophan, and a glutamine residue. However, in some GPXs, the SeCys residue is replaced by a Cys. Such substitution, characteristic of the so-called “non-selenium GPX-like enzymes” (nsGPXs), confers to these proteins decreased peroxidase activity (reviewed in ref. 48).

Elucidation of the genomes of trypanosomatids revealed that these organisms possess a genomic *locus* comprising a cluster of two to three genes coding for closely related nsGPXs, that we classify here as nsGPX-A (Table 1). In *T. cruzi*, one of these proteins, nsGPX-A1, is glycosomal, a small fraction possibly localizing also to the cytosol (109). Regarding the other two, the absence of any obvious organelle endorsement signal and the detection of a substantial level of glutathione peroxidase activity in the parasite cytosol, suggests that they are cytosolic (109). In both *T. brucei* and *Leishmania* spp., the nsGPX-A molecules are almost identical except in the very N- and C-terminal regions, which may possess signal peptides for glycosomes and for the mitochondrion. The *T. brucei* nsGPX-A3 and the *Leishmania* spp. nsGPX-A1 molecules carry both peptide endorsement sequences. In the case of the *T. brucei* enzyme, experimental evidence points towards its dual localization in both organelles in the insect stage of the parasite (24, 93). As for the other two *T. brucei* isoforms (nsGPX-A1 and 2), they are cytosolic (93).

The highest degree of similarity of Kinetoplastida nsGPXs is found with GPXs proteins from plants, which belong to the phospholipid hydroperoxide GPX (PHGPX) clade. Trypanosomatid nsGPXs, like PHGPXs, contain deletions in regions thought to mediate protein dimerization/tetramerization (12) (Fig. 4). Absence of such regions accounts for the monomeric nature of these enzymes and, as suggested for PHGPXs, may facilitate the access of complex hydroperoxides to the catalytic core. Accordingly, the *T. cruzi* nsGPX-A1 metabolizes fatty acid and phospholipids hydroperoxides (107). That is not the case, however, of the *T. brucei* nsGPX-A3, for which no activity towards lipid hydroperoxides was detected (93). Instead, this enzyme preferentially reacts with  $H_2O_2$ , additionally accepting linoleic acid, *t*-butyl, and thymine hydroperoxides as substrates. More recently, the monomeric nature of nsGPXs was postulated to be also a prerequisite for interaction with thioredoxin and related proteins containing a Cys–Xaa–Xaa–Cys motif (66).

One particularity of nsGPXs concerns their affinity for glutathione. In nsGPXs, similarly to that observed in PHGPXs, the residues involved in glutathione binding (33, Fig. 4) are



Hs_cGPX	-----MCAAKLAAAAAQSVTAFAARPLAGSGPVSLSGL	33
Hs_PHGPX	MSLGLRLCRLKLPALLCGALAAPGLAGTMCASRDWRCARSMHEFSAKDIDG-HMVNLDKY	59
Tb_nsGPX-A1	-----MSTIFDFEVLDAH-KPYNLVQH	22
Tb_nsGPX-A2	-----MSAASSIFDFEVLDAH-KPYNLVQH	25
Tb_nsGPX-A3	-----MLRSEKREMSAASSIFDFEVLDAH-KPYNLVQH	33
Tc_nsGPX-A1	-----MFRFGQLLRAAEKHSIYEFQVNAADG-KPYDLSC	34
Tc_nsGPX-A2	-----MTTVVDFQVNAADG-KPYDLSC	22
Tc_nsGPX-A3	-----MTSEG---TDNAHSSIYDFQILDADH-QLYDLSC	31
Lm_nsGPX-A1	-----MLRLPFFPRAAAAQAASSIYDFKVGSDH-QPYDLGQH	37
Lm_nsGPX-A2	-----MLRLPFFPRAAAAQAASSIYDFKVGSDH-QPYDLGQH	37
Lm_nsGPX-A3	-----MSIYDFKVGSDH-QPYDLGQH	21
Tb_nsGPX-B	-----NNGGAIFSHVLMDSN--AVNLSKY	23
Tc_nsGPX-B	-----NGSSTVFAYSALMSGK--SVALSNY	23
Lm_nsGPX-B	-----NAS--VETYSAVQNGK--TVVLQKY	21
Hs_cGPX	RGKVLLEINVASLGGTTVLDYTCQNHQLQRLGPRGLVVLGFPNCQFGRGENA <del>NEE</del> ILNS	93
Hs_PHGPX	RGFVCIVTNVASQGGKTEVNYTQLVDLHARYAEQGLRLAFPCNQFGHDEPGSNEEIK--	117
Tb_nsGPX-A1	EKSPLLIYNVASKCGYTKGGYETATALYNKYKSGGFTVLVFPNCFQGGCEAGNEEIEIK--	80
Tb_nsGPX-A2	EKSPLLIYNVASKCGYTKGGYETATTLYNKYKSGGFTVLAFPCNQFGGCEPGTEEEIEIK--	83
Tb_nsGPX-A3	EKSPLLIYNVASKCGYTKGGYETATALYNKYKSGGFTVLAFPCNQFGGCEPGTEEEIEIK--	91
Tc_nsGPX-A1	EKGHLLIYNVASKCGYTKGGYETATTLYNKYKSGGFTVLAFPCNQFAGCEPGTALEVK--	92
Tc_nsGPX-A2	EKGHLLIYNVASKCGYTKGGYETATTLYNKYKSGGFTVLAFPCNQFAGCEPGTAVEVK--	80
Tc_nsGPX-A3	EKGHLLIYNVASKCGYTKGGYETATTLYNKYKSGGFTVLAFPCNQFAGCEPGTALEVK--	89
Lm_nsGPX-A1	EKGHLLIYNVASKCGYTKGGYETATALYNKYKHLGFHVLAFPCNQFAGCEPGTEEEVK--	95
Lm_nsGPX-A2	EKGHLLIYNVASKCGYTKGGYETATALYNKYKHLGFHVLAFPCNQFAGCEPGTEEEVK--	95
Lm_nsGPX-A3	EKGHLLIYNVASKCGYTKGGYETATALYNKYKHLGFHVLAFPCNQFAGCEPGTEEEVK--	79
Tb_nsGPX-B	AGCVTVLVNTASLCSFTSSNIQHLLHVVQKWSRSTVLAFPCSQFGNCEPKRDEIC--	81
Tc_nsGPX-B	TGRVTVVVNTASLCSFANSSLQQLTHVQETYGPRGFTILAFPCAQFANCEPKSNEEIA--	81
Lm_nsGPX-B	SGYATLIVNVASCSLSTNIEMINEVQQAAGSRRTVLAFPCAQFANCEPLNTEIA--	79
Hs_cGPX	LKYVPPGGGFEPNEMLFKECEVNGAGAHPLFAFLREALPAPSDATAIMTDPKLITWSPV	153
Hs_PHGPX	---EFAAG-YNVEFDMEFKICVNGDDAHPLWQWQIQPKGG-----IL	157
Tb_nsGPX-A1	---DFVCTKFAEFFIMAKINVGNAHPLYEYMKKTKFG-----IL	119
Tb_nsGPX-A2	---DFVCTKFAEFFIMAKINVGNAHPLYEYMKKTKFG-----IL	122
Tb_nsGPX-A3	---DFVCTKFAEFFIMAKINVGNAHPLYEYMKKTKFG-----IL	130
Tc_nsGPX-A1	---DFACTRFKADFFIMEKIDVNGGKAHPLYEFMKATIFG-----LF	131
Tc_nsGPX-A2	---DFACTRFKADFFIMEKIDVNGGKAHPLYEFMKSLKFG-----IL	119
Tc_nsGPX-A3	---DFACTRFKADFFIMEKIDVNGGKAHPLYEFMKSLKFG-----SA	128
Lm_nsGPX-A1	---DFACTRFKADFFIMEKVCVNGEHEHPLYHYLKNTCKG-----IL	134
Lm_nsGPX-A2	---DFACTRFKADFFIMEKVCVNGEHEHPLYHYLKNTCKG-----IL	134
Lm_nsGPX-A3	---DFACTRFKADFFIMEKVCVNGEHEHPLYHYLKNTCKG-----IL	118
Tb_nsGPX-B	---CWVANGGINEPVPFDKVNHLKGNTHPLFQMIQSSLGK-----	117
Tc_nsGPX-B	---VWAQTCGLNPLFLDRVKVKGPDHPLFQHLQASLGK-----	117
Lm_nsGPX-B	---QWCEDLGLLFPVFDKVNKVGSSADPLFQHLQASLGK-----	115
Hs_cGPX	CRNDVANNFEKFLVGFPGVPLRYS <del>RL</del> FQTIDIEPDIEALLSQGPSCA-----	201
Hs_PHGPX	G-NAIKNNFTFLIDRNGCVVERFYSGASVKDIEKKLIPLESTQSA-----	197
Tb_nsGPX-A1	ATKAIKNNFTSFLIDRDGVVERFSPGASVKDIEKKLIPLESTQSA-----	166
Tb_nsGPX-A2	KTKAIKNNFTSFLIDRDGVVERFSPGASVKDIEKKLIPLESTQSA-----	169
Tb_nsGPX-A3	ATKAIKNNFTSFLIDRDGVVERFSPGASVKDIEKKLIPLLGSARI-----	176
Tc_nsGPX-A1	GTKAIKNNFTSFLIDRHGVVERFSPGASVEDIEKKLIPLLGSARI-----	177
Tc_nsGPX-A2	GTKAIKNNFTSFLIDRHGVVERFSPGASVEEIEKKLIPILNESN-----	164
Tc_nsGPX-A3	GVQAINNNFTSFLIDRHGVVARFSPGASVEEIEKLLPFLFEGEVVSGPS-----	178
Lm_nsGPX-A1	GTTLVKNNFTAFVLVDKDGHAVCRFAPGATVSEIEKKLIPLLQAAS <del>SKL</del> -----	183
Lm_nsGPX-A2	GTTLVKNNFTAFVLVDKDGHAVCRFAPGATVSEIEKKLIPLLQADSGDAKVLPSQS	190
Lm_nsGPX-A3	GTTLVKNNFTAFVLVDKDGHAVCRFAPGATVSEIEKKLIPLLQADSGDAKVLPSQS	174
Tb_nsGPX-B	----IRNNYTKVVCNRAGLPCVELQPGSSLEELERYVSQLCDE-----	156
Tc_nsGPX-B	----IRNNYTKICDKSGIPLVKLDSSFLLELRSEIQACGPQGA-----	160
Lm_nsGPX-B	----PLNNYTKYLCDSGGVPRKLEPGCSMDALRQSIECVL-----	152

**FIG. 4. Alignment of trypanosomatidal nonselenium glutathione peroxidases.** The trypanosomatidal nonselenium GPXs are aligned with the human cytosolic GPX (Hs\_cGPX, P07203) and the human phospholipid hydroperoxide GPX (Hs\_PHGPX, NP\_002076). Sequence accession numbers are the following: Tb\_nsGPX-A1, Q869A7; Tb\_nsGPX-A2, Q869A6; Tb\_nsGPX-A3, Q869A5; Tc\_nsGPX-A1, CAC85914; Lm\_nsGPX-A1, Q4Q9B4; Lm\_nsGPX-A2, Q4Q9B3; Lm\_nsGPX-A3, Q4Q9B2; Tc\_nsGPX-B, CAC85915; Lm\_nsGPX-B, Q4Q1B8; Tc\_nsGPX-A2, Tc\_nsGPX-A3 and Tb\_nsGPX-B sequences were derived from the genome projects. The conserved SeCys (represented by U), Gln and Trp residues characteristic of GPX are highlighted in *white letters on black*; in nsGPXs the SeCys is replaced by Cys. The consensus motif containing the distal redox active Cys of trypanosomatidal nsGPXs is shown in *white letters on gray*. Amino-terminus mitochondrial and C-terminus glycosomal targeting signals are marked on *light gray background*. Residues involved in glutathione binding are annotated in **bold on dark gray**. Tb, *T. brucei*; Tc, *T. cruzi*; Lm, *L. major*.

either mutated or deleted, rendering the affinity for this thiol weak (reviewed in ref. 48). This has led to the suggestion that glutathione is not their main reductant in the cell. In fact, some nsGPXs were shown to use thioredoxin as reducing substrate (reviewed in ref. 48). Trypanosomatid nsGPXs also lack the glutathione-binding residues and, consequently, display extremely low activities with this thiol, which translate into high  $K_M$  values, within the millimolar range (49, 107, 111). In the case of nsGPX-As, they clearly prefer trypanothione as source of reducing equivalents (49, 109) and at least the *T. brucei* nsGPX-A3 can also use thioredoxin as reductant, although with less efficiency (49). The catalytic mechanism for the reaction of the *T. brucei* nsGPX-A3 with the Cys-Xaa-Xaa-Cys motif of trypanothione was recently disclosed (94) and shown to comply with that of other nsGPXs with thioredoxin (48, 66). Two cysteine residues are directly involved in catalysis: (a) Cys47 (*T. brucei* nsGPX-A3 numbering), which replaces the SeCys of classical GPXs, and (b) Cys95, which is part of the (Phe/Tyr)-(Ala/Val)-Cys-Thr consensus motif present in all trypanosomatid nsGPX-As (Fig. 4). Cysteine47 is the residue that directly reacts with the hydroperoxide substrate, being oxidized to a sulfenic acid. This is then attacked by Cys95, the resulting intramolecular disulfide bridge being reduced by the Cys-Xaa-Xaa-Cys motif of the oxidoreductase. This mechanism differs from that of selenium-dependent GPXs and rather resembles that of atypical 2-Cys peroxiredoxins.

#### *Nonselenium glutathione peroxidases not dependent on trypanothione (nsGPX-B)*

Apart from the three identical clustered nsGPX-A genes, at least one additionally open reading frame, encoding a different nsGPX enzyme, can be found in a separate locus in the genomes of trypanosomatids (Fig. 4). These nsGPXs, which are classified here as nsGPX-B, share only 30% identity with nsGPX-As. One of these gene products was characterized in *T. cruzi* and assigned to the ER (111). As expected for a nsGPX, the *T. cruzi* enzyme metabolizes fatty acid and phospholipid hydroperoxides and reacts poorly with GSH. However, unlike trypanosomatid nsGPX-As, it does not accept TXN as reductant. This may, at least in part, be explained by the fact that the consensus motif containing the C-terminal Cys residue implicated in nsGPX-A catalysis with TXN (Cys95 of *T. brucei* nsGPX-A3, 94) is missing in trypanosomatid nsGPX-B (Fig. 4). Indeed, replacement of this Cys by a serine residue was shown to abolish the peroxidase activity of the *T. brucei* nsGPX-A3 in the presence of TXN (94). The possibility exists that other molecules, more efficient than GSH, react physiologically with nsGPX-Bs.

The activity of all trypanosomatid nsGPXs is likely dependent on trypanothione, even in the case of enzymes for which thioredoxin (49) or glutathione (111) might be the source of reducing electrons. In fact, in trypanosomatids, lacking thioredoxin reductase and glutathione reductase, the pool of reduced thioredoxin and glutathione is maintained by the TR/trypanothione system (95). In the case of glutathione, reduction may occur either via the direct interaction with the dithiol (56) or via a reaction catalyzed by a trypanothione-glutathione thioltransferase, such as Tc52 in *T. cruzi* (73, 74).

#### *Ascorbate peroxidase*

Ascorbic acid (or vitamin C) is a cofactor for a range of enzymes involved in diverse metabolic pathways, such as protein folding, iron absorption, and the synthesis of collagen, neurotransmitters, peptide hormones, and carnitine (reviewed in refs. 7 and 44). However, its major function is as a powerful antioxidant, either by reducing free radicals (e.g.,  $RO\cdot$ ,  $RO_2\cdot$ ,  $HO\cdot$ ,  $NO_2\cdot$ ), scavenging several nonradical reactive species (e.g., hypochlorous acid, ozone, and nitrating agents derived from peroxynitrite), maintaining  $\alpha$ -tocopherol (vitamin E) in its reduced form, or by acting as electron donor of the ascorbate peroxidase (APX) family of enzymes (44). Detection of ascorbate in trypanosomatids dates back to 1994 (23), but only recently has the capacity of synthesizing vitamin C been demonstrated in *T. brucei* and *T. cruzi*, the enzyme responsible for the terminal step of ascorbate biosynthesis having been characterized in both organisms (64, 113). Contrary to that observed for *T. cruzi*, which acquires ascorbate solely by *de novo* synthesis (64), the *T. brucei* mammalian stage can additionally take up this vitamin from the surrounding environment (113). Ascorbate biosynthesis is distinct in mammals, plants, and yeast. In *T. cruzi* and *T. brucei*, the synthesis of this vitamin likely resembles that of plants, as deduced from genome analysis (64, 113). In the case of *Leishmania* spp., the enzyme responsible for the last catalytic step in the biosynthetic pathway is also present, thus ascorbate metabolism should not significantly differ from that of the other trypanosomatids.

Ascorbate peroxidases (APXs) are class I heme-containing peroxidase enzymes, which catalyze the  $H_2O_2$ -dependent oxidation of ascorbate in photosynthetic organisms (reviewed in ref. 89). It was known since 1980 that *T. cruzi* epimastigote extracts exhibit ascorbate-dependent peroxidase activity (11, 30). Recently, this activity was assigned to the unusual plant-like APX enzyme (110). This protein has a counterpart in *Leishmania* spp. (2), but interestingly, not in *T. brucei*. As inferred from their amino acid sequences (2, 110) and/or gel filtration analysis (2), APXs from *T. cruzi* and *L. major* are monomers within the 30 kDa range. In addition, the *T. cruzi* APX kinetic behavior complies with the monomeric nature of the enzyme (80, 110). Trypanosomatid APXs differ from those of plants in possessing, near the C-terminus, a positively charged insert of unknown function. Also, they carry an N-terminal amino acid extension which is unique in coding for an amino acid topology that is reminiscent of signal sequences targeting proteins to the ER or plasma membrane. In the case of *T. cruzi*, the enzyme locates to the ER (110). The most significant difference to plant APXs, however, is that a residue crucial for ascorbate binding and oxidation, Arg172 (soya-bean cytosolic APX numbering; accession number T07056), is missing in the trypanosomatid enzymes. This may explain the fact that trypanosomatid APXs exhibit relatively low ascorbate utilization and also the nonsaturation kinetics observed for the *L. major* enzyme (2). Still, the fact that the affinity of both enzymes for ascorbate is within the micromolar range has led to the suggestion that an alternative ascorbate-binding mechanism could be operating in these molecules (2).

The catalytic mechanism of trypanosomatid APXs likely conforms to that of plant APXs (2, 96). This involves the very fast second-order reaction of ferric heme with  $H_2O_2$ , con-



sisting in the heterolytic cleavage of the O–O peroxide bond and release of a water molecule. The catalytic intermediate resulting from this step, known as Compound I, is subsequently reduced by ascorbate in two sequential single electron transfer reactions, initially to a second intermediate, Compound II, and then back to ferric heme, regenerating the ground state enzyme. Two monodehydroascorbate radicals result from these reactions, which under physiological conditions, are reduced back to ascorbate either (a) by the activity of a NADH-dependent reductase, or (b) by the rapid disproportionation to dehydroascorbic acid and subsequent recycling by glutathione or by glutathione-dependent enzymes (reviewed in ref. 43). Dehydroascorbate can also decompose irreversibly, yielding a range of products, such as oxalate and threonate (43). In Kinetoplastida, regeneration of ascorbate seems to be achieved nonenzymatically via the spontaneous reduction of dehydroascorbate by trypanothione (58, 110). Even though TXN can also reduce dehydroascorbate, there is no obvious advantage in this reaction (90).

With respect to the hydroperoxide substrates, the *T. cruzi* enzyme specifically reduces  $\text{H}_2\text{O}_2$  but not the organic hydroperoxides *t*-butyl and cumene hydroperoxides. Studies with mutant parasites overexpressing this molecule demonstrated that it can shield cells from exogenously added  $\text{H}_2\text{O}_2$  (110). In the case of the *L. major* APX, it was demonstrated that, besides serving as substrate,  $\text{H}_2\text{O}_2$  is also capable of acting as an inhibitor of this enzyme when the concentration of ascorbate is low (2).  $\text{H}_2\text{O}_2$ -mediated inactivation has also been described in another class I hemeperoxidases (50).

#### Other trypanothione-dependent pathways implicated in antioxidant function

Enzymes of the glutathione S-transferase class are important for the metabolism of phospholipid hydroperoxides in higher eukaryotes. They act by catalyzing the nucleophilic attack of GSH on a wide variety of hydrophobic substrates. In trypanosomatids, no such activity is present and, instead, these organisms possess trypanothione S-transferase activity associated to the eukaryotic translation elongation factor 1B (eEF1B) (102). The *L. major* eEF1B displays trypanothione-dependent peroxidase activity, reacting preferentially with linoleic acid hydroperoxide, but not with  $\text{H}_2\text{O}_2$  (103). This, added to the fact that eEF1B localizes to the ER, has led to the suggestion that this molecule may be involved in the elimination of oxidized lipids within this organelle.

Trypanothione is also an efficient reductant of ovothiol A (4). This mercaptohistidine, present in several trypanosomatids, can act as a nonenzymatic scavenger of free-radicals (reviewed in ref. 97) and can catalyze the decomposition of nitrosothiols (105).

Finally, apart from supplying reducing equivalents to enzymatic cascades that culminate in peroxide elimination, trypanothione can spontaneously reduce  $\text{H}_2\text{O}_2$  (4) and ONOO<sup>−</sup> (100). However, the second order rate constants estimated for these nonenzymatic reactions are low (5.37 and 7200 M<sup>−1</sup> s<sup>−1</sup> for  $\text{H}_2\text{O}_2$  and ONOO<sup>−</sup>, respectively), when compared to that of the peroxidases (Table 1), and are thus negligible under physiological conditions.

#### Complementary and Redundant Roles of Trypanosomatid Peroxidases

In cells, peroxide metabolism is largely ensured by peroxidases, and trypanosomatids are no exception to this. Peroxides are generated internally or derived from exogenous sources. The effect they produce in cells greatly depends on their nature and concentration. They can either cause oxidative damage on a range of biomolecules or act as important effectors in cell signaling pathways.

Within the cytosol of all three clinically relevant trypanosomatids, 2-Cys peroxiredoxins, and another less abundant class of tryparedoxin-dependent peroxidases, the nsGPX-A, are likely in charge of peroxide elimination. There are little doubts about the essentiality of these two sets of enzymes for trypanosomatids' survival, as directly demonstrated for *T. brucei* by RNA interference (93, 112). Still, in the case of nsGPX-As, the possibility that the observed defects are due to loss of functionality of the noncytosolic isoform cannot be discarded, due to similarity between all three sequences. On the whole, the emerging picture points to the existence of specific, nonredundant functions for each class of cytosolic peroxidases in trypanosomatids. This may be explained by the peculiarity of peroxiredoxins to metabolize ONOO<sup>−</sup> (Table 1).

The cytosolic location of an effective antioxidant system is most favorable to confer protection against exogenous sources of oxidants. Upon invasion of the mammalian host, the intramacrophagic pathogens *Leishmania* spp. and *T. cruzi* are challenged with reactive oxygen and nitrogen species derived from the immune response (10, 15, 63). The capacity of 2-Cys peroxiredoxins to protect trypanosomatids from externally derived oxidants does not seem to be compensated by any other hydroperoxide detoxifying pathway. Accordingly, the *T. brucei* ability to cope with  $\text{H}_2\text{O}_2$  added exogenously was impaired upon downregulation of the cytosolic peroxiredoxin, but not of nsGPX-A (112). The combined hydroperoxide and ONOO<sup>−</sup> metabolizing activities of 2-Cys peroxiredoxins are also likely to account for the increased infectivity of *Leishmania* mutants overexpressing a cytosolic peroxiredoxin, in an *ex vivo* model of infection (9). The significance of cytosolic peroxiredoxins for parasite pathogenicity was further proposed for *L. (Viannia) guyanensis*, wherein the parasite ability to disseminate from cutaneous lesions to the nasopharyngeal mucosa was related to upregulation of these enzymes (1, 106).

Additional sources of exogenous oxidative species may be provided by some of the drugs used clinically to treat trypanosomiasis and leishmaniasis (67, 88, 104, 117). Peroxiredoxin association with drug resistance was described for *L. amazonensis*, wherein a strain selected for arsenite resistance displayed upregulation of either the cytosolic or the mitochondrial enzyme (62). This likely happens to counteract an oxidative stress that occurs as a response to this metalloloid drug (68).

Apart from functioning as antioxidant devices, 2-Cys peroxiredoxins and nsGPXs may also be implicated in cell signaling pathways, as described for other organisms (reviewed in refs. 48, 91, 92). In 2-Cys peroxiredoxins, this function has been attributed to their propensity to undergo overoxidation by  $\text{H}_2\text{O}_2$ . This results in inactivation of the peroxidase and subsequent accumulation of the hydroperoxide, a require-

ment for signal propagation. As discussed previously, a number of features in trypanosomatid peroxiredoxins suggests that they might also be sensitive to hyperoxidation. Another peroxide that is being increasingly accepted as a redox-mediated intracellular messenger is ONOO<sup>-</sup> (reviewed in refs. 57, 101). At present, there is no direct evidence for ONOO<sup>-</sup> formation in trypanosomatids. However, its two precursors, superoxide anion and nitric oxide, may be generated in these cells (38, 39, 81, 98, 114). Thus, although rather speculative, the possibility remains that trypanosomatid peroxiredoxins may also participate in ONOO<sup>-</sup>-mediated regulation. Whether in Kinetoplastida 2-Cys peroxiredoxins act as regulators of signaling pathways is to be elucidated. The current belief is that they constitute the parasites' major antioxidant apparatus.

In eukaryotes, the ER is the main site for synthesis and post-translational modification of secretory and membrane proteins, a process that requires an oxidizing milieu. It is also in this organelle where fatty acid and phospholipids are synthesized and metabolized. Within the ER of *T. cruzi* and *Leishmania* spp., at least two enzymes are likely in charge of dealing with the hydroperoxide challenge that may arise therein, namely nsGPX-B and APX (110, 111). These peroxidases differ in terms of electron donor and substrate specificity (Table 1) and their combined activities may account for the parasite ability to metabolize a wide range of hydroperoxides. In this regard, APX may act by eliminating H<sub>2</sub>O<sub>2</sub>, while nsGPX-B may be crucial at repairing the oxidative damage of newly synthesized lipids. Interestingly, no APX coding sequence is annotated in the *T. brucei* genome. Wilkinson *et al.* (113) suggested that *T. brucei* may not require ascorbate-based antioxidant protection because, being an extracellular parasite, it is not exposed to the oxidative challenge produced by host immune cells in response to infection as do *T. cruzi* and *Leishmania* spp. Another molecule possibly in charge of eliminating oxidized lipid hydroperoxides in the endoplasmic reticulum is eEFB1 (102, 103).

Peroxisomes are major sources of reactive oxygen species (reviewed in ref. 29). In trypanosomatids these cell compartments are replaced by related organelles, the glycosomes, in which a number of biochemical processes take place (reviewed in ref. 69). In *T. cruzi* and *T. brucei* glycosomes, the presence of nsGPX-A enzymes was reported (24, 109), which likely constitute an important line of defense against oxidative damage. As for *Leishmania* spp., the presence of nsGPX-As in glycosomes has never been addressed. Although not yet experimentally demonstrated, *Leishmania* spp. possibly possesses a 2-Cys peroxiredoxin in glycosomes (PRX3 in Table 1).

Kinetoplastida organisms possess a single mitochondrion, wherein the electron transport chain leading to molecular oxygen reduction, constitutes one of the most relevant sources of oxidants. In homology to that described for the cytosol, two classes of TXN-dependent peroxidases (the 2-Cys PRXs and the nsGPX-As) are probably in charge of peroxide metabolism in the mitochondrion (Table 1). This situation may, nevertheless, differ with respect to the parasite species, because no nsGPX-A was found in the mitochondrial fractions of *T. cruzi* (109). Apart from general antioxidant defense, other functions have been postulated for 2-Cys peroxiredoxins operating in the mitochon-

dron of trypanosomatids. First, the enzyme may be indirectly implicated in regulation of mitochondrial DNA replication by competing with UMSBP for reducing equivalents derived from TXN (72). Second, in analogy to mitochondrial peroxiredoxins of higher eukaryotes (21), it could be involved in parasite protection from H<sub>2</sub>O<sub>2</sub>-induced programmed cell death, as reported for *L. donovani* (45). Another function proposed for the mitochondrial peroxiredoxin is to protect the mitochondrial genome from direct or indirect peroxide-mediated damage (108). The same role has been suggested for the mitochondrial nsGPX-A3 of *T. brucei*, for which thymine hydroperoxide was found to be a good substrate (93).

The importance of the mitochondrial 2-Cys peroxiredoxin for trypanosomatids has been investigated in two organisms. In the case of *L. infantum*, enzyme depletion was shown to significantly impair parasite infectivity in an animal model (H. Castro and Ana M. Tomás, unpublished). The situation may be somewhat different in *T. brucei*, wherein downregulation of the homologous enzyme does not seem to affect growth of the mammalian stage of the parasite (112). This could partially be explained by the fact that, unlike *Leishmania* spp. and *T. cruzi*, the mammalian stage of *T. brucei* lacks oxidative phosphorylation (87). Still, low levels of H<sub>2</sub>O<sub>2</sub> are constantly produced (35) and depletion of the mitochondrial peroxiredoxin is possibly complemented by other mechanisms, such as the mitochondrial nsGPX-A redox pathway. For *T. brucei*, the mitochondrial nsGPX-A was alleged to play an important role within the mitochondrion (93), although no definitive evidence for this exists.

## Concluding Remarks

In recent years, the combined effort of worldwide researchers has led to important findings concerning the pathways for peroxide elimination in trypanosomatids. Contradicting early convictions, it is presently recognized that these organisms possess an elaborate peroxide metabolism. This consists of a series of redox cascades, wherein reducing equivalents derived from trypanothione are transferred to enzymes belonging to three major families, (a) 2-cysteine peroxiredoxins, (b) nonselenium glutathione peroxidases, and (c) ascorbate peroxidases. These peroxidases possess different physiological reductants and substrate specificities (Fig. 2). In addition, they are strategically localized in specific parasite compartments, where they play precise and complementary roles. Despite all the advances in this area of research, the peroxidase repertoire of trypanosomatids may not yet be complete. Conclusion of the trypanosomatids genome projects will certainly add important findings to this, in the near future. Also, noteworthy is the possibility of exploring some of the unique characteristics of the peroxide metabolism of trypanosomatids to develop new chemotherapeutic agents for the treatment of these medically important parasites.

## Acknowledgments

This study was supported by Fundação para a Ciência e a Tecnologia (FCT) (grants POCTI/ESP/41982/2001 and POCI/SAU-IMI/59560/2004). H. Castro is recipient of a FCT postdoctoral fellowship (SFRH/BPD/20610/2004).

# Abbreviations

APX, ascorbate peroxidase; ER, endoplasmic reticulum; GPX, glutathione peroxidase; GSH, glutathione; GSPS, glutathionylspermidine synthetase;  $K_M$ , Michaelis–Menten constant; nsGPX, nonselenium glutathione peroxidase; PHGPX, phospholipid hydroperoxide glutathione peroxidase; PRX, peroxiredoxin; SeCys, selenocysteine; TR, trypanothione reductase; TRYS, trypanothione synthetase; TS<sub>2</sub>, trypanothione disulfide; T(SH)<sub>2</sub>, dihydrotrypanothione; TXN, trypanoperoxidase; UMSBP, Universal Minicircle Sequence Binding Protein

# References

1. Acestor N, Masina S, Ives A, Walker J, Saravia NG, and Fasel N. Resistance to oxidative stress is associated with metastasis in mucocutaneous leishmaniasis. *J Infect Dis* 194: 1160–1167, 2006.
2. Adak S and Datta AK. *Leishmania major* encodes an unusual peroxidase that is a close homologue of plant ascorbate peroxidase: a novel role of the transmembrane domain. *Biochem J* 390: 465–474, 2005.
3. Alphey MS, Bond CS, Tetaud E, Fairlamb AH, and Hunter WN. The structure of reduced trypanoperoxidase reveals a decamer and insight into reactivity of 2Cys-peroxiredoxins. *J Mol Biol* 300: 903–916, 2000.
4. Ariyanayagam MR and Fairlamb AH. Ovothioli and trypanothione as antioxidants in trypanosomatids. *Mol Biochem Parasitol* 115: 189–198, 2001.
5. Ariyanayagam MR, Oza SL, Mehlert A, and Fairlamb AH. Bis(glutathionyl)spermine and other novel trypanothione analogues in *Trypanosoma cruzi*. *J Biol Chem* 278: 27612–27619, 2003.
6. Ariyanayagam MR, Oza SL, Guthrie ML, and Fairlamb AH. Phenotypic analysis of trypanothione synthetase knock-down in the African trypanosome. *Biochem J* 391: 425–432, 2005.
7. Banhegyi G, Csala M, Szarka A, Varsanyi M, Benedetti A, and Mandl J. Role of ascorbate in oxidative protein folding. *Biofactors* 17: 37–46, 2003.
8. Barr SD and Gedamu L. Cloning and characterization of three differentially expressed peroxidase genes from *Leishmania chagasi*. Evidence for an enzymatic detoxification of hydroxyl radicals. *J Biol Chem* 276: 34279–34287, 2001.
9. Barr SD and Gedamu L. Role of peroxidases in *Leishmania chagasi* survival. Evidence of an enzymatic defense against nitrosative stress. *J Biol Chem* 278: 10816–10823, 2003.
10. Blos M, Schleicher U, Soares Rocha FJ, Meissner U, Rollinghoff M, and Bogdan C. Organ-specific and stage-dependent control of *Leishmania major* infection by inducible nitric oxide synthase and phagocyte NADPH oxidase. *Eur J Immunol* 33: 1224–1234, 2003.
11. Boveris A, Sies H, Martino EE, Docampo R, Turrens JF, and Stoppani AO. Deficient metabolic utilization of hydrogen peroxide in *Trypanosoma cruzi*. *Biochem J* 188: 643–648, 1980.
12. Brigelius-Flohé R, Aumann KD, Blöcker H, Gross G, Kiess M, Klöppel KD, Maiorino M, Roveri A, Schuckelt R, Ursini F, Wingender E, and Flohé L. Phospholipid-hydroperoxide glutathione peroxidase. Genomic DNA, cDNA, and deduced amino acid sequence. *J Biol Chem* 269: 7342–7348, 1994.
13. Brigelius-Flohé R and Flohé L. Is there a role of glutathione peroxidases in signaling and differentiation? *Biofactors* 17: 93–102, 2003.
14. Budde H, Flohé L, Hecht HJ, Hofmann B, Stehr M, Wissing J, and Lünsdorf H. Kinetics and redox-sensitive oligomerisation reveal negative subunit cooperativity in trypanoperoxidase of *Trypanosoma brucei brucei*. *Biol Chem* 384: 619–633, 2003.
15. Cardoni RL, Antunez MI, Morales C, and Nantes IR. Release of reactive oxygen species by phagocytic cells in response to live parasites in mice infected with *Trypanosoma cruzi*. *Am J Trop Med Hyg* 56: 329–334, 1997.
16. Carnieri EG, Moreno SN, and Docampo R. Trypanothione-dependent peroxide metabolism in *Trypanosoma cruzi* different stages. *Mol Biochem Parasitol* 61: 79–86, 1993.
17. Castro H, Budde H, Flohé L, Hofmann B, Lünsdorf H, Wissing J, and Tomás AM. Specificity and kinetics of a mitochondrial peroxiredoxin of *Leishmania infantum*. *Free Radic Biol Med* 33: 1563–1574, 2002.
18. Castro H, Sousa C, Santos M, Cordeiro-da-Silva A, Flohé L, and Tomás AM. Complementary antioxidant defense by cytoplasmic and mitochondrial peroxiredoxins in *Leishmania infantum*. *Free Radic Biol Med* 33: 1552–1562, 2002.
19. Castro H, Sousa C, Novais M, Santos M, Budde H, Cordeiro-da-Silva A, Flohé L, and Tomás AM. Two linked genes of *Leishmania infantum* encode trypanoperoxidases localised to cytosol and mitochondrion. *Mol Biochem Parasitol* 136: 137–147, 2004.
20. Chang TS, Jeong W, Choi SY, Yu S, Kang SW, and Rhee SG. Regulation of peroxiredoxin I activity by Cdc2-mediated phosphorylation. *J Biol Chem* 277: 25370–25376, 2002.
21. Chang TS, Cho CS, Park S, Yu S, Kang SW, and Rhee SG. Peroxiredoxin III, a mitochondrion-specific peroxidase, regulates apoptotic signaling by mitochondria. *J Biol Chem* 279: 41975–41984, 2004.
22. Chuang MH, Wu MS, Lo WL, Lin JT, Wong CH, and Chiou SH. The antioxidant protein alkylhydroperoxide reductase of *Helicobacter pylori* switches from a peroxide reductase to a molecular chaperone function. *Proc Natl Acad Sci USA* 103: 2552–2557, 2006.
23. Clark D, Albrecht M, and Arevalo J. Ascorbate variations and dehydroascorbate reductase activity in *Trypanosoma cruzi* epimastigotes and trypomastigotes. *Mol Biochem Parasitol* 66: 143–145, 1994.
24. Colasante C, Ellis M, Ruppert T, and Voncken F. Comparative proteomics of glycosomes from bloodstream form and procyclic culture form *Trypanosoma brucei brucei*. *Proteomics* 6: 3275–3293, 2006.
25. Comini M, Menge U, and Flohé L. Biosynthesis of trypanothione in *Trypanosoma brucei brucei*. *Biol Chem* 384: 653–656, 2003.
26. Comini M, Menge U, Wissing J, and Flohé L. Trypanothione synthesis in *Crithidia* revisited. *J Biol Chem* 280: 6850–6860, 2005.
27. Comini MA, Guerrero SA, Haile S, Menge U, Lünsdorf H, and Flohé L. Validation of *Trypanosoma brucei* trypanothione synthetase as drug target. *Free Radic Biol Med* 36: 1289–1302, 2004.
28. Comini MA, Krauth-Siegel RL, and Flohe L. Depletion of the thioredoxin homologue trypanoperoxidase impairs antioxidant defence in African trypanosomes. *Biochem J* 402: 43–49, 2007.
29. Dansen TB and Wirtz KW. The peroxisome in oxidative stress. *IUBMB Life* 51: 223–230, 2001.



30. Docampo R, de Boiso JF, Boveris A, and Stoppani AO. Localization of peroxidase activity in *Trypanosoma cruzi* microbodies. *Experientia* 32: 972–975, 1976.
31. Dormeyer M, Reckenfelderbäumer N, Lüdemann H, and Krauth-Siegel RL. Trypanothione-dependent synthesis of deoxyribonucleotides by *Trypanosoma brucei* ribonucleotide reductase. *J Biol Chem* 276: 10602–10606, 2001.
32. el-Sayed NM, Alarcon CM, Beck JC, Sheffield VC, and Donelson JE. cDNA expressed sequence tags of *Trypanosoma brucei rhodesiense* provide new insights into the biology of the parasite. *Mol Biochem Parasitol* 73: 75–90, 1995.
33. Epp O, Ladenstein R, and Wendel A. The refined structure of the selenoenzyme glutathione peroxidase at 0.2-nm resolution. *Eur J Biochem* 133: 51–69, 1983.
34. Fairlamb AH, Blackburn P, Ulrich P, Chait BT, and Cerami A. Trypanothione: a novel bis(glutathionyl)spermidine cofactor for glutathione reductase in trypanosomatids. *Science* 227: 1485–1487, 1985.
35. Fang J and Beattie DS. Alternative oxidase present in procyclic *Trypanosoma brucei* may act to lower the mitochondrial production of superoxide. *Arch Biochem Biophys* 414: 294–302, 2003.
36. Flohé L, Hecht HJ, and Steinert P. Glutathione and trypanothione in parasitic hydroperoxide metabolism. *Free Radic Biol Med* 27: 966–984, 1999.
37. Flohé L, Budde H, Bruns K, Castro H, Clos J, Hofmann B, Kansal-Kalavar S, Krumme D, Menge U, Plank-Schumacher K, Sztajer H, Wissing J, Wylegalla C, and Hecht HJ. Tryparedoxin peroxidase of *Leishmania donovani*: molecular cloning, heterologous expression, specificity, and catalytic mechanism. *Arch Biochem Biophys* 397: 324–335, 2002.
38. Genestra M, Guedes-Silva D, Souza WJ, Cysne-Finkelstein L, Soares-Bezerra RJ, Monteiro FP, and Leon LL. Nitric oxide synthase (NOS) characterization in *Leishmania amazonensis* axenic amastigotes. *Arch Med Res* 37: 328–333, 2006.
39. Genestra M, Souza WJ, Guedes-Silva D, Machado GM, Cysne-Finkelstein L, Bezerra RJ, Monteiro F, and Leon LL. Nitric oxide biosynthesis by *Leishmania amazonensis* promastigotes containing a high percentage of metacyclic forms. *Arch Microbiol* 185: 348–354, 2006.
40. Gommel DU, Nogoceke E, Morr M, Kiess M, Kalisz HM, and Flohé L. Catalytic characteristics of tryparedoxin. *Eur J Biochem* 248: 913–918, 1997.
41. Grondin K, Haimeur A, Mukhopadhyay R, Rosen BP, and Ouellette M. Co-amplification of the gamma-glutamylcysteine synthetase gene *gsh1* and of the ABC transporter gene *pgpA* in arsenite-resistant *Leishmania tarentolae*. *EMBO J* 16: 3057–3065, 1997.
42. Guerrero SA, Lopez JA, Steinert P, Montemartini M, Kalisz HM, Colli W, Singh M, Alves MJ, and Flohé L. His-tagged tryparedoxin peroxidase of *Trypanosoma cruzi* as a tool for drug screening. *Appl Microbiol Biotechnol* 53: 410–414, 2000.
43. Halliwell B. Vitamin C: poison, prophylactic or panacea? *Trends Biochem Sci* 24: 255–259, 1999.
44. Halliwell B. Vitamin C and genomic stability. *Mutat Res* 475: 29–35, 2001.
45. Harder S, Bente M, Isermann K, and Bruchhaus I. Expression of a mitochondrial peroxiredoxin prevents programmed cell death in *Leishmania donovani*. *Eukaryot Cell* 5: 861–870, 2006.
46. Heby O, Persson L, and Rentala M. Targeting the polyamine biosynthetic enzymes: a promising approach to therapy of African sleeping sickness, Chagas' disease, and leishmaniasis. *Amino Acids* 33: 359–366, 2007.
47. Henderson GB, Fairlamb AH, and Cerami A. Trypanothione dependent peroxide metabolism in *Crithidia fasciculata* and *Trypanosoma brucei*. *Mol Biochem Parasitol* 24: 39–45, 1987.
48. Herbette S, Roeckel-Drevet P, and Drevet JR. Seleno-independent glutathione peroxidases. More than simple antioxidant scavengers. *FEBS J* 274: 2163–2180, 2007.
49. Hillebrand H, Schmidt A, and Krauth-Siegel RL. A second class of peroxidases linked to the trypanothione metabolism. *J Biol Chem* 278: 6809–6815, 2003.
50. Hiner AN, Rodriguez-Lopez JN, Arnao MB, Lloyd Raven E, Garcia-Canovas F, and Acosta M. Kinetic study of the inactivation of ascorbate peroxidase by hydrogen peroxide. *Biochem J* 348: 321–328, 2000.
51. Huynh TT, Huynh VT, Harmon MA, and Phillips MA. Gene knockdown of gamma-glutamylcysteine synthetase by RNAi in the parasitic protozoa *Trypanosoma brucei* demonstrates that it is an essential enzyme. *J Biol Chem* 278: 39794–39800, 2003.
52. Jang HH, Lee KO, Chi YH, Jung BG, Park SK, Park JH, Lee JR, Lee SS, Moon JC, Yun JW, Choi YO, Kim WY, Kang JS, Cheong GW, Yun DJ, Rhee SG, Cho MJ, and Lee SY. Two enzymes in one; two yeast peroxiredoxins display oxidative stress-dependent switching from a peroxidase to a molecular chaperone function. *Cell* 117: 625–635, 2004.
53. Jang HH, Kim SY, Park SK, Jeon HS, Lee YM, Jung JH, Lee SY, Chae HB, Jung YJ, Lee KO, Lim CO, Chung WS, Bahk JD, Yun DJ, Cho MJ, and Lee SY. Phosphorylation and concomitant structural changes in human 2-Cys peroxiredoxin isotype I differentially regulate its peroxidase and molecular chaperone functions. *FEBS Lett* 580: 351–355, 2006.
54. Jirata D, Kuru T, Genetu A, Barr S, Hailu A, Aseffa A, and Gedamu L. Identification, sequencing and expression of peroxidoxin genes from *Leishmania aethiopica*. *Acta Trop* 99: 88–96, 2006.
55. Jönsson TJ and Lowther WT. The peroxiredoxin repair proteins. In: *Peroxiredoxin Systems*, edited by Flohé L and Harris JR. New York: Springer, 2007, pp. 115–141.
56. Kelly JM, Taylor MC, Smith K, Hunter KJ, and Fairlamb AH. Phenotype of recombinant *Leishmania donovani* and *Trypanosoma cruzi* which over-express trypanothione reductase. Sensitivity towards agents that are thought to induce oxidative stress. *Eur J Biochem* 218: 29–37, 1993.
57. Klotz LO, Schroeder P, and Sies H. Peroxynitrite signaling: receptor tyrosine kinases and activation of stress-responsive pathways. *Free Radic Biol Med* 33: 737–743, 2002.
58. Krauth-Siegel RL and Lüdemann H. Reduction of dehydroascorbate by trypanothione. *Mol Biochem Parasitol* 80: 203–208, 1996.
59. Krauth-Siegel RL, Bauer H, and Schirmer RH. Dithiol proteins as guardians of the intracellular redox milieu in parasites: old and new drug targets in trypanosomes and malaria-causing plasmodia. *Angew Chem Int Ed Engl* 44: 690–715, 2005.
60. Krauth-Siegel RL, Comini MA, and Schlecker T. The trypanothione system. In: *Peroxiredoxin Systems*, edited by Flohé L and Harris JR. New York: Springer, 2007, pp. 231–251.
61. Levick MP, Tetaud E, Fairlamb AH, and Blackwell JM. Identification and characterisation of a functional peroxidoxin from *Leishmania major*. *Mol Biochem Parasitol* 96: 125–137, 1998.
62. Lin YC, Hsu JY, Chiang SC, and Lee ST. Distinct overexpression of cytosolic and mitochondrial tryparedoxin per-

- oxidases results in preferential detoxification of different oxidants in arsenite-resistant *Leishmania amazonensis* with and without DNA amplification. *Mol Biochem Parasitol* 142: 66–75, 2005.
63. Linares E, Giorgio S, Mortara RA, Santos CX, Yamada AT, and Augusto O. Role of peroxynitrite in macrophage microbicidal mechanisms *in vivo* revealed by protein nitration and hydroxylation. *Free Radic Biol Med* 30: 1234–1242, 2001.
  64. Logan FJ, Taylor MC, Wilkinson SR, Kaur H, and Kelly JM. The terminal step in vitamin C biosynthesis in *Trypanosoma cruzi* is mediated by a FMN-dependent galactonolactone oxidase. *Biochem J* 407: 419–426, 2007.
  65. Lüdemann H, Dormeyer M, Sticherling C, Stallmann D, Follmann H, and Krauth-Siegel RL. *Trypanosoma brucei* trypanoperoxidase, a thioredoxin-like protein in African trypanosomes. *FEBS Lett* 431: 381–385, 1998.
  66. Maiorino M, Ursini F, Bosello V, Toppo S, Tosatto SC, Mauri P, Becker K, Roveri A, Bulato C, Benazzi L, De Palma A, and Flohé L. The thioredoxin specificity of *Drosophila* GPx: a paradigm for a peroxiredoxin-like mechanism of many glutathione peroxidases. *J Mol Biol* 365: 1033–1046, 2007.
  67. Mandal G, Wyllie S, Singh N, Sundar S, Fairlamb AH, and Chatterjee M. Increased levels of thiols protect antimony unresponsive *Leishmania donovani* field isolates against reactive oxygen species generated by trivalent antimony. *Parasitology* 134: 1679–1687, 2007.
  68. Mehta A and Shaha C. Mechanism of metalloid-induced death in *Leishmania* spp.: role of iron, reactive oxygen species,  $\text{Ca}^{2+}$ , and glutathione. *Free Radic Biol Med* 40: 1857–1868, 2006.
  69. Michels PA, Bringaud F, Herman M, and Hannaert V. Metabolic functions of glycosomes in trypanosomatids. *Biochim Biophys Acta* 1763: 1463–1477, 2006.
  70. Monod J, Wyman J, and Changeux JP. On the nature of allosteric transitions: a plausible model. *J Mol Biol* 12: 88–118, 1965.
  71. Moon JC, Hah YS, Kim WY, Jung BG, Jang HH, Lee JR, Kim SY, Lee YM, Jeon MG, Kim CW, Cho MJ, and Lee SY. Oxidative stress-dependent structural and functional switching of a human 2-Cys peroxiredoxin isotype II that enhances HeLa cell resistance to  $\text{H}_2\text{O}_2$ -induced cell death. *J Biol Chem* 280: 28775–28784, 2005.
  72. Motyka SA, Drew ME, Yildirim G, and Englund PT. Overexpression of a cytochrome b5 reductase-like protein causes kinetoplast DNA loss in *Trypanosoma brucei*. *J Biol Chem* 281: 18499–18506, 2006.
  73. Moutiez M, Aumercier M, Schoneck R, Meziane-Cherif D, Lucas V, Aumercier P, Ouassii A, Sergheraert C, and Tartar A. Purification and characterization of a trypanothione-glutathione thioltransferase from *Trypanosoma cruzi*. *Biochem J* 310: 433–437, 1995.
  74. Moutiez M, Quemeneur E, Sergheraert C, Lucas V, Tartar A, and Davioud-Charvet E. Glutathione-dependent activities of *Trypanosoma cruzi* p52 makes it a new member of the thiol:disulphide oxidoreductase family. *Biochem J* 322: 43–48, 1997.
  75. Müller S, Liebau E, Walter RD, and Krauth-Siegel RL. Thiol-based redox metabolism of protozoan parasites. *Trends Parasitol* 19: 320–328, 2003.
  76. Nogoceke E, Gommel DU, Kiess M, Kalisz HM, and Flohé L. A unique cascade of oxidoreductases catalyses trypanothione-mediated peroxide metabolism in *Crithidia fasciculata*. *Biol Chem* 378: 827–836, 1997.
  77. Onn I, Milman-Shtepel N, and Shlomai J. Redox potential regulates binding of universal minicircle sequence binding protein at the kinetoplast DNA replication origin. *Eukaryot Cell* 3: 277–287, 2004.
  78. Oza SL, Tetaud E, Ariyanayagam MR, Warnon SS, and Fairlamb AH. A single enzyme catalyses formation of Trypanothione from glutathione and spermidine in *Trypanosoma cruzi*. *J Biol Chem* 277: 35853–35861, 2002.
  79. Oza SL, Shaw MP, Wyllie S, and Fairlamb AH. Trypanothione biosynthesis in *Leishmania major*. *Mol Biochem Parasitol* 139: 107–116, 2005.
  80. Patterson WR and Poulos TL. Characterization and crystallization of recombinant pea cytosolic ascorbate peroxidase. *J Biol Chem* 269: 17020–17024, 1994.
  81. Paveto C, Pereira C, Espinosa J, Montagna AE, Farber M, Esteva M, Flawia MM, and Torres HN. The nitric oxide transduction pathway in *Trypanosoma cruzi*. *J Biol Chem* 270: 16576–16579, 1995.
  82. Penketh PG and Klein RA. Hydrogen peroxide metabolism in *Trypanosoma brucei*. *Mol Biochem Parasitol* 20: 111–121, 1986.
  83. Penketh PG, Kennedy WP, Patton CL, and Sartorelli AC. Trypanosomatid hydrogen peroxide [corrected] metabolism. *FEBS Lett* 221: 427–431, 1987.
  84. Piacenza L, Peluffo G, Alvarez MN, Kelly JM, Wilkinson SR, and Radi R. Peroxiredoxins play a major role in protecting *Trypanosoma cruzi* against macrophage- and endogenously-derived peroxynitrite. *Biochem J* 410: 359–368, 2008.
  85. Piñeyro MD, Pizarro JC, Lema F, Pritsch O, Cayota A, Bentley GA, and Robello C. Crystal structure of the trypanoperoxidase from the human parasite *Trypanosoma cruzi*. *J Struct Biol* 150: 11–22, 2005.
  86. Poole LB. The catalytic mechanism of peroxiredoxins. In: *Peroxiredoxin Systems*, edited by Flohé L and Harris JR. New York: Springer, 2007, pp. 66–81.
  87. Priest JW and Hajduk SL. Developmental regulation of mitochondrial biogenesis in *Trypanosoma brucei*. *J Bioenerg Biomembr* 26: 179–191, 1994.
  88. Rais S, Perianin A, Lenoir M, Sadak A, Rivollet D, Paul M, and Deniau M. Sodium stibogluconate (Pentostam) potentiates oxidant production in murine visceral leishmaniasis and in human blood. *Antimicrob Agents Chemother* 44: 2406–2410, 2000.
  89. Raven EL. Understanding functional diversity and substrate specificity in haem peroxidases: what can we learn from ascorbate peroxidase? *Nat Prod Rep* 20: 367–381, 2003.
  90. Reckenfelderbäumer N and Krauth-Siegel RL. Catalytic properties, thiol pK value, and redox potential of *Trypanosoma brucei* trypanoperoxidase. *J Biol Chem* 277: 17548–17555, 2002.
  91. Rhee SG, Chae HZ, and Kim K. Peroxiredoxins: a historical overview and speculative preview of novel mechanisms and emerging concepts in cell signaling. *Free Radic Biol Med* 38: 1543–1552, 2005.
  92. Rhee SG, Kang SW, Jeong W, Chang TS, Yang KS, and Woo HA. Intracellular messenger function of hydrogen peroxide and its regulation by peroxiredoxins. *Curr Opin Cell Biol* 17: 183–189, 2005.
  93. Schlecker T, Schmidt A, Dirdjaja N, Voncken F, Clayton C, and Krauth-Siegel RL. Substrate specificity, localization, and essential role of the glutathione peroxidase-type trypanoperoxidase in *Trypanosoma brucei*. *J Biol Chem* 280: 14385–14394, 2005.

94. Schlecker T, Comini MA, Melchers J, Ruppert T, and Krauth-Siegel RL. Catalytic mechanism of the glutathione peroxidase-type trypanodoxin peroxidase of *Trypanosoma brucei*. *Biochem J* 405: 445–454, 2007.
95. Schmidt H and Krauth-Siegel RL. Functional and physico-chemical characterization of the thioredoxin system in *Trypanosoma brucei*. *J Biol Chem* 278: 46329–46336, 2003.
96. Sharp KH, Moody PCE, and Raven EL. A new framework for understanding substrate binding and functional diversity in haem peroxidases. *Dalton Trans*: 4208–4215, 2003.
97. Steenkamp DJ. Trypanosomal antioxidants and emerging aspects of redox regulation in the trypanosomatids. *Antioxid Redox Signal* 4: 105–121, 2002.
98. Temperton NJ, Wilkinson SR, and Kelly JM. Cloning of an Fe-superoxide dismutase gene homologue from *Trypanosoma cruzi*. *Mol Biochem Parasitol* 76: 339–343, 1996.
99. Tetaud E, Giroud C, Prescott AR, Parkin DW, Baltz D, Biteau N, Baltz T, and Fairlamb AH. Molecular characterisation of mitochondrial and cytosolic trypanothione-dependent trypanodoxin peroxidases in *Trypanosoma brucei*. *Mol Biochem Parasitol* 116: 171–183, 2001.
100. Trujillo M, Budde H, Piñeyro MD, Stehr M, Robello C, Flohé L, and Radi R. *Trypanosoma brucei* and *Trypanosoma cruzi* trypanodoxin peroxidases catalytically detoxify peroxynitrite via oxidation of fast reacting thiols. *J Biol Chem* 279: 34175–34182, 2004.
101. Ullrich V and Kissner R. Redox signaling: bioinorganic chemistry at its best. *J Inorg Biochem* 100: 2079–2086, 2006.
102. Vickers TJ and Fairlamb AH. Trypanothione S-transferase activity in a trypanosomatid ribosomal elongation factor 1B. *J Biol Chem* 279: 27246–27256, 2004.
103. Vickers TJ, Wyllie S, and Fairlamb AH. *Leishmania major* elongation factor 1B complex has trypanothione S-transferase and peroxidase activity. *J Biol Chem* 279: 49003–49009, 2004.
104. Viode C, Bettache N, Cenas N, Krauth-Siegel RL, Chauviere G, Bakalara N, and Perie J. Enzymatic reduction studies of nitroheterocycles. *Biochem Pharmacol* 57: 549–557, 1999.
105. Vogt RN and Steenkamp DJ. The metabolism of S-nitrosothiols in the trypanosomatids: the role of ovoid thiol A and trypanothione. *Biochem J* 371: 49–59, 2003.
106. Walker J, Acestor N, Gongora R, Quadroni M, Segura I, Fasel N, and Saravia NG. Comparative protein profiling identifies elongation factor-1 $\beta$  and trypanodoxin peroxidase as factors associated with metastasis in *Leishmania guyanensis*. *Mol Biochem Parasitol* 145: 254–264, 2006.
107. Wilkinson SR, Meyer DJ, and Kelly JM. Biochemical characterization of a trypanosome enzyme with glutathione-dependent peroxidase activity. *Biochem J* 352: 755–761, 2000.
108. Wilkinson SR, Temperton NJ, Mondragon A, and Kelly JM. Distinct mitochondrial and cytosolic enzymes mediate trypanothione-dependent peroxide metabolism in *Trypanosoma cruzi*. *J Biol Chem* 275: 8220–8225, 2000.
109. Wilkinson SR, Meyer DJ, Taylor MC, Bromley EV, Miles MA, and Kelly JM. The *Trypanosoma cruzi* enzyme TcGPXI is a glycosomal peroxidase and can be linked to trypanothione reduction by glutathione or trypanodoxin. *J Biol Chem* 277: 17062–17071, 2002.
110. Wilkinson SR, Obado SO, Mauricio IL, and Kelly JM. *Trypanosoma cruzi* expresses a plant-like ascorbate-dependent hemoperoxidase localized to the endoplasmic reticulum. *Proc Natl Acad Sci USA* 99: 13453–13458, 2002.
111. Wilkinson SR, Taylor MC, Touitha S, Mauricio IL, Meyer DJ, and Kelly JM. TcGPXII, a glutathione-dependent *Trypanosoma cruzi* peroxidase with substrate specificity restricted to fatty acid and phospholipid hydroperoxides, is localized to the endoplasmic reticulum. *Biochem J* 364: 787–794, 2002.
112. Wilkinson SR, Horn D, Prathalingam SR, and Kelly JM. RNA interference identifies two hydroperoxide metabolizing enzymes that are essential to the bloodstream form of the african trypanosome. *J Biol Chem* 278: 31640–31646, 2003.
113. Wilkinson SR, Prathalingam SR, Taylor MC, Horn D, and Kelly JM. Vitamin C biosynthesis in trypanosomes: a role for the glycosome. *Proc Natl Acad Sci USA* 102: 11645–11650, 2005.
114. Wilkinson SR, Prathalingam SR, Taylor MC, Ahmed A, Horn D, and Kelly JM. Functional characterisation of the iron superoxide dismutase gene repertoire in *Trypanosoma brucei*. *Free Radic Biol Med* 40: 198–209, 2006.
115. Wood ZA, Poole LB, and Karplus PA. Peroxiredoxin evolution and the regulation of hydrogen peroxide signaling. *Science* 300: 650–653, 2003.
116. Wood ZA, Schröder E, Harris RJ, and Poole LB. Structure, mechanism and regulation of peroxiredoxins. *Trends Biochem Sci* 28: 32–40, 2003.
117. Wyllie S, Cunningham ML, and Fairlamb AH. Dual action of antimonial drugs on thiol redox metabolism in the human pathogen *Leishmania donovani*. *J Biol Chem* 279: 39925–39932, 2004.

Address reprint requests to:

Ana M. Tomás  
IBMC  
Rua do Campo Alegre, 823  
150-180 Porto, Portugal

E-mail: atomas@ibmc.up.pt

Date of first submission to ARS Central, February 20, 2008; date of acceptance, March 6, 2008.