

Africa's Tropical Forests

Francis J. Adair, 2025



Photo 1: Evidence of logging along the Mpivié river.

Contents

Introduction.....	3
Part 1 — Forests and Gorillas.....	4
Central Africa.....	4
Paleo-history.....	4
Gorillas.....	5
Western.....	5
Eastern.....	5
Habitat.....	6
Introduction of Humans.....	7
Gorilla Extinction.....	7
Threat of Humans.....	8
Part 2 — Forests and Climate.....	8
Limiting Factors.....	9
Atmospheric Effects.....	10
Carbon Dioxide.....	11
Forest Loss.....	12
Logging.....	12
Part 3 — People of the Forest.....	14
Forest Resources.....	14
Adaptation and Mitigation.....	15
Mitigation Mechanisms.....	16
Adaptation Strategy.....	16
Governance & Tourism (Gabon & Cameroon).....	17
Viability of Ecotourism.....	18
Part 4 — Ecosystem Conservation.....	20
Classifying Spatial Costs.....	21
Migratory Response to Climate Change.....	21
Protected Areas and Forest Conservation.....	21
Evaluating Costs.....	22
Habitat Fragmentation.....	23
Range Expansion (Primates).....	23
Technology.....	23
Wildlife.....	24
Forests.....	24
Local Context.....	24
Summary.....	25
References.....	26

Introduction

In 2018 I went on an expedition to the Loango coastal forest region of southwestern Gabon as a voluntary field agent under the auspices of Stuart Nixon, Chester Zoo's Regional Field Programme Manager. I would like to thank Stuart and Charlie Childs for taking care of me on that expedition and everyone else I met who inspired and supported me while I wrote this review. The contents are split into four parts to understand how forest conservation is tied to wildlife, climate, and human existence in Central Africa.

- Part 1 - The history of forests and gorillas in equatorial Africa to the modern day.
- Part 2 - Tropical forest interactions with a changing climate, hydrology, and humans.
- Part 3 - Governments, peoples, and communities utilisation and valuation of forests.
- Part 4 - Complexities, methods (e.g. ecosystem services), and contexts for conservation.



Photo 2: Adult antlion in a grassland patch of secondary rainforest mosaic near Fernan Vaz lagoon.

Part 1 — Forests and Gorillas

Central Africa

The Central Africa region includes the countries of Cameroon, Gabon, Equatorial Guinea, the Democratic Republic of the Congo (DRC), the Republic of Congo (Congo), the Central African Republic (CAR), and the Angolan exclave of Cabinda (Abernethy et al., 2016). The region covers around 2.28 million km² (Hoare, 2007) and includes parts of central and southern Cameroon, the CAR, Republic of Congo, Gabon, Equatorial Guinea, and the DRC (Kimbu, 2011). Dense rainforest presides across this equatorial belt region but not equally (White, 2001a; Abernethy et al., 2016) as we shall see. The Congolian rainforests extend from the eastern Albertine Rift Valley to the western coast and include areas in DRC, southern CAR, Congo, and Cabinda (Abernethy et al., 2016).

As for forests, the Congo Basin contains 1.8 million km² of forest that reaches across DRC, most of Congo, the southeastern parts of Cameroon, the southern CAR, as well as Gabon and Equatorial Guinea (Nogherotto et al., 2013). The basin is home to the second largest (Kleinschroth et al., 2019) and most intact tropical rainforest after the Amazon (Nkem et al., 2013) and represents approximately 20% of the world's remaining tropical forest (Mayaux et al., 2004). The most extensive stretch of rainforest encompasses the Congo Basin, the adjoining lower Guinean region, and smaller block in West Africa (White, 2001a).

To the west of the Congo River's drainage, the contiguous Lower Guinean forests extend east and south from the Cross River through southern Cameroon, Gabon, and Equatorial Guinea (Abernethy et al., 2016). These forests are collectively referred to as the Guineo-Congolian rainforests but are biologically distinct from the West African rainforests that stretch from southern Senegal to the Dahomey Gap (White, 1985). Additionally, further distinct swamp forests grow along the south Nigerian coasts and Niger River Delta that span the coast from the Dahomey Gap to the Cross River (Olson et al., 2001) (Figure 1).

Paleo-history

The world transitioned from the Mesozoic to the Cenozoic Era 66 million years before-present (Ma YBP) and we are now in the seventh epoch (Cohen et al., 2023). Starting from the Paleocene (66.00 Ma YBP), through to the Eocene (56 Ma YBP), Oligocene (33.9 Ma YBP), Miocene (23.03 Ma YBP), Pliocene (5.333 Ma YBP), Pleistocene (2.58 Ma YBP), up to our present Holocene (11,700 YBP) epoch (Cohen et al., 2023). During the Eocene epoch about 50 Ma YBP, CO₂ rose to over 1000 ppm and it was the last time forests reached their greatest extent (Willis & McElwain, 2014), then by the Oligocene CO₂ had fallen and grasses began replacing forests (Bond & Midgley, 2012). During the same time, C4 type grasses evolved that, unlike their ancestral C3 pathway, grow well in conditions of high heat, drought, salinity (Sage, 2004), sunlight, and fire frequency (Ratnam et al., 2011).

The Last Glacial Maximum (LGM) is the last time ice sheets reached their greatest extent and lasted until around 10,000 YBP (Maley, 1996). During the LGM equatorial Africa experienced a significant drop in temperature (Maley, 2001), approximately 4 °C lower than current levels (Bonnefille et al., 1990; Tierney et al., 2008) and CO₂ was much lower than during interglacial periods (Bond & Midgley, 2012). Pollen studies from Lake Bosumtwi in Ghana and Lake Barombi Mbo in West Cameroon suggest the LGM occurred between 20,000 and 15,000 YBP (e.g. Maley 1989, 1991) (Maley, 1996). But upon different evidence it is posited that the LGM may have occurred anywhere between 26,500 to 19,000 YBP (Clark et al., 2009a) or 18,000 to 12,000 YBP (Maley, 2001). In addition to lower temperatures during the LGM, monsoon circulation was weakened and moisture transport from the Atlantic Ocean was diminished (Gasse, 2000; Schefuß et al., 2005; Tierney et al., 2008). These effects collectively led to a decrease in rainfall (Bonnefille and Chalieu, 2000) and a subsequent increase in aridity (Hamilton, 1982), that in turn caused forests to reduce in extent (Maley, 1996) [see *Part 2 - Forests and Climate*].

When there is a reduction in forest extent grasses move in to colonise, and with a smaller leaf area index than trees, evapotranspiration decreases and causes surface albedo to increase (Nogherotto et al., 2013) (i.e. increased radiation reflectance). Evergreen rainforests seemingly expand during wet intervals and shrink during periods of drought (Ngomanda et al., 2007). Moreover, today's Central African forests originate from islands of 'refugia' that managed to persist throughout the LGM (Maley, 1996, 2001; Willis et al., 2013; Städele et al., 2022). Therefore, the oldest forests are located within former refugia dating to over 18,000 YBP (Abernethy et al., 2016) that include the massive swamps in central DRC and the mountain ranges of the western equatorial forests (Abernethy et al., 2016). Conversely, to complete the picture, the youngest forests are only some decades old and can be found hundreds of kilometres north and south of the refugia, transitioning to and engulfing savannahs (White, 2001b).

Gorillas

Looking more closely at gorillas, Western gorillas are genetically and morphologically distinct from gorillas found 1000 kilometres to the east with two named species in each group; as supported by a comprehensive literature review spanning 85 years (Tocheri et al., 2016). It is estimated that western and eastern gorillas diverged only 1.2 to 3.0 Ma YBP (Thalmann et al., 2007; Becquet and Przeworski, 2007) with gene flow continuing up to around 200,000 to 80,000 YBP (Thalmann et al., 2007), or possibly as recently as 20,000 YBP (Becquet & Przeworski, 2007; Mailund et al., 2012; McManus et al., 2015; Scally et al., 2012; Thalmann et al., 2007; Xue et al., 2015), indicating a gradual separation (Städele et al., 2022). Interestingly, genetic studies of western gorillas exhibit significantly larger past effective population sizes, greater genetic diversity, and a more pronounced population genetic substructure compared to their eastern counterparts (Thalmann et al., 2007, 2011; Scally et al., 2012, 2013; Prado-Martinez et al., 2013; Roy et al., 2014a; Xue et al., 2015; Fünfstück and Vigilant, 2015).

Western

In western Africa, **Western lowland gorillas** (*Gorilla gorilla gorilla*) span western central Sub-Saharan Africa and have the largest geographical distribution of all gorilla subspecies, with an estimated population size of 362,000 individuals (Strindberg et al., 2018). **Cross River gorillas** (*G. g. diehli*) are another western gorilla subspecies that inhabit a highly fragmented and confined area straddling the border between Cameroon and Nigeria (Städele et al., 2022), with the smallest estimated population size of the gorilla species ranging between 200 to 300 (Dunn et al. 2014).

The limited and fragmented Cross River gorilla habitats likely contribute to reduced genetic diversity and greater differentiation (propagation of certain traits within the population) due to limited gene flow and mating opportunities (Städele et al., 2022). Cross River gorillas diverged from Western lowland gorillas between 17,800 (Thalmann et al., 2011) and 68,000 YBP (McManus et al., 2015; Prado-Martinez et al., 2013; Thalmann et al., 2011) during a long cold period that lasted from 75,000 to 10,000 YBP during which forest regression was significant (Maley, 1996). However, gene flow between the two groups did not stop till approximately 420 YBP and was followed by a genetic bottleneck in Cross River gorillas just 320 YBP (Thalmann et al., 2011).

Eastern

In eastern Africa, **Mountain gorillas** (*G. beringei beringei*) are only found in two small, discrete populations, totalling around 1,000 individuals in the Virunga Massif spanning Rwanda, Uganda, and the DRC and including Bwindi Impenetrable National Park in Uganda (Granjon et al., 2020). **Grauer or Eastern lowland gorillas** (*G. b. graueri*) are located in the eastern DRC and they have the second-largest spatial distribution among the gorilla subspecies with an estimated population of approximately 6,800 individuals (Plumptre et al., 2021). Grauers are thought to have split from mountain gorillas between 20,000 (Jensen-Seaman & Kidd, 2001; Roy et al., 2014a) and 10,000 YBP (Roy et al., 2014a).

Mountain gorillas, along with some Grauer gorilla populations inhabit high-altitude areas characterised by afro-montane forests (Mehlman, 2008). Lowland variety Grauers consume a larger amount of fruit and climb more often, with more extensive and seasonally variable ranging patterns aligning more closely with western lowland gorillas (Casimir, 1975; Goodall and Groves, 1977; Yamagiwa et al., 1992; Yamagiwa and Mwanza, 1994; Uchida, 1996; Yamagiwa et al., 1996; Doran and McNeillage, 1998, 2001; Robbins and McNeillage, 2003). Yet, despite having a larger distribution than mountain gorillas, Grauer's genetic diversity and differentiation still suffers from potential geographic and environmental barriers (Städele et al., 2022).



Photo 3: Long-legged katydid in logged primary rainforest along the Mpivié river.

Habitat

Gorillas diverged from the evolutionary line leading to modern humans, chimpanzees, and bonobos only between 12.0-8.8 Ma YBP (Sclay et al., 2012). The middle Miocene occurred 15.98 Ma YBP (Cohen et al., 2023) and during this time East Africa's dense forest cover gave way to a mosaic landscape of forest and savannah (e.g. Van Zinderen Bakker & Mercer 1986; Hamilton 1991; Harris 1993) (Maley, 1996). The upper Miocene started 11.63 Ma YBP (Cohen et al., 2023) as indicated by the introduction of Asian and European flora and fauna to Africa 10.5 Ma YBP, whereby the African climatic zonation began to resemble present-day patterns (Maley, 1996). During the upper Miocene about 8.0 Ma YBP, C4 grasses became very prominent (Cerling et al., 1997) possibly facilitated by low CO₂ levels (Ehleringer et al., 1997; Osborne, 2008; Edwards et al., 2010) such as which occurred in East Africa during the last glaciation (Wooller et al., 2000; Damsté et al., 2011).

Fast-forward to the LGM and the landscape of the highlands along the Albertine Rift is transformed with an arid glacial climate and lower atmospheric CO₂ concentrations that depress montane forest belts by over 1,500 metres (Hamilton, 1982). That led to the advancement of glaciers in the Rwenzori mountains to elevations as low as 2,000 metres above sea level, 2,700 metres lower than today (Livingstone, 1967; Kelly et al., 2014). On East African mountains such as the Rwenzori, these LGM conditions allowed for alpine grasslands to become the dominant vegetation type until about 15,000 YBP (Livingstone, 1967) when the early deglacial period began (Maley, 1996). These cold and arid conditions in east Africa during the LGM made gorilla habitation largely unsuitable with active populations being confined to refugia; as demonstrated in an extensive literature review (Tocheri et al., 2016).

The African Humid Period (AHP) then lasted from 15,000-5,000 years YBP (de Menocal et al., 2000) or possibly 11,000-4,000 YBP (Willis et al., 2013). The AHP brought with it increased temperatures and precipitation, punctuated by the near-glacial Younger Dryas period between 12,800-11,700 YBP (Zhang et al., 2014). The conclusion of the AHP 5000 YBP (e.g. de Menocal et al., 2000; Tierney et al., 2008) marked a period of rainforest decline and the expansion of grasslands and savannahs in the lowlands (Beuning et al., 1997). In East Africa moderate drying is noted at 6,200-5,500 YBP that abruptly shifts to arid conditions at 4,900-4,500 YBP (Tierney et al., 2010).

Increased precipitation during the AHP (e.g. Tierney et al., 2011; Kageyama et al., 2013) allowed for forests to remerge that likely provided the necessary habitat for Grauers to split from mountain gorillas either just before or just after the Younger Dryas period (Tocheri et al., 2016). The AHP reached its maximum at 8,500 YBP and during this time tropical forest species are noted as reaching up into the Sahara/ Sahelian by growing along watercourses (Watrin et al., 2009). The proto-Grauer may have dispersed westward from the Virunga forests alongside an expanding forest habitat into what is now known as the grauer gorilla range, located in the highlands northwest of Lake Kivu (Tocheri et al., 2016).

The late Holocene then starts 4,200 YBP to the present day (Cohen et al., 2023) during which rainforests across western equatorial Africa are then characterised by successive millennial-scale changes (Maley & Brenac, 1998; Vincens et al., 1998; Giresse et al., 1994, 2005; Delègue et al., 2001; Wirmann et al., 2001; Nguetsop et al., 2004) driven by natural climatic variability (Vincens et al., 1999; Elenga et al., 2004). Increasing aridity from 4,000 to 1,300 YBP caused a further reduction in African lowland rainforests (Ngomanda et al., 2007) that largely resulted in the modern vegetational landscape (Vincens et al., 1999).

Tropical rainforests extended their range 1,200 to 800 YBP that may be attributed to increased moisture during the "Medieval Warm Period" (MWP) ~1,100 to 800 YBP (Ngomanda et al., 2007) that unevenly affected parts of the globe (Bradley, 2000; Broecker, 2002). For example, from 3,200 YBP onwards West Africa is noted as undergoing a decrease in wetness and an increase in temperature (Schefuß et al., 2005). Between 1,300 to 600 YBP equatorial west Africa experienced low water levels that may be due to decreased rainfall (e.g. Nguetsop et al., 2004; Wirmann et al., 2001; Giresse et al., 2005) (Ngomanda et al., 2007) during the MWP.

Introduction of Humans

In parts of Central and central West Africa 2,500 YBP an arid event is registered that may have caused forest retreat (Vincens et al., 1999; Maley, 2002) that has also been attributed to human induced land-use change (e.g. Neumann et al., 2012) (Bayon et al., 2012). From 2,700 YBP East African tree-like species have declined and grasses have increased (Vincens, 1993). In Uganda of East Africa 2,000 YBP, a decline in montane tree taxa and rise in grass pollen indicates human-induced deforestation (Livingstone, 1967; Taylor, 1990) and the human clearance of the Virunga forests are documented to have begun around 1,000 YBP (McGlynn et al., 2013).

Over the past 1000 years there has been minor variation in the Guineo-Congolian rainforest cover suggesting that global-scale climatic variability has had little effect (Ngomanda et al., 2007). In fact, various pollen records indicate that vegetation fluctuations in equatorial western Africa during this time are resultant from human activities (Elenga et al., 1994; Reynaud-Farrera et al., 1996; Vincens et al., 1998, 1999). However, the data is too patchy to say with certainty at the decadal or centennial resolution and more archaeological investigations are required (Ngomanda et al., 2007). In my opinion, this is inconsequential to the contemporary evidence explored later.

Gorilla Extinction

Gorillas live in groups where reproduction is dominated by one (a silverback) or a few males (Schaller, 1963; Fossey, 1974; Harcourt, 1979; Robbins, 1995; Watts, 1996; Bradley et al., 2005; Breuer et al., 2010; Masi et al., 2021; Vigilant et al., 2015). For example about half of mountain gorilla social groups have multiple resident adult males but the dominant silverback may sire up to 72% of the offspring (Bradley et al., 2005; Vigilant et al., 2015). High male reproductive skew like that has implications for genetic diversity (range of traits within the population) and differentiation (propagation of certain traits within the population) (Tocheri et al., 2016).

Isolation reduces genetic diversity and increases differentiation (Bradley et al., 2005; Breuer et al., 2010), whereas long-distance male dispersal increases genetic diversity and reduces differentiation (Douadi et al., 2007; Roy et al., 2014b; Masi et al., 2021). Research on gorilla morphology, particularly with grauer gorillas, points toward low levels of genetic diversity among eastern gorilla populations (Ruvolo et al., 1994; Garner and Ryder, 1996; Ruvolo, 1997; Saltonstall et al., 1998; Yu et al., 2004; Prado-Martinez et al., 2013). Low genetic diversity adversely impacts adult survivorship (Xue et al., 2015; Fünfstück & Vigilant, 2015), thereby heightening the risk of extinction for gorillas (Tocheri et al., 2016).

Today, Grauer gorilla habitat is being reduced and fragmented by forest clearance, compounded by bushmeat hunting and the Ebola virus outbreak which have drastically reduced present-day population sizes (Walsh et al., 2003; Plumptre et al., 2016; Strindberg et al., 2018). Ebola has also caused a reduction in the western gorilla population size (Walsh et al., 2003; Leroy et al., 2004; Bermejo et al., 2006; Rizkalla et al., 2007). Currently, western lowland gorillas exhibit the greatest genetic diversity across all measures, but a reduction in genetic diversity can affect individual fitness and survivability (Städle et al., 2022). Of mountain and Grauer's gorillas, 34% and 39% of their genomes respectively, are homozygous, which indicates excessive inbreeding (Xue et al., 2015; van der Valk et al., 2019).

Threat of Humans

Deforestation of gorilla habitat and bushmeat trafficking reduce gorilla populations and exacerbate the risks of extinction for gorillas (Harcourt, 1996; Walsh et al., 2003; Yamagiwa, 2003; McNeilage et al., 2006; Robbins et al., 2011). Rural and urban communities in tropical Africa depend significantly on wild meat for protein, fuelling a growing commercial bushmeat trade (Milner-Gullans & Bennnett, 2003). Bushmeat is a primary source of protein for many households, especially forest dependent people who are traditionally not engaged with animal husbandry (Nasi et al., 2011). Grauers, western lowland, and Cross River gorillas are already classified as Critically Endangered, and mountain gorillas as Endangered (Maisels et al., 2018; Plumptre et al., 2019).

Commercial hunting in Central Africa has significantly impacted wildlife populations, such as with the loss of 62% of Central Africa's elephants from 2002 to 2011 (Maisels et al., 2013). With products like ivory becoming scarcer, hunters shift their focus to other species to remain profitable (Fa & Brown, 2009). In the Congo basin, 60% of 57 studied mammal species are being exploited unsustainably (Fa et al., 2002). In West Africa, persistent overhunting has nearly caused the bushmeat trade to collapse, with some primate species facing likely extinction (Oates et al., 2000). Hunters in forest communities generally do not hunt beyond a 10 km radius from a village (Van Vliet, 2010) but as increasing trade opportunities emerge they are being driven to extend further and exert an increasing pressure into the forest (Kümpel et al., 2010). These complexities will be explored further in Part 3.

Part 2 — Forests and Climate

Climate change is arguably the most persistent threat to global stability in the coming century (Adger et al., 2003). Human influence has unequivocally warmed the planet with global surface temperature increasing since 1970 faster than any other 50-year period in at least 2000 years (Allan et al., 2023). The IPCC's Fifth Assessment Report has identified Representative Concentration Pathways (RCPs) to evaluate projected emission scenarios (Lyon et al., 2022). RCP 2.6 is the most low-impact scenario (Stocker et al., 2013) predicting low-emissions into the future (Lyon et al., 2022) and RCP 8.5 is a high-emission trajectory (Van Vuuren et al., 2011). RCP 6.0 falls between these two extremes with global temperatures predicted to increase from 2.2 °C to 3.6 °C between 2100 and 2200 (Lyon et al., 2022).

In tropical rainforests temperatures are expected to increase by 2-5 °C over this century (Hulme et al., 2001; Giorgi, 2002; Cramer et al., 2004). Global warming last reached 5 °C between 11,700-11,500 YBP (Kobashi et al., 2008) with Greenland ice cores even suggesting a 20-60 year period of 10 °C of warming during that time (Steffensen et al., 2008). Already in the last decade [2011-2020] temperatures have exceeded the most recent multi-century warm period of around 6,500 YBP (Allan et al., 2023). Africa is predicted to warm at a faster rate than the global average (James & Washington, 2013), so even under RCP 2.6 overall warming across Central Africa may become over 3 °C by the end of the century (United Nations, 2015).

Empirically observed climatic trends suggest an increased seasonality and heightened water stress across the northern and western Lower Guinean forests that may alter both forest function and extent (Abernethy et al., 2016). For the central, southern, and eastern forests of the Congo Basin, the situation is less clear, but eastern DRC is expected to experience a wetter climate (Zelazowski et al., 2011). A strong drying trend has been observed in the northern Congo, whereas the southern Congo has seen less of a reduction in rainfall (Zhou et al., 2014). Additionally, West Africa has been subjected to two decades of persistent drought and seen a shift towards drier conditions (Nicholson, 1994; Fauset et al., 2012) despite increases in monsoon precipitation since 1980 (Allan et al., 2023).



Photo 4: Thread-waisted wasp (*Genus Chlorion*) on Fernan Vaz lagoon's banks.

The Congolese forest is drier than the rainforests of Amazonia and may be more tolerant of short-term rainfall reduction than wetter tropical forests (Asefi-Najafabady & Saatchi, 2013). Also, Certain Central African trees possess a higher resilience compared to their neotropical (New World) counterparts that is attributed to their history of enduring post-Pleistocene epoch drought periods (Willis et al., 2013) (e.g. Vincens et al., 1999). For example, rainforest soils such as those of the Lake Nguène catchment in Gabon can be hydromorphic, which may have allowed them to survive through decadal wet-dry episodes (Ngomanda et al., 2007). A relatively high proportion of large, old evergreen trees with high wood density add to forest resilience, but may also be most vulnerable to an ecological tipping point resulting from physiological stress (Lindenmayer et al., 2012).

Drying may be associated with forest retreat (Malhi et al., 2008), particularly at 4°C (James et al., 2013) and especially since drier conditions heighten the risk of forest-fires (Nepstad et al., 2004; Randerson et al., 2005; Poulter et al., 2010). The rainforests of central Africa have been experiencing long-term drying conditions (Malhi & Wright, 2004; Asefi-Najafabady & Saatchi, 2013) and northern African tropics have been experiencing rapid warming since at least 1970 (Malhi & Wright, 2004). There has been a persistent browning in the Congolese forests that may be a reflection of a gradual adjustment to long-term drying, rather than the sporadic droughts seen in the Amazon (Zhou et al., 2014). A strong drying trend has also been observed in the central (Malhi & Wright, 2004; Asefi-Najafabady & Saatchi, 2013), western (Malhi & Wright, 2004; James et al., 2013), and northern African tropics, despite an overall increase of precipitation in East Africa and globally (James et al., 2013).

Limiting Factors

Soil water is the most important factor determining the distribution of forest, savannah, and grassland but is depleted through transpiration (Tinley, 1982). It is over timescales of thousands to tens of thousands of years that soil generally improves in suitability for supporting plant growth (Van Breemen, 1993). The soil parent material influences vegetation and soil chemistry more so than in-situ tree species do (Augusto et al., 2003). That said, the distribution of trees in a forest can have a large bearing on the amelioration, or degradation of its soil (Zinke, 1961; Binkley & Giardina, 1998; Augusto et al., 2003; Frelich et al., 2015). In deforested land both soil moisture and evapotranspiration decline, after which the soil is less able to extract and use moisture (Nogherotto et al., 2013). Highly flammable C4 grasses are the first to colonise cleared forests and recurrent grass-fuelled fires can strongly suppress tree regrowth (Bond et al., 2003). Grasses typically have a higher surface albedo than forests from retaining dead reflective leaves, but transpire less water due to their smaller roots, leaf area, and canopy roughness (Bond & Midgley, 2012).

Increasing photosynthetic active radiation (energy) can enhance radiation-limited canopy photosynthesis (Nemani et al., 2003; Huete et al., 2006) and evapotranspiration, while energy remains the limiting factor (Woodward, 2008; Zhou et al., 2014). But, when soil moisture becomes the limiting factor, increasing energy may enhance water stress for plant growth (Zhou et al., 2014). Water stress results in less turgid (wilting) leaves, to their shedding altogether, that in the short term is recoverable once rainfall returns (Zhou et al., 2014). Thinning and/ or retreating forests then reduce evapotranspiration and increase run-off, resulting in decreases to precipitation rates (James et al., 2013). This may be accelerated as higher temperatures increase evapotranspiration (Woodward, 2008) and decrease productivity (Delire et al., 2008). Long-term ground observations and drought manipulative experiments are needed to understand the impacts of climate change on forests (Nepstad et al., 2007) but research of this kind is not available for Congolese forests (Zhou et al., 2014).

Water deficit over an extended period of time may alter forest species composition and structure from a gradual loss of photosynthetic capacity and water content (Lewis, 2006; Enquist & Enquist, 2011; Fauset, 2012) that may have knock-on effects for biodiversity and carbon storage (Lewis, 2006; Valentini et al., 2014). Long-term drought can reach a critically low threshold of water availability whereby closed-canopy forests transition to more open, lower-biomass forests (Lewis, 2006; Brncic et al., 2007; Enquist & Enquist, 2011; Fauset et al., 2012). A strong drying trend may also cause the photosynthetic capacity of trees and their moisture content to be reduced, which could favour the spread of drought-tolerant species (Lewis, 2006; Delire et al., 2008; Enquist & Enquist, 2011). As seen in Ghanaian forest species shifting from wetter evergreen trees towards drier deciduous species (Fauset et al., 2012). Once open land is established, higher rainfall promotes the growth of wooded savannah (Sankaran et al., 2005) instead of forests, which differ in their species composition, structure, and function (Ratnam et al., 2011).

Atmospheric Effects

Forests have a low surface albedo and so contribute to planetary warming through increased heat absorption on land, but they also cool the climate through evapotranspiration as part of the hydrologic cycle (Bonan, 2008). The level of radiation required for plant growth and transpiration is determined by atmospheric conditions (clouds and aerosols) and latitude (Zhou et al., 2014). In this instance, surface warming associated with the low albedo of forests is offset by strong evaporative cooling from transpiration, but can become inhibited via drought (Bonan, 2008). Inversely, increased land evapotranspiration due to human-induced climate change has contributed to increases in agricultural and ecological droughts (Allan et al., 2023).

The combined effects tropical deforestation has on surface albedo, evapotranspiration, and cloud cover may accelerate global-scale warming in equatorial tropical forests (Bala et al., 2007). Observed water deficits across Central Africa are driven by factors such as lower rainfall, higher post-storm runoff (Beyene et al., 2012), and increased evapotranspiration due to warmer temperatures (Woodward, 2008; Abernethy et al., 2016). Notwithstanding, reduced evapotranspiration in a drier climate may lead to the loss of tropical forests (Baldocchi et al., 2000). Making biosphere evapotranspiration, surface roughness, and albedo all key controls in the general circulation of the atmosphere which could affect the regional climate (Hayden, 1998).

Regional climates in Central Africa have been identified as particularly sensitive to the exchange of water and energy between the surface, the atmosphere (Koster et al., 2004) and biogeophysical processes (Bonan, 2008). In particular, extreme events like cyclones, fire weather, heatwaves, heavy precipitation, and droughts are expected to increase in frequency and intensity in direct relation to increasing global warming (Allan et al., 2023). Whereas the impact of changes in precipitation over short and long timescales on tropical rainforests remains poorly understood (Malhi & Wright, 2004; Brando et al., 2010; Atkinson et al., 2011; Xu et al., 2011; Samanta et al., 2012; Asefi-Najafabady & Saatchi, 2013; Saatchi et al., 2013; Morton et al., 2014).

Carbon Dioxide

The central African evergreen tropical forests act as a carbon sink (Saatchi et al., 2011a), with the Congo Basin alone estimated to contain between 25 and 30 billion tons of carbon (Hoare, 2007). Over the last six decades, land and ocean have consistently taken-up about 56% per year of human induced CO₂ emissions but this capacity will decrease under higher CO₂ emissions (Allan et al., 2023). Plants can be influenced by CO₂ by altering water use efficiency, photosynthetic rates, light and nutrient efficiency, and the relative performance of C3 and C4 photosynthesis (Ratnam et al., 2011). Decreased CO₂ during the LGM drove the retreat of forests and expansion of flammable grasslands (Wooler et al., 2000). But, in 2019, global CO₂ concentrations were higher than at any time in at least 2 million years and CH₄ and N₂O were the highest in at least 800,000 years (Allan et al., 2023). Increasing CO₂ can result in a CO₂ fertilisation process whereby alongside warming, photosynthesis and plant productivity are enhanced (Norby et al., 2005; Woodward, 2008; Willis & Bhagwat, 2009; Willis & MacDonald, 2011).

Increasing CO₂ also often results in reduced transpiration/ leaf area that slows the depletion of soil water (Polley et al., 1997) and favours the recruitment of tree seedlings in place of savannah grasses (Bond & Midgley, 2012). Unlike tropical forests, the evaporative cooling generated by temperate and boreal forests have "little to none" and "counterproductive" mitigative effects on global warming respectively (Bala et al., 2007). Such that, when temperature increases in boreal forests terrestrial net primary production (vegetation growth) is enhanced, whereas it decreases for tropical forests due to greater evaporative demand that dries soil (Fung et al., 2005). Despite this, in recent decades tropical forests have shown an increase in productivity, tree growth, recruitment, mortality rates, and forest biomass (Lewis et al., 2009) probably due to increases in CO₂ (Lloyd & Farquhar, 2008). Subsequently, global warming may exacerbate under a positive feedback loop between decreased evaporative cooling and the amount of released CO₂ (Betts et al., 2004).

Larger forest trees are much more resistant to fire than their smaller counterparts (Bond & Midgley, 2012) and as larger trees establish themselves in place of grasses, there will be large impacts on fluxes of water, nutrients, carbon, and even weathering (Pagani et al., 2009). However, forest trees and savannah trees are considered as alternative ecosystem states between pyrophobic (fire repellant) and pyrophytic (fire resistant) respectively (Warman & Moles, 2009; Lehmann et al., 2011; Staver et al., 2011). Wildfires can play a natural part of the carbon cycle whereby CO₂ uptake during the growing season alleviates the emissions from fires (Valentini et al., 2014). Increasing CO₂ has indeed resulted in young savannah trees relative to 100 YBP, possessing faster growth, stronger regrowth, massive root systems, and larger starch reserves (Bond & Midgley, 2012). Since savannahs cover a large spatial extent, even a small increase in woody biomass would result in a substantially increased carbon sink (Archer et al., 2001; Asner et al., 2003; Williams et al., 2005).



Photo 5: Hairy Golden Orb-weaving Spider (*Trichonephila fenestrata*) in secondary rainforest

Climate model projections predict a drier Atlantic coast that may result in further decreases of rainforest extent (Delire et al., 2008). Large-scale conversion of tropical forest to pasture also creates a warmer, drier climate (Bonan, 2008) and between 1980 and 2000, nearly 60% of new agricultural land in Africa was derived from intact forests (Gibbs et al., 2010). Compared with pastureland, tropical forests maintain high rates of evapotranspiration, lesser surface air temperature, and greater precipitation (Bonan, 2008). Water stress affects the spatiotemporal probability of rapid land use changes, particularly regarding the suitability of land for agricultural purposes (Abernethy et al., 2016). Moreover, deforestation is coupled with the local and global climate, and as drier regions are more likely to be deforested (Malhi & Wright, 2004) global warming is accelerated (Cox et al., 2000). Therefore, in Central Africa climate change has the potential to accelerate forest loss through land conversion to agriculture (Laurance et al., 2014; Bele et al., 2015).

Forest Loss

A reduction in the absolute range of a species will likely increase the risk of local extinction (Thomas et al., 2004; Thuiller et al., 2005) and tropical rainforests are vanishing more rapidly than any other biome (Laurance, 1999; Achard et al., 2002). In Central Africa the main drivers of deforestation may be attributed to clearance for smallholder agriculture (Damanika & Wheeler, 2015), firewood and charcoal extraction (Mayaux et al., 2013), timber logging (often illegally) (Hoare, 2015), and mining activities (Edwards et al., 2014a). In addition to the forest habitat, native wildlife is also under threat from industrial logging, slash-and-burn farming, overhunting, and increased infrastructure development (Laurance et al., 2006).

A significant cause of deforestation in central Africa is the combination of rapid population growth and extensive road building (Wilkie & Laporte, 2001; Naughton-Treves & Weber, 2001) (Figure 1). The Congo Basin is critically impacted by the effects of road construction (Kleinschroth et al., 2019) with the World Bank predicting that over the next 15 years it will be the driving factor for deforestation in the basin (Megevan, 2013). The extraction of timber and precious minerals start near urban centres and major transport routes and pave the way for lower value extraction such as charcoal and smallholder agriculture (Ahrends et al., 2010). As seen within 10-15 km of villages and roads, whereby wildlife populations, such as duikers (forest antelope), bushpigs, elephants, and diurnal primates, experience significant declines (Barnes et al., 1991; Lahm et al., 1998; Fa et al., 2000; Wilkie et al., 2000; Lahm, 2001).

Over time, concentric haloes of deforestation form around cities and transport routes that drive the demand for yet more resources (Abernethy et al., 2016). Beyond a critical density of 8 people per square kilometre canopy loss accelerates and deforestation haloes around villages become permanent (Mayaux et al., 2013). Few species can survive in the mosaic of farmland and degraded scrub left behind from a large-scale forest invasion (Laurance, 2001a). These haloes may even merge in regions with high rural population densities such as parts of the DRC, Equatorial Guinea, and southern Cameroon (Abernethy et al., 2016). Road construction also serves as a gateway for oftentimes illegal mining, land colonisation, poaching, and exacerbated bushmeat hunting (Laurance et al., 2009; Barber et al., 2014; Kleinschroth & Healey, 2017).

Logging

It is estimated that only 1 person per square kilometre may be supported while sustaining tropical forest biomass (Robinson & Bennett, 2000). In terms of rural populations, all countries in the Central Africa region have surpassed this figure, except for Gabon (Abernethy et al., 2016). The deforestation rate in Central Africa is projected to increase by between 0.3% and 0.5% per year by 2020-2030 (Nogherotto et al., 2013). West Africa has only about 12% of its original rainforest left, Eastern Africa has just 8%, and Central Africa retains about 59% (Naughton-Treves & Weber, 2001). Logging may result in major deforestation (Kleinschroth et al., 2019) and frequently opens up forested areas to be encroached upon by shifting agriculture (Ernst et al., 2013; Molinario et al., 2017), which is the primary cause of forest loss across Africa (Curtis et al., 2018).

Without severe hunting pressure, wildlife may persist in logged forests albeit with reduced abundance (Johns, 1997). Nation states in the Congo Basin grant concessions (long-term leases) to logging companies (Karsenty & Ferron, 2017). For example the DRC has allocated extensive timber leases to Zimbabwe, Germany, Malaysia, and China (Laurance et al., 2006) with the most substantial lease spanning 1.5 times the size of Great Britain, 34 Ma hectares (Laurance, 2001b). Within well-managed concessions outside of DRC, the impact of logging roads on forest loss is mitigated by a fourfold higher rate of road abandonment than in areas outside concessions (Kleinschroth et al., 2019). While concessions do maintain much higher carbon stocks and wildlife habitat than cleared areas can (Goetz et al., 2015) most logging operations in the tropics are poorly managed as reduced-impact logging can be more expensive than traditional methods (Putz et al., 2000).

Unsustainably logged forests are prone to catastrophic wildfires (Cochrane et al., 1999; Nepstad et al., 1999), especially during droughts (Nepstad et al., 2004; Randerson et al., 2005; Poulter et al., 2010). Whereas selective logging can have positive impacts on tropical ecosystems (Laurance & Laurance, 1996; Johns, 1997; Malcolm & Ray, 2000; Fimbel et al., 2001;) and within concessions can provide economic value so long as they remain forested (Putz et al., 2012; Edwards et al., 2014b). However, results of a study across an array of African tropical forests, have found that over the course of 30 years selective logging significantly reduces carbon levels in forests (Valentini et al., 2014). Furthermore, the maze of tracks left by selective logging serves to enhance accessibility into the forested areas by hunters, miners, and farmers that subsequently destroy or severely degrade forests (Wilkie et al., 1992, 2001; Laurance, 2001a). Nevertheless, reduced impact logging should be promoted to reduce damage to the forest and maintain biodiversity (Sonwa et al., 2011).

Part 3 — People of the Forest

Forests are crucial to the well-being of a large number of poor people (Arnold & Ruiz-Pérez, 1998; Byron & Arnold 1999; Sunderlin et al., 2005; Tieguhong & Ndoye, 2006; Powell et al., 2011; Wan et al., 2011) yet they are rarely considered in national planning and policy (Bele et al., 2015) and neither are indigenous communities (Salick & Ross, 2018; Wiben et al., 2019). Across the Congo basin's 113 Ma population, many of those in rural settings still maintain a partly hunter-gatherer lifestyle, along with small-scale shift-and-burn agriculture for subsistence (Abernethy et al., 2016). The DRC contains anywhere between 600,000 to 2 million indigenous peoples, Burundi and Chad are home to hundreds of thousands, and Cameroon, Rwanda, and the CAR range in the tens of thousands at least (Wiben et al., 2019).

"Pygmies" are a collection of indigenous groups within the Congo Basin with a unique culture at risk of extinction (Kimbu, 2011). 'Pygmy' (Pygmée) is a stereotypical term (Moïse, 2011), but is used here to describe those who are directly dependent on the forest (Nkem et al., 2013). However, it should be noted that reductive categorisations such as "hunter-gatherer"/ "farmer" and "pygmy"/ "villager" transform the narrative into contrasting subsistence strategies and polarised relationships of power (Woodburn, 1997; Rupp, 2003). The Mbuti, Baka, and Batwa/ Twa peoples of Central Africa consider the term "Pygmies" derogatory and discriminatory (Wiben et al., 2019). Even reference to 'nomadic lifestyles' can be considered problematic when it comes to determining indigenous peoples' rights to property and land (Vinding, 2019). Therefore, long standing Pygmy relations with their 'exploiters' is important to consider when developing their platform for autonomy (Moïse, 2011).

Pygmies must use forests other than those bordering their settlements, which brings them into contact with other communities that influence their livelihood strategies such as with bushmeat sales (Nkem et al., 2013). They easily become victims of discrimination (Woodburn, 1997; Hewlett, 2000; Lewis, 2000; Jackson & Payne, 2003) and the powerful (Rupp, 2003, 2011) as they are lacking in social status (Lewis, 2000; Klieman, 2003) material wealth, development, and basic human and citizen rights (Hewlett, 2000). The Baka speaking people live in the tropical forests of southern Cameroon, northern Congo, and Gabon (Figure 2) but are exploited by Bantu farmers, villagers, and government officials (Hewlett, 2000). Notwithstanding, the Batwa peoples in the Tanganyika province of DRC face issues of violence and displacement and rebel groups in CAR are subjugating Pygmies to enact unsustainable hunting (Wiben et al., 2019).

Forest Resources

Forests provide refuges for biodiversity, food, medicinal, and forest products, regulation of the hydrological cycle, protection of soil resources, recreational uses, spiritual needs, and aesthetic values (Bonan, 2008). Millions of people are dependent on the forest for their livelihoods (Sunderlin et al., 2005) medicinal plants, and other resources (World Bank, 2004). Many rural communities in the Congo Basin also rely on shifting agriculture and forest resources for their livelihoods (Pelletier et al., 2018). Therefore forests play a crucial role in reducing the vulnerability of natural and social systems (Bele et al., 2015).

Many developing countries' national development plans and poverty reduction strategies must depend on tropical-forests as natural resource pools (Smith & Scherr, 2003; Nkem et al. 2010; 2013). Forests are used for in-situ forest hunting (Nasi et al., 2011) and gathering (Bahuchet, 1994; Wan et al., 2011), such as the forest communities of southern Cameroon whom depend on non-timber forest products to survive (Ndoye et al., 1998; van den Berg & Biesbrouck 2000; Sassen & Jum, 2007; Nkem et al. 2010). However, the provision of tropical-forest ecosystem goods and services (bushmeat, crops, timber, herbal medicine, honey, etc...) are rapidly declining and are predicted to be severely altered due to climate change impacts (Bele et al., 2015) such as global warming (Delire et al., 2008).

It is highly dispersed communities, predominantly indigenous ones, that are most vulnerable to the impacts of climate change (Nkem et al., 2010; Bele et al., 2011). For example, aforementioned CO₂ fertilisation effects can contribute to woody thickening in savannahs and cause land users to work much harder to maintain ecosystems such as grasslands (Bond & Midgley, 2012). Notwithstanding, some forms of small-scale agriculture result in an over-extraction of biomass that leads to land degradation and lower crop yields, which then further exacerbates the agricultural pressure on the land (Bennett et al., 2009). Mining, mechanised-agriculture, logging, and over-exploitation of forests for fuel-wood also threaten the livelihoods of forest-dwellers (Pearce, 2012). If forest communities' vulnerability to climate variability is addressed, the negative impacts of climate change may be reduced (Ribot et al., 1996).

Adaptation and Mitigation

Mitigation loosely defined reduces the accumulation of greenhouse gases (e.g. by enhancing carbon sinks), while adaptation reduces the vulnerability of societies and ecosystems to climate change (Ravindranath, 2007; Locatelli et al., 2008). Climate change impacts can either be mitigated by reducing greenhouse gases and enhancing carbon sinks (e.g. forests) (Klein et al., 2005; Allan et al., 2023) or adapted for by adjusting natural or human systems to ensure a sustainable continuation of peoples livelihoods (Ravindranath, 2007). While mitigation has global benefits and may well be implemented on the same local or regional scale, adaptation typically works on the scale of an impacted system, which is regional at best, but mostly local (Klein et al., 2005).

The United Nations Framework Convention on Climate Change has identified adaptation and mitigation (A/M) as potential policy responses to climate change (Klein et al., 2005; Ravindranath, 2007; Somorin et al., 2012). Adaptation is paramount as the poorest regions are anticipated to be hardest hit (Adger et al., 2003; Tol et al., 2004; Thomas & Twyman, 2005; Bele et al., 2011) but society cannot adapt indefinitely (King, 2004). Yet for many developing countries, adaptation is a priority due to a high vulnerability to climate impacts (Ayers and Huq, 2009; Somorin, 2010). Therefore, in 2015 the IPCCs Sixth Assessment Report used Shared Socio-economic Pathways (SSPs) to model different climate policy scenarios while integrating the challenges of A/M (Allan et al., 2023).

People's perspective on adaptation shapes the discourse around it and vice-versa, meaning adaptation policy is sensitive to the way in which it is perceived (Somorin et al., 2012). Climate change will affect temperature and rainfall patterns, which will expose forest-dependent groups to physical, mental, and health risks (Costello et al., 2009). Many forest communities have deep-grained social structures (van den Berg & Biesbrouck, 2000) and are relatively isolated, which makes adaptation difficult or impracticable without causing disruption (Nkem et al., 2013). Improperly implemented adaptation actions e.g. draining wetlands or moving peoples, could damage natural ecosystems and invoke maladaptation in the long-term (Campbell, 2009).

Developed countries contain the vast majority of the world's scientific and technological capacity, so it is in their interest to help developing countries leapfrog into non-carbon emissions technologies (King, 2004). State, market, and civil institutions play an important role in determining the response to conditions and risks associated with the environment and policy (Adger, 2000; Adger and Vincent, 2005; Smit and Wandel, 2006). Mechanisms such as the Global Environmental Facility and Kyoto Protocol's Adaptation Fund support adaptation strategies (Ravindranath, 2007). However, the Clean Development Mechanism, the Kyoto Protocol, and other global mechanisms mostly focus on mitigation (Ravindranath, 2007).



Photo 6: Netwing Beetle (Genus Lycus) in logged primary rainforest along the Mpivié river.

Mitigation Mechanisms

Economic studies indicate that reducing tropical deforestation is cost-effective for climate change mitigation (Enkvist et al., 2007; Strassburg et al., 2012). And, in 2005 Gabon and New Guinea spearheaded the REDD+ framework introduced by the UN Framework Convention on Climate Change, which was then internationally adopted at the Paris COP21 in 2015 (Abernethy et al., 2016). The framework aims to establish a global system for rewarding carbon storage and forest preservation (Abernethy et al., 2016) whereby an additional 1% of protected forests in Central Africa equates to over USD~\$500M (Bele et al., 2015). Such an approach has the potential to reduce damaging deforestation (O'Sullivan, 2009) and thereby reduce the vulnerability of forest-dependent communities to climate change (Brown et al., 2010).

Forest preservation and conservation are mitigation measures (Kremen et al., 2000; Thompson et al., 2009) that generate carbon credits (Somorin et al., 2012). The Congo Basin governments have shown a preference towards policy agenda such as REDD+ since it provides a source of revenue (Somorin et al., 2012). Investing in small-scale technologies that enhance soil moisture can reverse land degradation, curb erosion, boost carbon sequestration, and increase crop yields simultaneously (Enfors & Gordon 2007; Enfors et al. 2008). Payments for environmental services could then be used to supplement agricultural approaches and technologies that reduce emissions (Campbell, 2009).

Forests are able to sequester large amounts of carbon annually (Bonan, 2008) and REDD+ acts both to encourage carbon sequestration while simultaneously reducing our reliance upon natural forests (Somorin et al., 2012). However, while forests may be globally valued for climate regulation (Saatchi et al., 2011a), keeping forests can reduce the ability of rural households to escape poverty (Campbell, 2009). Forests may be locally valued for fuelwood but timber harvesting by one individual can lead to habitat decline, impacting another person's access to bushmeat (Fisher et al., 2009). Poorly implemented, the REDD+ framework could deprive people of the main source for sustaining their livelihoods (Locatelli et al., 2008) or endanger previously undisturbed forest types (Nkem et al., 2013).

Logging is expected to cause severe and potentially irreversible damage within the next decade (Bele et al. 2015) but the private forestry sector is generally against adaptation and consider it instead to be a responsibility of the government (Brown et al., 2010; Somorin et al., 2012). Moreover, the economic returns of industrial logging creates more immediate wealth than conservation initiatives (Kremen et al., 2000). The \$5 per ton rate for avoided carbon emissions set by REDD+ is significantly lower than competitive pricing against other land uses (Abernethy et al., 2016). REDD may work as a financial incentive for shifting cultivators and extensive cattle ranchers in the humid tropics, but not for commercial agricultural crops, logging, or even smallholder intensive agroforestry (Campbell, 2009).

Future land transformation in Africa is unknown, but a number of socioeconomic drivers may either increase or decrease pressure on the expansion of land-use activities (Thuiller et al., 2006). Mitigation efforts tend to operate within the context of international obligations, whereas adaptation is usually more local in nature, concerning economies and livelihoods (Martens et al., 2009). REDD+ can be said to target local communities (Somorin et al., 2012) as adaptation opportunities exist in forest conservation, afforestation, and reforestation (Ravindranath, 2007). But these solutions are ultimately framed within the context of a mitigation agenda (Campbell, 2009).

Adaptation Strategy

National adaptation policies and programmes focused on forest dependent-communities should consider livelihood strategies and natural resource management (Sonwa et al., 2011; 2012). In Pygmy societies human development could choose to focus on autonomy (e.g. in education, hunting, gathering, agriculture, and wage labour), entrepreneurialism (e.g. as a diviner/ healer, or master hunter), and creativity/ play (e.g. storytelling, comedic performances, graphic arts) (Moïse, 2011). The effects of climate change on the natural resource base must also be evaluated (Nkem et al., 2013). Therefore, better downscaled climate models are needed for seasonal forecasting and longer term planning under new climate regimes (Campbell, 2009).

The UNFCCC's 'New Collective Quantified Goal on Climate Finance' funding (UNFCCC, 2015) has been criticised for not meeting targets or the needs of those most affected by climate change and lacking understanding of climate finance (Pauw, 2025). No mitigation activity should reduce adaptation and so the global effort should attempt to focus on A/M synergies (Ravindranath, 2007). Moreover, international conventions have been known to deal with interrelated issues such as desertification, biodiversity loss, and climate change separately (Cowie et al., 2007). Our understanding may be improved by an interdisciplinary framework combining climate change science, climate impacts on ecosystems, and climate change mitigation policy (Bonan, 2008).

Institutions and networks allow for fair power distribution and so can enhance adaptive capacity to climate change (Tompkins and Adger, 2004; Walker et al., 2006; Pahl-Wostl, 2009). For instance, Cameroon's adaptive capacity would be enhanced by institutional linkages and coordination across the government, private sector and civil society (Brown et al., 2010). An integrated approach to A/M may combine adaptation and mitigation where synergies exist e.g. using REDD+ income to finance adaptation, but then treat A/M separately where trade-offs are inevitable (Somorin et al., 2012). To achieve a more integrated approach to A/M new forms of engagement between scientists, policy-makers, and wider stakeholder communities are required (Martens et al., 2009).

Governance & Tourism (Gabon & Cameroon)

Gabon is roughly the size of Italy at 268,000 km² and is situated on the equator just south of Cameroon (Laurance et al., 2006). The country contains dense rainforest, swamps, mangroves, steppes, savannahs (Lahm, 2001) and about 40% of the world's gorillas (Walsh, 2006). Ecotourism aims to improve protected area management and provide economic benefits to local residents but struggles to compete with the competitive pricing of timber companies for access to old-growth trees (Wilkie & Carpenter, 1999). Gabon designated thirteen national parks with the aim of creating a sustainable ecotourism model (Laurance et al., 2006) and unlike Cameroon, has an efficient tourism development, promotion, and marketing framework in place (Kimbu, 2011).



Photo 7: Variegated grasshopper (*Zonocerus variegatus*) within smallholder agriculture.

In the 1990s the French oil company Elf made substantial payments to political leaders in countries including Gabon and Cameroon (Henley, 2003). Crude oil exports contribute to over 40% of Gabon's gross domestic product (GDP) but are declining, which may result in increased timber exploitation (Laurance et al., 2006). Logging practices in Gabon frequently result in significant environmental damage due to inadequate regulation (Laurance et al., 2006), leading to extensive changes in forest structure, microclimatic shifts, soil compaction, and erosion (Fimbel et al., 2001; Johns, 1997). In Gabon alone less than 3% of the 221 registered logging companies have even submitted the required management plans by law (Laurance et al., 2006).

Cameroon covers 475,440 km² (Molua, 2007), situated on the west coast of Africa, it is the leading economy in the Central African region (Harilal et al., 2019) and is often described as an "Africa in miniature" (Mbenda et al., 2014). Cameroon has been ranked as one of the world's most corrupt according to the Transparency International Corruption Perception Index (Alemagi & Kozak, 2010). Despite this, the country receives the highest number of international visitors in the subregion compared to other Central African countries (Kimbu, 2011). Pygmies represent ~0.4% of Cameroon's 24 Ma population, with the most prominent being the 40,000 Baka living in the east and southern regions but the Bagyeli/ Bakola [Kola] and Bedzang [Medzan] are also present across southern Cameroon (Wiben et al., 2019). Cameroon has over 20 protected areas and many local communities depend on the natural resources within these protected areas for livelihood sustenance and cultural practices (Tchindjang et al., 2005).

Therefore, restricting local access to resources in protected areas may affect forest dependent communities' source of livelihood (Brown et al., 2010) and may undermine support for tourism initiatives (Harilal et al., 2019). In DRC the Kahuzi Biega National Park has been criticised for the expulsion and permanent exclusion of the Batwa indigenous peoples (Wiben et al., 2019). Cameroon validated the REDD+ National Strategy in 2018 following the creation of the 'Platform REDD+ and Indigenous Peoples of Cameroon' (Wiben et al., 2019). Then, over the next 2000 years global mean sea level will rise by 2 to 3 metres if warming is limited to just 1.5 °C, but could reach 22 metres under 5 °C of warming (Allan et al., 2023). Gabon and Cameroon had already surpassed over 50% of their population living in urban areas by 1985 (Abernethy et al., 2016) and even under RCP 2.6, rising sea levels may result in the loss of coastal forests and wetlands, costing billions of dollars by the end of the century (Brown et al., 2011).

Viability of Ecotourism

Central African nations have commonly experienced civil wars, political coups, rebellions, and harassment from police, immigration, and customs officers (Wilkie & Carpenter, 1999). Political instability, recurring wars, and widespread corruption pose challenges for forest conservation (Laurance et al., 2006). For example, the mass displacements and violence in CAR (Wiben et al., 2019) and the DRC (Samset, 2002; Chilunjika, 2024). Additionally, Central African government representatives have been known to be unprofessional and uncourteous (Hewlett, 2000). And, in the period leading up to 2011, the Central Africa subregion trailed behind Eastern and Southern Africa in terms of growth in tourism (Kimbu, 2011).

Gabon and to some extent Cameroon stand out as not having experienced violent internal conflicts over the past fifty years (Abernethy et al., 2016). Since 2004, the Cameroonian government has aimed to reduce its reliance on oil and other extractive industries by developing other sectors such as tourism (Kimbu, 2011). Which has become increasingly difficult since northern Cameroon has been subjected to terrorist attacks from groups such as Boko Haram (Tichaawa, 2017). Fifteen percent of Cameroon's land area is dedicated to nature conservation areas (Riley & Riley, 2005) and maintain relative peace and political stability (Shackley, 2007), which lays a solid foundation for developing a dynamic ecotourism industry (Shackley, 2007; Harilal et al., 2019).

Ecotourism can become a major source of revenue for protected areas if certain criteria are met as historically seen with Gorilla tourism in the DRC, Rwanda, and Uganda (Wilkie & Carpenter, 1999). However, the revenue from ecotourism may be insufficient to cover management costs as seen with the Dzanga-Sangha Special Reserve in the CAR, and the Lope Reserve in Gabon (Wilkie & Carpenter, 1999). For ecotourism to be successful in Cameroon a sustainable tourism industry must be present (Harilal et al., 2019) which requires advancements in infrastructure and collaboration amongst varied stakeholders (Murphy et al., 2000; Khadaroo & Seetanah, 2007; Kimbu, 2011).




Figure 2: Map of Cameroon depicting Sentinel-2 Land Cover and forest dwelling linguistic groups.

Part 4 — Ecosystem Conservation

Ecosystem Services (ES) may be loosely defined as the benefits derived from ecosystems (Bennett et al., 2009). That said, humans have engineered many ecosystems for the efficient production of services such as food, timber, and fibre (Foley et al. 2005; Kareiva et al. 2007; Monfreda et al. 2008; Ramankutty et al. 2008). Tropical and subtropical African biomes such as forest and savannah are examples of important ES that carry associated human benefits (Willis et al., 2013). African forests provide regulating services, such as carbon sequestration (Saatchi et al., 2011a) and provisioning services, such as timber, non-timber forest products (Hoare, 2007), and biofuels (Walter & Segerstedt, 2012). In East Africa it is noted that climate change threatens to significantly impact the provision of ES into the future (Conway et al., 2005). ES is therefore an anthropocentric term tied to human existence and so concerns stakeholders across all of society (Fisher et al., 2009).

The Millennium Ecosystem Assessment (MA) is a global study concerning ES that tries to classify services into supporting, regulating, provisioning, and cultural categories (Alcamo et al., 2005). Fisher et al., (2009) states that the MA's definition of ES therefore includes cultural meaning, recreation, and spiritual fulfilment. This can result in errors such as double-counting in overlapping categories like in both the drinking water and spiritual fulfilment derived from a sacred forest (Fisher et al., 2009). Therefore, ES may alternatively be categorised by Organisation (ecosystems), Operation (process/ functioning of ecosystems), and Outcome (benefit provided to humans) (Fisher et al., 2009): For example, the pattern of water stress across Central African forests will play a crucial role in shaping the future of land use in the region (Bell et al., 2015). Forest cover is tied to soil moisture, which is influenced by rainfall and evapotranspiration (Zelazowski et al., 2011). In this case the Organisation is forests, an Operation is rainfall and evapotranspiration, and an Outcome is soil moisture levels.

By simply analysing trade-offs and synergies between ESs at a single point in time or only examining spatial concordance, it risks misunderstanding the underlying mechanisms at play (Bennett et al., 2009). ES can exhibit spatio-temporal dynamism, whereby an example might be water regulation services provided by a mountain forest which will benefit downstream areas over time (Fisher et al., 2009). Moisture availability in the root zone of plants is determined by the sum of rainfall, the level of runoff and evapotranspiration (Reager & Famiglietti, 2013). Notwithstanding, the trees of some forests have deep roots that grant access to groundwater that may delay the effects of water stress (Zhou et al., 2014) but inhibit their expansion (Drake et al., 2011). Therefore, maintaining relationships between ES are crucial for bolstering their resilience and reducing their decline (Bennett et al., 2009) yet the MA found that 15 out of 24 studied ES were declining globally (Fisher et al., 2009).

Focusing only on a limited set of ES such as soil moisture alone, risks regime shifts that could result in the unexpected and sudden loss of other ES (Gordon et al., 2008). As an example, pollination is considered an indirect ES and is crucial for the reproduction of many plants, including crops like almonds (Fisher et al., 2009) and coffee (Ricketts et al., 2004). Integrating forest patches in agricultural landscapes can improve pollination and pest control, which in turn improves crop yield (e.g. Olschewski et al., 2006; Bodin et al. 2006; Priess et al., 2007; Kremen et al. 2007; Ricketts et al. 2008) (Bennett et al., 2009). Small forest patches of moderately-managed landscapes have been demonstrated to conserve tree diversity and forest structure (Hernández-Ruedas et al., 2014). Planting trees also enhances carbon sequestration, but the presence of trees can increase evapotranspiration which may lead to exacerbated water stress (Fahey & Jackson 1997; Engel et al. 2005). Meaning that already dry agricultural plots could actually suffer from reduced moisture levels by planting forest patches nearby.

A major limitation of ES comes in the form of our own limited understanding of their workings (Kremen & Ostfeld, 2005; Carpenter et al. 2009) and of biodiversity itself (Midgley & Thuiller, 2005). Ecosystems exhibit complex lagged relationships between rainfall, vegetation phenology and photosynthetic activity (Lewis, 2006; Enquist & Enquist, 2011; Lewis et al., 2013) and large spatiotemporal variabilities in rainfall (Malhi & Wright, 2004). In Africa forest against savannah distribution doesn't change over a rainfall gradient, which suggests that the two vegetation types are largely determined by local processes (Staver et al., 2011). ES are characterised by feedback loops, time lags, and nested phenomena (Limburg et al., 2002) and differing ecosystems have differing responses from various plant species to drought (Lucht et al., 2006).

Classifying Spatial Costs

Diminishing environmental quality has a disproportionate impact on those marginalised by the market economy (Dasgupta, 2001; Adger et al., 2003; Nkem et al., 2010). However, the complexity is amplified when differing interests in ES across various spatial scales are considered (Hein et al., 2006). For example, a crucial aspect of ES is in understanding how they contribute to human welfare and equity (e.g. Aylward & Barbier, 1992) (Fisher et al., 2009). Therefore, a major challenge exists in reconciling values derived from public opinions on biodiversity with market-based values for commodities like timber (e.g. Aldred, 2002) (Fisher et al., 2009). Another example might be with integrating the valuation of tangible services like water provision with intangible aspects like spiritual benefits from a sacred forest (Fisher et al., 2009). Conservation costs can vary dramatically across different spatial extents (Ando et al., 1998; Ferraro, 2003; Balmford et al., 2003) and intangible ones like biodiversity's existence value are challenging to integrate (Naidoo et al., 2006).

Species' risk of extinction increases under a combination of threats such as environmental warming, overexploitation, and habitat fragmentation (Mora et al., 2007). Therefore, in conservation planning, economic costs are integral for conducting cost-benefit analysis (Margules & Pressey, 2000) and plans that take only a random sampling approach are unlikely to avert extinctions (Woodroffe & Ginsberg, 1998). Estimation methods for conservation costs traditionally have made use of aggregate measures such as total area, units, or constraints (e.g. Dobson et al., 1997; Araújo et al., 2002) that assume an equal cost (Naidoo et al., 2006). Initially conservation goals may be achieved at a relatively low cost and effort, but they exhibit diminishing returns as the cost and effort required to achieve complete conservation objectives escalates (Montgomery & Adams, 1994; Ando et al., 1998; Polasky et al., 2001; Naidoo & Adamowicz, 2005).

Migratory Response to Climate Change

Climate, particularly rainfall conditions, has an influence on mammal distribution and abundances in Africa (Owen-Smith, 1990; Mills et al., 1995). Tropical forests are more resilient to climate impacts than monoculture plantations or any artificial forest, especially when they are rich in biodiversity (Ravindranath, 2007). Species-climate envelope models (CEM) predict species distribution under differing climate scenarios but are highly sensitive to their chosen algorithm and context (Araújo et al., 2005; Hijmans & Graham, 2006). Also, the historical ranges of species hundreds or thousands of years ago is lacking in available data that may limit model predictions under future scenarios (Carvalho et al., 2021). Despite this, in the forthcoming decades CEMs predict the extinction of many African mammals (Thuiller et al., 2006) and sub-Saharan plant species (McClean et al., 2005).

A key research challenge is in quantifying the impact of climate change on African tropical and subtropical plant and animal species that are important for current and future ES provision (Willis et al., 2013). Land transformation by humans restricts species range migrations under climate change and only then does the risk of extinction become a real possibility (Thuiller et al., 2006). Climate scenarios with low greenhouse gas emissions strongly limit the negative impacts of a changing climate (Allan et al., 2023). Also, during the AHP forest expansion was extensive but nonlinear as different species underwent range shifts at drastically different rates and directions (Watrin et al., 2009; Drake et al., 2011; Lézine et al., 2011). Therefore, understanding the nonlinear behaviour of African biota in response to climate change is essential in understanding the reliability of current and future predictions (Willis et al., 2013).

A major conservation challenge is to assess what effects climate and land use changes will have on species distributions under a range of environmental scenarios (Carvalho et al., 2021). Many terrestrial organisms are shifting their range to a higher latitude or elevation in response to climate change (Chen et al., 2011). Climate-driven crop migration and yield reductions have already been observed (Zhang et al., 2017; Sloat et al., 2020) and are projected for the future (King et al., 2018; Ceglar et al., 2019). As range shifts occur, new ecological relationships will be forged that may unpredictably transform the character of species interactions and fundamental ecosystem processes (Post et al., 1999; Walther et al., 2002; Schmitz et al., 2003) that could lead to species loss (Root et al., 2003).

Protected Areas and Forest Conservation

National parks and bio-reserves are key conservation tools that can be used to protect species and habitats within the confines of their boundaries (Thuiller et al., 2006). International agreements like REDD or the Kyoto Protocol are deemed necessary for managing congestible public goods (e.g. clean air) at a global scale, but differ from local-scale management strategies (e.g. Sandler, 1998) (Fisher et al., 2009). At a global scale cost per unit area for protection decreases with increasing area size, indicating that larger areas provide more benefits for a given investment (Naidoo et al., 2006). When a 'protected area' is formed and effectively managed, it can protect biodiversity and prevent forest degradation or conversion (Ravindranath, 2007). However, in designating a new national park as a protected area, decision makers are often faced with the challenge of balancing the equitable distribution of benefits with other objectives, such as education (Fisher et al., 2009).

Climate change will alter the distribution of many habitats and component species, subsequently threatening the effectiveness of protected areas to meet conservation needs (Burns et al., 2003). Climate mitigation policies must recognise the influences of forests on climate, their differing spatial and temporal scales, and their long-term effectiveness and sustainability in a changing climate (Bonan, 2008). However, focusing ecological management and monitoring on provisioning or cultural services alone, threatens to overlook the various interactions between ES (Carpenter et al. 2006). Linking protected areas together is an adaptive strategy for flora and fauna to migrate in response to a changing climate (Ravindranath, 2007). Therefore, other objectives may include landscape management for preserving biodiversity and economic valuations such as ecosystem services and conservation value (Fisher et al., 2009).

Legally protected areas in Central Africa play a crucial role in forest preservation by largely withstanding the impacts of extractive industries, land conversion, and local hunting (Abernethy et al., 2016). However, evidence suggests that protected areas in equatorial Africa aren't able to withstand current poaching levels (Walsh et al., 2003; Bennett, 2011). Without adequate funding, governments may permit or overlook illegal and lucrative activities, such as logging, bushmeat hunting, and gold or diamond mining within protected areas (Wilkie & Carpenter, 1999). Logging concessions are an example of what might form a buffer zone in the vicinity of protected areas (Clark et al., 2009b). Without effective education, hunting regulations, buffer zones, and a reduction in logging leases, national parks risk becoming isolated pockets (Woodroffe & Ginsberg, 1998) of protected land.



Photo 8: Round-backed millipede on Fernan Vaz lagoon's banks.

Evaluating Costs

In developing countries there is often a lack of available data on land prices available at scales necessary for detailed conservation planning (Naidoo et al., 2006). Designating areas for conservation may elevate the value and, consequently, the price of adjacent lands, potentially increasing the likelihood of habitat conversion in nearby areas (Wu, 2000; Armsworth et al., 2006). For instance, protected areas may be considered as simply displacing logging to other areas (Barber et al., 2014). Which would mean increased pressure on non-protected areas and may even cause a net decrease in biodiversity (Wu, 2000; Naidoo et al., 2006). The challenge of lacking spatial data in valuations may be partly overcome by modelling cost-maps using inputs such as soil type, climate, current land cover, and proximity to roads (e.g. Chomitz et al., 2005) (Naidoo et al., 2006).

Conservation initiatives will likely only work where the value of ES exceeds the opportunity costs forgone in abandoning the first best land-use option, plus the transaction costs (Wunder, 2008). The spatial variability of these costs may be partly accounted for by considering factors like the vulnerability or threats to habitats, or proximity to infrastructure and population density (Naidoo et al., 2006). The needs of many species may be approximated using just a few habitats (Krosby et al., 2015). Yet, protection, restoration, and conservation decisions are likely to consider habitats individually, despite some species being dependent on multiple priority habitats (Travers et al., 2021).

Habitat Fragmentation

Fragmented and isolated patches of forest are vulnerable to invasive species and hold little value for mammalian biodiversity (Gibson et al., 2013) and a reduction in the absolute range of a species will likely increase the risk of local extinction (Thomas et al., 2004; Thuiller et al., 2005). Possible landscape configurations may raise the problem of optimising where to restore over what to conserve (Hodgson et al., 2016). For example, why not high-quality 'refugia' habitats that have remained stable over long periods of time and are expected to be resilient towards climate change (Graham et al., 2010; Shoo et al., 2011). There are still many intact ecosystems about the globe that offer excellent conservation opportunities (Riggio et al., 2020) but the conservation of forests may benefit from directing efforts at large expanses of habitat (Morcatty et al., 2013).

About half of terrestrial protected areas are lacking in connectedness across the globe (Saura et al., 2018) with only 9.7% of Earth's terrestrial protected network actually considered structurally connected (Ward et al., 2020). Habitat fragmentation has been found to reduce biodiversity to anywhere between 13% and 75% while decreasing biomass and altering nutrient cycles (Haddad et al., 2015). Habitat loss and climate change pose a synergistic threat to biodiversity (Oliver & Morecroft, 2014; Synes et al., 2020; Mancini et al., 2022) and without proper designation, the likelihood of degradation or destruction increases and may severely impact the connectivity of habitat patches (Travers et al., 2021). Despite wildlife habitats across the world consisting mainly of small patches of poorly connected (semi-) natural habitat (Mancini et al., 2022), patches critical to species range expansion have generally not been prioritized and are underprotected (Travers et al., 2021).

Range Expansion (Primates)

Habitat loss and fragmentation, land use change and hunting, and the effects of commodity growth and trade is causing many primate species to approach extinction (Estrada et al., 2018). Climate change too will expose species, especially forest-dwelling primates to unsuitable conditions and limit their dispersal range (Carvalho et al., 2019). There are lots of uncertainties in modelling species responses to global environmental changes (Araújo & New, 2007; Thuiller et al., 2019). But over the past twenty years African apes have faced dramatic declines in suitable environment (Junker et al., 2012) in addition to large population losses (Walsh et al., 2003; Plumptre et al., 2016; Kühl et al., 2017; Strindberg et al., 2018) which is predicted to continue into the future due to climate change (Carvalho et al., 2021). Notwithstanding, different species have different dispersal abilities (Mancini et al., 2022) and range expansion may not necessarily be achieved through successful colonisation only (MacArthur & Wilson, 2001; Travers et al., 2021).

The limited dispersal capacity of many mammals means that range gains and losses can operate at timescales in the hundreds to thousands of years (Schloss et al., 2012). An ape generation lasts under 30 years (Plumptre et al., 2016; Kühl et al., 2017) which is unlikely to see any significant migration (Carvalho et al., 2021). Range expansion may be facilitated through improving the quality of existing habitat patches, increasing the permeability of the surrounding landscape, restoring degraded habitat, creating new 'stepping-stone' habitat patches, and by enlarging existing patches (MacArthur & Wilson, 2001; Crick et al., 2020; Synes et al., 2020). It is recommended that for great apes effective conservation strategies should be taxon specific and focus on existing protected areas, and proposing new ones based on habitat suitability models (Carvalho et al., 2021).

Technology

The drivers of rapid environmental change in Central Africa are primarily external economic forces that invest in resources like (Lewis et al., 2015) oil, minerals (Edwards et al., 2014a), timber (Hoare, 2015), agricultural land (Feintrenie, 2014), and wildlife products (UNEP & IUCN, 2013). Whereas, internally the surge in prices for ivory and other wildlife products since the mid-2000s has drawn well-funded criminal organisations to Africa, equipped with high-tech navigational tools, powerful rifles, and mobile communications (Abernethy et al., 2016). Synchronously, conservation technology has become increasingly important in the 21st century, with non-invasive techniques being at the forefront of these efforts (Piel et al., 2021). Therefore, conservation management must integrate a dynamic framework to strategically sequence conservation actions tailored to varying threat levels faced by different conservation targets (Naidoo et al., 2006).

Technology enhances the field of conservation by providing innovative solutions for monitoring, early-warning damage mitigation, and policy implementation (Berger-Tal & Lahoz-Monfort, 2018). For instance, mapping forest carbon stocks may assist countries in assessing the likely carbon emissions avoided by implementing policies aimed at reducing deforestation and forest degradation (Saatchi et al., 2011a). Furthermore, modelled predictions of land suitability for crops is important to determine how climate-driven tree loss might influence governmental decisions and land use allocation (Abernethy et al., 2016). Therefore, better data collection is key to informing management decisions (Berger-Tal & Lahoz-Monfort, 2018), especially in a world where catastrophe mitigation can be financed by such mechanisms as crop-weather insurance (Campbell, 2009), grants, loans, and even bonds (Motlagh et al., 2024).

Obtaining high temporal resolution data series required for predictive climate models, may be facilitated by high-resolution paleoecological data and independent climate records (Willis et al., 2013). However, the sparse observational network of the Congo basin severely inhibits climate change projections (Delire et al., 2008; Washington et al., 2013) and tropical Africa is lacking in high-resolution paleoecological data such as pollen records (Ngomanda et al., 2007). Therefore it may be necessary to implement a network of radiosonde stations, weather radar (Washington et al., 2013), ground observations and drought manipulative experiments across the region (Zhou, 2014). Though even with strong predictive capabilities, unpredictable events such as a series of large volcanic eruptions, may cause several decades of substantial global and regional climate perturbations (Allan et al., 2023).

Wildlife

Automating the annotation of camera trap images/ videos (Rowcliffe et al., 2016; Chen et al., 2019) and sound recordings (Kalan et al., 2015; Spillmann et al., 2015, 2017) using machine learning, could allow researchers greater efficiency and accuracy when monitoring wildlife (Piel et al., 2021; Tuia et al., 2022). Machine learning algorithms can be incredibly powerful for analysing small datasets, but they can lose resolution when applied to larger and more complex datasets (Joppa, 2015). Camera traps and especially passive acoustic monitoring (PAM), may be set up to collect data remotely and are deployable in remote areas, making them ideal for studying primates (Enari et al., 2019; Cruncheon et al., 2020) like gorillas. PAM may be used in dense tropical forests or at night, and can be deployed for months or years at multiple locations (Piel et al., 2021). But, while PAM offers more effective detection, high intracaller acoustic variability makes it difficult to implement automated detection (Zimmermann & Lerch, 1993; Bouchet et al., 2012).

Forests

Drones may be utilised to facilitate detailed land-cover classification and monitor changes in certain variables, but fall short of detecting individuals at distance or those obscured by overhanging canopy (Piel et al., 2021). More generally, remote sensors (RS) are mounted to drones, aircraft, satellites, environmental features, or animals and can capture detailed real-world data (Tuia et al., 2022). For example, project and national scale mapping of carbon stocks traditionally use satellite imagery and ground-based inventory samples of forest carbon density (Saatchi et al., 2007; Asner et al., 2010; Neigh et al., 2013; Hu et al., 2016). Light detection and ranging (LiDAR) and radar have also been used in RS to provide higher-resolution forest carbon density estimates (Drake et al., 2002; Asner et al., 2010; Saatchi et al., 2011b; Lang et al., 2021). RS are therefore essential in the systematic monitoring of forests to understand their response to climate change (Malhi & Wright, 2004; Asefi-Najafabady & Saatchi, 2013).

Local Context

Certain technologies may be viewed as “neoliberal conservation” whereby local communities are excluded from decision making (Arrighi et al., 2003; Igoe & Brockington 2007). However, there is a limitation of computational technology in conservation and a lack of funding and expertise in certain areas, such as with drones (Joppa, 2015). In the case of great apes in the wild, researchers have traditionally relied on non-invasive sampling, such as collecting faeces in order to carry out genetic analysis (Städele et al., 2022). Other conventional wildlife monitoring systems involve manual counting and observations of animals, which are labour-intensive, time-consuming, expensive (Witmer, 2005), and prone to errors or bias (Burghardt et al., 2012; McEvoy et al., 2016). For local communities to become responsible for the nature on their own lands, easy access to necessary skills and technology is required (Goldman, 2001) in addition to an appropriate strategy, such as crowdsourcing (Pimm et al., 2015). Thankfully, this has become possible with the advent of cheap prototyping and designing tools such as mobile phones, 3D printers, affordable PCBs, microcontrollers, and single-board computers (Berger-Tal & Lahoz-Monfort, 2018).



Photo 9: Dwarf crocodile (*Osteolaemus tetraspis*) along the Mpivié river.

Summary

Central Africa is home to the Guineo-Congolian rainforests that have persisted in one form or another for millions of years. Gorillas have evolved alongside the forests but habitat loss and the bushmeat trade threaten their continued existence. Increasing temperatures from climate change is drying soil and excessive atmospheric CO₂ is altering vegetation compositions. Suitable agricultural land is adversely affected and so more forest is cleared in a feedback loop as fertile soils become increasingly rare. Logging practices could be made more sustainable but are generally destructive and pave the way for large-scale human habitation at the expense of biodiversity. In response to local, regional, and global pressures forest dependent and indigenous communities' are being driven to unsustainably exploit the forest too. Since their livelihoods and access to land are being drastically altered, developing a strategy to support these people must go beyond reducing pollution to meet climate goals. If correctly perceived and implemented then the conservation of tropical rainforest can contribute to both adaptation and mitigation actions.

Gabon and Cameroon have designated a number of national parks but wildlife and forest degradation is ongoing and climate change is making matters worse. Ecotourism offers a source of revenue for keeping ecosystems intact, but cannot compete with the economic returns of extractive industries like logging. Ecosystem Services are a school of thought that try to value the conservation of ecosystems but contrasting facts and opinions make defining ES difficult. Nevertheless, conservationists must consider a cost-benefit analysis when deciding where or what to attempt to conserve which means also understanding the effects of climate change and local contexts. The biota of protected areas may need to migrate in response to climate change and so linking protected areas together or conserving large expanses of habitat could help to avoid extinctions. Models are needed that can accurately predict land use change and species migration but high quality data is lacking for Central Africa. The emergence of cheap technologies that can be deployed to collect lots of data in the field is making conservation aims more achievable.

References

- Abernethy, K., Maisels, F., & White, L. J. (2016). Environmental issues in central Africa. *Annual Review of Environment and Resources*, 41, 1-33.
- Achard, F., Eva, H. D., Stibig, H. J., Mayaux, P., Gallego, J., Richards, T., & Malingreau, J. P. (2002). Determination of deforestation rates of the world's humid tropical forests. *Science*, 297(5583), 999-1002.
- Adger, W. N., Huq, S., Brown, K., Conway, D., & Hulme, M. (2003). Adaptation to climate change in the developing world. *Progress in development studies*, 3(3), 179-195.
- Adger, W. N. (2000). Institutional adaptation to environmental risk under the transition in Vietnam. *Annals of the Association of American Geographers*, 90(4), 738-758.
- Adger, W. N., & Vincent, K. (2005). Uncertainty in adaptive capacity. *Comptes Rendus Geoscience*, 337(4), 399-410.
- Ahrends, A., Burgess, N. D., Milledge, S. A., Bulling, M. T., Fisher, B., Smart, J. C., ... & Lewis, S. L. (2010). Predictable waves of sequential forest degradation and biodiversity loss spreading from an African city. *Proceedings of the National Academy of Sciences*, 107(33), 14556-14561.
- Alcamo, J., Ash, N. J., Butler, C. D., Callicott, J. B., Capistrano, D., Carpenter, S. R., Castilla, J. C., Chambers, R., Chopra, K., Cropper, A., Daily, G. C., Dasgupta, P., de Groot, R., Dietz, T., Duraiappah, A. K., Gadgil, M., Hamilton, K., Hassan, R., Lambine, E. F., ... Zurek, M. B. (2005). Millennium Ecosystem Assessment: Ecosystems and Human Well-being. Island Press.
- Aldred, J. (2002). Cost-benefit analysis, incommensurability and rough equality. *Environmental values*, 11(1), 27-46.
- Alemagi, D., & Kozak, R. A. (2010). Illegal logging in Cameroon: Causes and the path forward. *Forest Policy and Economics*, 12(8), 554-561.
- Allan, R. P., Arias, P. A., Berger, S., Canadell, J. G., Cassou, C., Chen, D., ... & Zickfeld, K. (2023). Intergovernmental Panel on Climate Change (IPCC). Summary for Policymakers. In Climate change 2021: The physical science basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change (pp. 3-32). Cambridge University Press.
- Ando, A., Camm, J., Polasky, S., & Solow, A. (1998). Species distributions, land values, and efficient conservation. *Science*, 279(5359), 2126-2128.
- Araújo, M. B., Williams, P. H., & Fuller, R. J. (2002). Dynamics of extinction and the selection of nature reserves. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 269(1504), 1971-1980.
- Araújo, M. B., Pearson, R. G., Thuiller, W., & Erhard, M. (2005). Validation of species-climate impact models under climate change. *Global change biology*, 11(9), 1504-1513.
- Araújo, M. B., & New, M. (2007). Ensemble forecasting of species distributions. *Trends in ecology & evolution*, 22(1), 42-47.
- Archer, S., Boutton, T. W., & Hibbard, K. A. (2001). Trees in grasslands: biogeochemical consequences of woody plant expansion. In Schulze, E.-D., Heimann, M., Harrison, S., Holland, E., Lloyd, J., Prentice I. C., & Schimel, D. (Eds.), *Global biogeochemical cycles in the climate system*. Academic Press, 115-137.
- Armstrong, P. R., Daily, G. C., Kareiva, P., & Sanchirico, J. N. (2006). Land market feedbacks can undermine biodiversity conservation. *Proceedings of the National Academy of Sciences*, 103(14), 5403-5408.
- Arnold, J. E. M., & Ruiz-Pérez, M. (1998). The role of non-timber forest products in conservation and development. In Wollenberg E., & Ingles, A. (Eds.), *Incomes from the forest: methods for the development and conservation of forest products for local communities*. Centre for International Forestry Research, 17-42.
- Arrighi, G., Silver, B. J., & Brewer, B. D. (2003). Industrial convergence, globalization, and the persistence of the North-South divide. *Studies in comparative international development*, 38, 3-31.
- Asefi-Najafabadi, S., & Saatchi, S. (2013). Response of African humid tropical forests to recent rainfall anomalies. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1625), 20120306.
- Asner, G. P., Archer, S., Hughes, R. F., Ansley, R. J., & Wessman, C. A. (2003). Net changes in regional woody vegetation cover and carbon storage in Texas drylands, 1937-1999. *Global Change Biology*, 9(3), 316-335.
- Asner, G. P., Powell, G. V., Mascaró, J., Knapp, D. E., Clark, J. K., Jacobson, J., ... & Hughes, R. F. (2010). High-resolution forest carbon stocks and emissions in the Amazon. *Proceedings of the National Academy of Sciences*, 107(38), 16738-16742.
- Atkinson, P. M., Dash, J., & Jeganathan, C. (2011). Amazon vegetation greenness as measured by satellite sensors over the last decade. *Geophysical Research Letters*, 38(19).
- Augusto, L., Dupouey, J. L., & Ranger, J. (2003). Effects of tree species on understory vegetation and environmental conditions in temperate forests. *Annals of Forest Science*, 60(8), 823-831.
- Ayers, J. M., & Huq, S. (2009). The value of linking mitigation and adaptation: a case study of Bangladesh. *Environmental Management*, 43, 753-764.
- Aylward, B., & Barbier, E. B. (1992). Valuing environmental functions in developing countries. *Biodiversity & Conservation*, 1, 34-50.
- Bakker, E. M. V. Z., & Mercer, J. H. (1986). Major late Cainozoic climatic events and palaeoenvironmental changes in Africa viewed in a world wide context. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 56(3-4), 217-235.
- Bala, G., Caldeira, K., Wickett, M., Phillips, T. J., Lobell, D. B., Delire, C., & Mirin, A. (2007). Combined climate and carbon-cycle effects of large-scale deforestation. *Proceedings of the National Academy of Sciences*, 104(16), 6550-6555.
- Balmford, A., Gaston, K. J., Blyth, S., James, A., & Kapos, V. (2003). Global variation in terrestrial conservation costs, conservation benefits, and unmet conservation needs. *Proceedings of the National Academy of Sciences*, 100(3), 1046-1050.
- Baldocchi, D., Kelliher, F. M., Black, T. A., & Jarvis, P. (2000). Climate and vegetation controls on boreal zone energy exchange. *Global change biology*, 6(S1), 69-83.
- Bele, M. Y., Sonwa, D. J., & Tiani, A. M. (2015). Adapting the Congo Basin forests management to climate change: Linkages among biodiversity, forest loss, and human well-being. *Forest Policy and Economics*, 50, 1-10.
- Bele, M. Y., Somorin, O., Sonwa, D. J., Nkem, J. N., & Locatelli, B. (2011). Forests and climate change adaptation policies in Cameroon. *Mitigation and Adaptation Strategies for Global Change*, 16(3), 369-385.
- Barber, C. P., Cochrane, M. A., Souza Jr, C. M., & Laurance, W. F. (2014). Roads, deforestation, and the mitigating effect of protected areas in the Amazon. *Biological conservation*, 177, 203-209.
- Barnes, R. F., Barnes, K. L., Alers, M. P. T., & Blom, A. (1991). Man determines the distribution of elephants in the rain forests of northeastern Gabon. *African Journal of Ecology*, 29(1), 54-63.
- Bayon, G., Dennielou, B., Etoubleau, J., Ponzevera, E., Toucanne, S., & Bermell, S. (2012). Intensifying weathering and land use in Iron Age Central Africa. *Science*, 335(6073), 1219-1222.
- Becquet, C., & Przeworski, M. (2007). A new approach to estimate parameters of speciation models with application to apes. *Genome research*, 17(10), 1505-1519.
- Bell, J. P., Tompkins, A. M., Bouka-Biona, C., & Sanda, I. S. (2015). A process-based investigation into the impact of the Congo basin deforestation on surface climate. *Journal of Geophysical Research: Atmospheres*, 120(12), 5721-5739.
- Bennett, E. M., Peterson, G. D., & Gordon, L. J. (2009). Understanding relationships among multiple ecosystem services. *Ecology letters*, 12(12), 1394-1404.
- Bennett, E. L. (2011). Another inconvenient truth: the failure of enforcement systems to save charismatic species. *Oryx*, 45(4), 476-479.
- Berger-Tal, O., & Lahoz-Monfort, J. J. (2018). Conservation technology: The next generation. *Conservation Letters*, 11 (6), e12458.
- Bermejo, M., Rodríguez-Tejedor, J. D., Illera, G., Barroso, A., Vilà, C., & Walsh, P. D. (2006). Ebola outbreak killed 5000 gorillas. *Science*, 314(5805), 1564-1564.
- Betts, R. A., Cox, P. M., Collins, M., Harris, P. P., Huntingford, C., & Jones, C. D. (2004). The role of ecosystem-atmosphere interactions in simulated Amazonian precipitation decrease and forest dieback under global climate warming. *Theoretical and applied climatology*, 78, 157-175.
- Beuning, K. R., Talbot, M. R., & Kelts, K. (1997). A revised 30,000-year paleoclimatic and paleohydrologic history of Lake Albert, East Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 136(1-4), 259-279.
- Beyene, T., Ludwig, F., & Franssen, W. H. P. (2012). The potential consequences of climate change in hydrology regime of the Congo River Basin. In *Climate change scenarios for the Congo Basin* (No. 11, pp. 59-104). Climate Service Centre.
- Binkley, D. A. N., & Giardina, C. (1998). Why do tree species affect soils? The warp and woof of tree-soil interactions. *Plant-induced soil changes: Processes and feedbacks*, 89-106.
- Bradley, R. (2000). 1000 years of climate change. *Science*, 288(5470), 1353-1355.

46. Bradley, B. J., Robbins, M. M., Williamson, E. A., Steklis, H. D., Steklis, N. G., Eckhardt, N., ... & Vigilant, L. (2005). Mountain gorilla tug-of-war: silverbacks have limited control over reproduction in multimale groups. *Proceedings of the National Academy of Sciences*, 102(26), 9418-9423.
47. Brando, P. M., Goetz, S. J., Baccini, A., Nepstad, D. C., Beck, P. S., & Christman, M. C. (2010). Seasonal and interannual variability of climate and vegetation indices across the Amazon. *Proceedings of the National Academy of Sciences*, 107(33), 14685-14690.
48. Breuer, T., Robbins, A. M., Olejniczak, C., Parnell, R. J., Stokes, E. J., & Robbins, M. M. (2010). Variance in the male reproductive success of western gorillas: acquiring females is just the beginning. *Behavioral Ecology and Sociobiology*, 64, 515-528.
49. Brncic, T. M., Willis, K. J., Harris, D. J., & Washington, R. (2007). Culture or climate? The relative influences of past processes on the composition of the lowland Congo rainforest. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 362(1478), 229-242.
50. Broecker, W. S. (2002). Was the Medieval Warm Period Global? Climate Change: Evaluating recent and future climate change, 4, 270.
51. Brown, H. C. P., Nkem, J. N., Sonwa, D. J., & Bele, Y. (2010). Institutional adaptive capacity and climate change response in the Congo Basin forests of Cameroon. Mitigation and adaptation strategies for global change, 15(3), 263-282.
52. Brown, S., Kebede, A. S., & Nicholls, R. J. (2011). Sea-level rise and impacts in Africa, 2000 to 2100. School of Civil Engineering and the Environment University of Southampton, UK.
53. Bodin, Ö., Tengö, M., Norman, A., Lundberg, J., & Elmqvist, T. (2006). The value of small size: loss of forest patches and ecological thresholds in southern Madagascar. *Ecological applications*, 16(2), 440-451.
54. Bonan, G. B. (2008). Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefits of Forests. *Science*, 320(5882), 1444-1449.
55. Bond, W. J., & Midgley, G. F. (2012). Carbon dioxide and the uneasy interactions of trees and savannah grasses. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1588), 601-612.
56. Bond, W. J., Midgley, G. F., & Woodward, F. I. (2003). The importance of low atmospheric CO₂ and fire in promoting the spread of grasslands and savannas. *Global Change Biology*, 9(7), 973-982.
57. Bonnefille, R., Roeland, J. C., & Guiot, J. (1990). Temperature and rainfall estimates for the past 40,000 years in equatorial Africa. *Nature*, 346(6282), 347-349.
58. Bonnefille, R., & Chalieu, F. (2000). Pollen-inferred precipitation time-series from equatorial mountains, Africa, the last 40 kyr BP. *Global and Planetary Change*, 26(1-3), 25-50.
59. Bouchet, H., Blois-Heulin, C., Pellier, A. S., Zuberbühler, K., & Lemasson, A. (2012). Acoustic variability and individual distinctiveness in the vocal repertoire of red-capped mangabeys (*Cercocebus torquatus*). *Journal of Comparative Psychology*, 126(1), 45.
60. Burghardt, G. M., Bartmess-LeVasseur, J. N., Browning, S. A., Morrison, K. E., Stec, C. L., Zachau, C. E., & Freeberg, T. M. (2012). Perspectives-minimizing observer bias in behavioral studies: a review and recommendations. *Ethology*, 118(6), 511-517.
61. Burns, C. E., Johnston, K. M., & Schmitz, O. J. (2003). Global climate change and mammalian species diversity in US national parks. *Proceedings of the National Academy of Sciences*, 100(20), 11474-11477.
62. Byron, N., & Arnold, M. (1999). What futures for the people of the tropical forests? World development, 27(5), 789-805.
63. Campbell, B. M. (2009). Beyond Copenhagen: REDD+, agriculture, adaptation strategies and poverty. *Global Environmental Change*, 19(4), 397-399.
64. Carpenter, S. R., DeFries, R., Dietz, T., Mooney, H. A., Polasky, S., Reid, W. V., & Scholes, R. J. (2006). Millennium ecosystem assessment: research needs. *Science*, 314(5797), 257-258.
65. Carpenter, S. R., Mooney, H. A., Agard, J., Capistrano, D., DeFries, R. S., Diaz, S., ... & Whyte, A. (2009). Science for managing ecosystem services: Beyond the Millennium Ecosystem Assessment. *Proceedings of the National Academy of Sciences*, 106(5), 1305-1312.
66. Carvalho, J. S., Graham, B., Bocksberger, G., Maisels, F., Williamson, E. A., Wich, S., ... & Kühl, H. S. (2021). Predicting range shifts of African apes under global change scenarios. *Diversity and Distributions*, 27(9), 1663-1679.
67. Carvalho, J. S., Graham, B., Rebelo, H., Bocksberger, G., Meyer, C. F., Wich, S., & Kühl, H. S. (2019). A global risk assessment of primates under climate and land use/cover scenarios. *Global Change Biology*, 25(9), 3163-3178.
68. Casimir, M. J. (1975). Feeding ecology and nutrition of an eastern gorilla group in the Mt. Kahuzi region (Republique du Zaïre). *Folia Primatologica*, 24(2-3), 81-136.
69. Ceglár, A., Zampieri, M., Toreti, A., & Dentener, F. (2019). Observed northward migration of agro-climate zones in Europe will further accelerate under climate change. *Earth's Future*, 7(9), 1088-1101.
70. Cerling, T. E., Harris, J. M., MacFadden, B. J., Leakey, M. G., Quade, J., Eisenmann, V., & Ehleringer, J. R. (1997). Global vegetation change through the Miocene/Pliocene boundary. *Nature*, 389(6647), 153-158.
71. Chen, I. C., Hill, J. K., Ohlemüller, R., Roy, D. B., & Thomas, C. D. (2011). Rapid range shifts of species associated with high levels of climate warming. *Science*, 333(6045), 1024-1026.
72. Chen, R., Little, R., Mihaylova, L., Delahay, R., & Cox, R. (2019). Wildlife surveillance using deep learning methods. *Ecology and evolution*, 9(17), 9453-9466.
73. Chilunjika, A. (2024). Understanding the Roots of the Recurrent Eastern Democratic Republic of Congo (DRC) Conflict. *African Renaissance*, 21(4), 53.
74. Chomitz, K. M., Alger, K., Thomas, T. S., Orlando, H., & Nova, P. V. (2005). Opportunity costs of conservation in a biodiversity hotspot: the case of southern Bahia. *Environment and Development Economics*, 10(3), 293-312.
75. Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B., ... & McCabe, A. M. (2009a). The last glacial maximum. *Science*, 325(5941), 710-714.
76. Clark, C. J., Poulsen, J. R., Malonga, R., & ELKAN, Jr, P. W. (2009b). Logging concessions can extend the conservation estate for Central African tropical forests. *Conservation biology*, 23(5), 1281-1293.
77. Cochrane, M. A., Alencar, A., Schulze, M. D., Souza Jr, C. M., Nepstad, D. C., Lefebvre, P., & Davidson, E. A. (1999). Positive feedbacks in the fire dynamic of closed canopy tropical forests. *Science*, 284(5421), 1832-1835.
78. Cohen, K.M., Harper, D.A.T., Gibbard, P.L. (2023). ICS International Chronostratigraphic Chart 2023/09. International Commission on Stratigraphy, IUGS. Retrieved May 13, 2024, from www.stratigraphy.org
79. Cramer, W., Bondeau, A., Schaphoff, S., Lucht, W., Smith, B., & Stith, S. (2004). Tropical forests and the global carbon cycle: impacts of atmospheric carbon dioxide, climate change and rate of deforestation. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 359(1443), 331-343.
80. Crick, H., Crosher, I., Mainstone, C., Taylor, S., Wharton, A., Langford, P., ... & Macgregor, N. (2020). Nature networks evidence handbook.
81. Conway, D., Allison, E., Felstead, R., & Goulden, M. (2005). Rainfall variability in East Africa: implications for natural resources management and livelihoods. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 363(1826), 49-54.
82. Costello, A., Abbas, M., Allen, A., Ball, S., Bell, S., Bellamy, R., ... & Patterson, C. (2009). Managing the health effects of climate change: lancet and University College London Institute for Global Health Commission. *The Lancet*, 373(9676), 1693-1733.
83. Cowie, A., Schneider, U. A., & Montanarella, L. (2007). Potential synergies between existing multilateral environmental agreements in the implementation of land use, land-use change and forestry activities. *environmental science & policy*, 10(4), 335-352.
84. Cruncheon, A. S., Borchers, D., Kühl, H., & Piel, A. (2020). Listening and watching: Do camera traps or acoustic sensors more efficiently detect wild chimpanzees in an open habitat? *Methods in Ecology and Evolution*, 11(4), 542-552.
85. Curtis, P. G., Slay, C. M., Harris, N. L., Tyukavina, A., & Hansen, M. C. (2018). Classifying drivers of global forest loss. *Science*, 361(6407), 1108-1111.
86. Damania, R., & Wheeler, D. (2015). Road improvement and deforestation in the Congo Basin countries. *World Bank Policy Research Working Paper*, (7274).
87. Damsté, J. S. S., Verschuren, D., Ossebaer, J., Blokker, J., van Houten, R., van der Meer, M. T., ... & Schouten, S. (2011). A 25,000-year record of climate-induced changes in lowland vegetation of eastern equatorial Africa revealed by the stable carbon-isotopic composition of fossil plant leaf waxes. *Earth and Planetary Science Letters*, 302(1-2), 236-246.
88. Dasgupta, P. (2001). *Human well-being and the natural environment*. Oxford University Press.
89. Deléue, M. A., Fuhr, M., Schwartz, D., Mariotti, A., & Nasi, R. (2001). Recent origin of a large part of the forest cover in the Gabon coastal area based on stable carbon isotope data. *Oecologia*, 129, 106-113.
90. Delire, C., Ngomanda, A., & Jolly, D. (2008). Possible impacts of 21st century climate on vegetation in Central and West Africa. *Global and Planetary Change*, 64(1-2), 3-15.
91. de Menocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L., & Yarusinsky, M. (2000). Abrupt onset and termination of the African Humid Period: rapid climate responses to gradual insolation forcing. *Quaternary science reviews*, 19(1-5), 347-361.
92. Dobson, A. P., Rodriguez, J. P., Roberts, W. M., & Wilcove, D. S. (1997). Geographic distribution of endangered species in the United

- States. *Science*, 275(5299), 550-553.
93. Doran, D. M., & McNeillage, A. (1998). Gorilla ecology and behavior. *Evolutionary Anthropology: Issues, News, and Reviews: Issues, News, and Reviews*, 6(4), 120-131.
 94. Doran, D. M., & McNeillage, A. (2001). Subspecific variation in gorilla behavior: the influence of ecological and social factors. In Robbins, M. M., Sicotte, P., Stewart K. J. (Eds.), *Mountain Gorillas: Three Decades of Research at Karisoke*. Cambridge University Press, 123-150.
 95. Douadi, M. I., Gatti, S., Levvero, F., Duhamel, G., Bermejo, M., Vallet, D., ... & Petit, E. J. (2007). Sex-biased dispersal in western lowland gorillas (*Gorilla gorilla gorilla*). *Molecular Ecology*, 16(11), 2247-2259.
 96. Drake, J. B., Dubayah, R. O., Clark, D. B., Knox, R. G., Blair, J. B., Hofton, M. A., ... & Prince, S. (2002). Estimation of tropical forest structural characteristics using large-footprint lidar. Remote sensing of environment, 79(2-3), 305-319.
 97. Drake, N. A., Blench, R. M., Armitage, S. J., Bristow, C. S., & White, K. H. (2011). Ancient watercourses and biogeography of the Sahara explain the peopling of the desert. *Proceedings of the National Academy of Sciences*, 108(2), 458-462.
 98. Dunn, A., Bergl, R., Byler, D., Eben-Ebai, S., Etiendem, D. N., Fotso, R., ... & Williamson, E. A. (2014). *Revised regional action plan for the conservation of the Cross River Gorilla (Gorilla gorilla diehli)* 2014–2019. IUCN/SSC Primate Specialist Group and Wildlife Conservation Society.
 99. Edwards, E. J., Osborne, C. P., Strömberg, C. A., Smith, S. A., C4 Grasses Consortium, Bond, W. J., ... & Tipple, B. (2010). The origins of C4 grasslands: integrating evolutionary and ecosystem science. *science*, 328(5978), 587-591.
 100. Edwards, D. P., Sloan, S., Weng, L., Dirks, P., Sayer, J., & Laurance, W. F. (2014a). Mining and the African environment. *Conservation Letters*, 7(3), 302-311.
 101. Edwards, D. P., Tobias, J. A., Shell, D., Meijaard, E., & Laurance, W. F. (2014b). Maintaining ecosystem function and services in logged tropical forests. *Trends in ecology & evolution*, 29(9), 511-520.
 102. Ehleringer, J. R., Cerling, T. E., & Helliker, B. R. (1997). C 4 photosynthesis, atmospheric CO 2, and climate. *Oecologia*, 112, 285-299.
 103. Elenga, H., Maley, J., Vincens, A., & Farrera, I. (2004). Palaeoenvironments, palaeoclimates and landscape development in Atlantic Equatorial Africa: a review of key sites covering the last 25 kyrs. Past climate variability through Europe and Africa, 181-198.
 104. Elenga, H., Schwartz, D., & Vincens, A. (1994). Pollen evidence of late Quaternary vegetation and inferred climate changes in Congo. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 109(2-4), 345-356.
 105. Enari, H., Enari, H. S., Okuda, K., Maruyama, T., & Okuda, K. N. (2019). An evaluation of the efficiency of passive acoustic monitoring in detecting deer and primates in comparison with camera traps. *Ecological Indicators*, 98, 753-762.
 106. Enfors, E. I., & Gordon, L. J. (2007). Analysing resilience in dryland agro-ecosystems: A case study of the Makanya catchment in Tanzania over the past 50 years. *Land degradation & development*, 18(6), 680-696.
 107. Enfors, E. I., Gordon, L. J., Peterson, G. D., & Bossio, D. (2008). Making investments in dryland development work: participatory scenario planning in the Makanya catchment, Tanzania. *Ecology and Society*, 13(2).
 108. Engel, V., Jobbágy, E. G., Stieglitz, M., Williams, M., & Jackson, R. B. (2005). Hydrological consequences of Eucalyptus afforestation in the Argentine Pampas. *Water Resources Research*, 41(10).
 109. Enkvist, P., Nauclér, T., & Rosander, J. (2007). A cost curve for greenhouse gas reduction. *McKinsey Quarterly*, 1, 34.
 110. Enquist, B. J., & Enquist, C. A. (2011). Long-term change within a Neotropical forest: Assessing differential functional and floristic responses to disturbance and drought. *Global Change Biology*, 17(3), 1408-1424.
 111. Ernst, C., Mayaux, P., Verhegghen, A., Bodart, C., Christophe, M., & Defourny, P. (2013). National forest cover change in Congo Basin: deforestation, reforestation, degradation and regeneration for the years 1990, 2000 and 2005. *Global change biology*, 19(4), 1173-1187.
 112. Estrada, A., Garber, P. A., Mittermeier, R. A., Wich, S., Gouveia, S., Dobrovolski, R., ... & Setiawan, A. (2018). Primates in peril: the significance of Brazil, Madagascar, Indonesia and the Democratic Republic of the Congo for global primate conservation. *PeerJ*, 6, e4869.
 113. Fa, J. E., & Brown, D. (2009). Impacts of hunting on mammals in African tropical moist forests: a review and synthesis. *Mammal Review*, 39(4).
 114. Fa, J. E., Peres, C. A., & Meeuwig, J. (2002). Bushmeat exploitation in tropical forests: an intercontinental comparison. *Conservation biology*, 16(1), 232-237.
 115. Fa, J. E., Yuste, J. E. G., & Castelo, R. (2000). Bushmeat markets on Bioko Island as a measure of hunting pressure. *Conservation biology*, 14(6), 1602-1613.
 116. Fahey, B., & Jackson, R. (1997). Hydrological impacts of converting native forests and grasslands to pine plantations, South Island, New Zealand. *Agricultural and Forest Meteorology*, 84(1-2), 69-82.
 117. Fauset, S., Baker, T. R., Lewis, S. L., Feldpausch, T. R., Affum-Baffoe, K., Folli, E. G., ... & Swaine, M. D. (2012). Drought-induced shifts in the floristic and functional composition of tropical forests in Ghana. *Ecology letters*, 15(10), 1120-1129.
 118. Feintrenie, L. (2014). Agro-industrial plantations in Central Africa, risks and opportunities. *Biodiversity and conservation*, 23(6), 1577-1589.
 119. Ferraro, P. J. (2003). Assigning priority to environmental policy interventions in a heterogeneous world. *Journal of Policy Analysis and Management*, 22(1), 27-43.
 120. Fimbel, R. A., Robinson, J., & Grajal, A. (Eds.). (2001). *The cutting edge: conserving wildlife in logged tropical forests*. Columbia University Press.
 121. Fisher, B., Turner, R. K., & Morling, P. (2009). Defining and classifying ecosystem services for decision making. *Ecological economics*, 68(3), 643-653.
 122. Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., ... & Snyder, P. K. (2005). Global consequences of land use. *science*, 309(5734), 570-574.
 123. Fossey, D. (1974). Observations on the home range of one group of mountain gorillas (*Gorilla gorilla beringel*). *Animal Behaviour*, 22(3), 568-581.
 124. Frelich, L. E., Montgomery, R. A., & Oleksyn, J. (2015). Northern temperate forests. In *Routledge handbook of forest ecology* (pp. 30-45). Routledge.
 125. Fünfstück, T., & Vigilant, L. (2015). The geographic distribution of genetic diversity within gorillas. *American Journal of Primatology*, 77(9), 974-985.
 126. Fung, I. Y., Doney, S. C., Lindsay, K., & John, J. (2005). Evolution of carbon sinks in a changing climate. *Proceedings of the National Academy of Sciences*, 102(32), 11201-11206.
 127. Garner, K. J., & Ryder, O. A. (1996). Mitochondrial DNA diversity in gorillas. *Molecular Phylogenetics and Evolution*, 6(1), 39-48.
 128. Gasse, F. (2000). Hydrological changes in the African tropics since the Last Glacial Maximum. *Quaternary Science Reviews*, 19(1-5), 189-211.
 129. Gibbs, H. K., Ruesch, A. S., Achard, F., Clayton, M. K., Holmgren, P., Ramankutty, N., & Foley, J. A. (2010). Tropical forests were the primary sources of new agricultural land in the 1980s and 1990s. *Proceedings of the National Academy of Sciences*, 107(38), 16732-16737.
 130. Gibson, L., Lynam, A. J., Bradshaw, C. J., He, F., Bickford, D. P., Woodruff, D. S., ... & Laurance, W. F. (2013). Near-complete extinction of native small mammal fauna 25 years after forest fragmentation. *Science*, 341(6153), 1508-1510.
 131. Giorgi, F. (2002). Variability and trends of sub-continental scale surface climate in the twentieth century. Part I: observations. *Climate Dynamics*, 18(8), 675-691.
 132. Giresse, P., Maley, J., & Brenac, P. (1994). Late Quaternary palaeoenvironments in the Lake Barombi Mbo (West Cameroon) deduced from pollen and carbon isotopes of organic matter. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 107(1-2), 65-78.
 133. Giresse, P., Maley, J., & Kossioni, A. (2005). Sedimentary environmental changes and millennial climatic variability in a tropical shallow lake (Lake Ossa, Cameroon) during the Holocene. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 218(3-4), 257-285.
 134. Goetz, S. J., Hansen, M., Houghton, R. A., Walker, W., Laporte, N., & Busch, J. (2015). Measurement and monitoring needs, capabilities and potential for addressing reduced emissions from deforestation and forest degradation under REDD+. *Environmental Research Letters*, 10(12), 123001.
 135. Goldman, M. (2001). Constructing an environmental state: eco-governmentality and other transnational practices of a 'green'World Bank. *Social problems*, 48(4), 499-523.
 136. Goodall, A. G., & Groves, C. P. (1977). The conservation of eastern gorillas. In Prince Rainier III H. R. H., Bourne G. H. (Eds.), *Primate conservation*. NY: Academic Press, 599-637.
 137. Gordon, L. J., Peterson, G. D., & Bennett, E. M. (2008). Agricultural modifications of hydrological flows create ecological surprises. *Trends in ecology & evolution*, 23(4), 211-219.
 138. Graham, C. H., VanDerWal, J., Phillips, S. J., Moritz, C., & Williams, S. E. (2010). Dynamic refugia and species persistence: tracking spatial shifts in habitat through time. *Ecography*, 33(6), 1062-1069.

139. Granjon, A. C., Robbins, M. M., Arinaitwe, J., Cranfield, M. R., Eckardt, W., Mburanumwe, I., ... & Hickey, J. R. (2020). Estimating abundance and growth rates in a wild mountain gorilla population. *Animal Conservation*, 23(4), 455-465.
140. Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., ... & Townshend, J. R. (2015). Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science advances*, 1(2), e1500052.
141. Hamilton AC. (1982). Environmental history of East Africa: a study of the Quaternary. London: Academic Press.
142. Hamilton, A. C., & Taylor, D. (1991). History of climate and forests in tropical Africa during the last 8 million years. *Climatic change*, 19(1), 65-78.
143. Harcourt, A. H. (1979). Social relationships among adult female mountain gorillas. *Animal Behaviour*, 27, 251-264.
144. Harcourt, A. H. (1996). Is the gorilla a threatened species? How should we judge?. *Biological Conservation*, 75(2), 165-176.
145. Harilal, V., Tichaawa, T. M., & Saarinen, J. (2019). "Development without policy": Tourism planning and research needs in Cameroon, Central Africa. *Tourism Planning & Development*, 16(6), 696-705.
146. Harris, J. (1993). Ecosystem structure and growth of the African savanna. *Global and planetary change*, 8(4), 231-248.
147. Hayden, B. P. (1998). Ecosystem feedbacks on climate at the landscape scale. Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences, 353(1365), 5-18.
148. Hein, L., Van Koppen, K., De Groot, R. S., & Van Ierland, E. C. (2006). Spatial scales, stakeholders and the valuation of ecosystem services. *Ecological economics*, 57(2), 209-228.
149. Henley, J. (2003, November, 13). Gigantic sleaze scandal winds up as former Elf oil chiefs are jailed. Retrieved from <https://www.theguardian.com/business/2003/nov/13/france.oilandpetrol>
150. Hernández-Ruedas, M. A., Arroyo-Rodríguez, V., Meave, J. A., Martínez-Ramos, M., Ibarra-Manriquez, G., Martínez, E., ... & Santos, B. A. (2014). Conserving tropical tree diversity and forest structure: the value of small rainforest patches in moderately-managed landscapes. *PLoS one*, 9(6), e98931.
151. Hewlett, B. S. (2000). Central African governments' and international NGOs' perceptions of Baka Pygmy development. In Shweitzer, P. P., Bieseke, M., & Hitchcock, R. K. (Eds.), *Hunters and gatherers in the modern world: Conflict, resistance, and self-determination*, 380-390. Berghahn Books.
152. Hijmans, R. J., & Graham, C. H. (2006). The ability of climate envelope models to predict the effect of climate change on species distributions. *Global change biology*, 12(12), 2272-2281.
153. Hoare, A. L. (2007). The use of non-timber forest products in the Congo Basin: constraints and opportunities. London, UK: Rainforest Foundation.
154. Hoare, A. (2015). Tackling illegal logging and the related trade: *What progress and where next*. The Royal Institute of International Affairs.
155. Hodgson, J. A., Wallis, D. W., Krishna, R., & Cornell, S. J. (2016). How to manipulate landscapes to improve the potential for range expansion. *Methods in Ecology and Evolution*, 7(12), 1558-1566.
156. Hu, T., Su, Y., Xue, B., Liu, J., Zhao, X., Fang, J., & Guo, Q. (2016). Mapping global forest aboveground biomass with spaceborne LiDAR, optical imagery, and forest inventory data. *Remote Sensing*, 8(7), 565.
157. Huete, A. R., Didan, K., Shimabukuro, Y. E., Ratana, P., Saleska, S. R., Hutya, L. R., ... & Myneni, R. (2006). Amazon rainforests green-up with sunlight in dry season. *Geophysical research letters*, 33(6).
158. Hulme, M., Doherty, R., Ngara, T., New, M., & Lister, D. (2001). African climate change: 1900-2100. *Climate research*, 17(2), 145-168.
159. Igoe, J., & Brockington, D. (2007). Neoliberal conservation: a brief introduction. *Conservation and society*, 5(4), 432-449.
160. Jackson, D., & Payne, K. (2003). *Twa women, Twa rights in the Great Lakes region of Africa*. London, United Kingdom: Minority Rights Group International.
161. James, R., & Washington, R. (2013). Changes in African temperature and precipitation associated with degrees of global warming. *Climatic change*, 117, 859-872.
162. James, R., Washington, R., & Rowell, D. P. (2013). Implications of global warming for the climate of African rainforests. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1625), 1-8.
163. Jensen-Seaman, M. I., & Kidd, K. K. (2001). Mitochondrial DNA variation and biogeography of eastern gorillas. *Molecular Ecology*, 10(9), 2241-2247.
164. Johns, A. G. (1997). *Timber production and biodiversity conservation in tropical rain forests*. Cambridge University Press.
165. Joppa, L. N. (2015). Technology for nature conservation: An industry perspective. *Ambio*, 44(Suppl 4), 522-526.
166. Junker, J., Blake, S., Boesch, C., Campbell, G., Toit, L. D., Duvall, C., ... & Kuehl, H. S. (2012). Recent decline in suitable environmental conditions for African great apes. *Diversity and Distributions*, 18(11), 1077-1091.
167. Kageyama, M., Braconnot, P., Bopp, L., Mariotti, V., Roy, T., Woillez, M. N., ... & Marti, O. (2013). Mid-Holocene and last glacial maximum climate simulations with the IPSL model: part II: model-data comparisons. *Climate dynamics*, 40, 2469-2495.
168. Kalan, A. K., Mundry, R., Wagner, O. J., Heinicke, S., Boesch, C., & Kühl, H. S. (2015). Towards the automated detection and occupancy estimation of primates using passive acoustic monitoring. *Ecological Indicators*, 54, 217-226.
169. Kareiva, P., Watts, S., McDonald, R., & Boucher, T. (2007). Domesticated nature: shaping landscapes and ecosystems for human welfare. *Science*, 316(5833), 1866-1869.
170. Karsenty, A., & Ferron, C. (2017). Recent evolutions of forest concessions status and dynamics in Central Africa. *International Forestry Review*, 19(4), 10-26.
171. Kelly, M. A., Russell, J. M., Baber, M. B., Howley, J. A., Loomis, S. E., Zimmerman, S., ... & Lukaye, J. (2014). Expanded glaciers during a dry and cold Last Glacial Maximum in equatorial East Africa. *Geology*, 42(6), 519-522.
172. Khadaroo, A. J., & Seetanah, B. (2007). Research note: Does transport infrastructure matter in overall tourism development? Evidence from a sample of island economies. *Tourism Economics*, 13(4), 675-687.
173. Kimbu, A. N. (2011). The challenges of marketing tourism destinations in the Central African subregion: The Cameroon example. *International Journal of Tourism Research*, 13(4), 324-336.
174. King, D. A. (2004). Climate change science: adapt, mitigate, or ignore?. *Science* 303(5655), 176-177.
175. King, M., Altdorff, D., Li, P., Galagedara, L., Holden, J., & Unc, A. (2018). Northward shift of the agricultural climate zone under 21st-century global climate change. *Scientific reports*, 8(1), 7904.
176. Klein, R. J., Schipper, E. L. F., & Dessai, S. (2005). Integrating mitigation and adaptation into climate and development policy: three research questions. *Environmental science & policy*, 8(6), 579-588.
177. Kleinschroth, F., & Healey, J. R. (2017). Impacts of logging roads on tropical forests. *Biotropica*, 49(5), 620-635.
178. Kobashi, T., Severinghaus, J. P., & Barnola, J. M. (2008). 4±1.5 C abrupt warming 11,270 yr ago identified from trapped air in Greenland ice. *Earth and Planetary Science Letters*, 268(3-4), 397-407.
179. Koster, R. D., Dirmeyer, P. A., Guo, Z., Bonan, G., Chan, E., Cox, P., ... & Yamada, T. (2004). Regions of strong coupling between soil moisture and precipitation. *Science*, 305(5687), 1138-1140.
180. Kremen, C., Niles, J. O., Dalton, M. G., Daily, G. C., Ehrlich, P. R., Fay, J. P., ... & Guillery, R. P. (2000). Economic incentives for rain forest conservation across scales. *Science*, 288(5472), 1828-1832.
181. Kremen, C., Williams, N. M., Aizen, M. A., Gemmill-Herren, B., LeBuhn, G., Minckley, R., ... & Ricketts, T. H. (2007). Pollination and other ecosystem services produced by mobile organisms: a conceptual framework for the effects of land-use change. *Ecology letters*, 10(4), 299-314.
182. Kremen, C., & Ostfeld, R. S. (2005). A call to ecologists: measuring, analyzing, and managing ecosystem services. *Frontiers in Ecology and the Environment*, 3(10), 540-548.
183. Krosby, M., Breckheimer, I., John Pierce, D., Singleton, P. H., Hall, S. A., Halupka, K. C., ... & Schuett-Hames, J. P. (2015). Focal species and landscape "naturalness" corridor models offer complementary approaches for connectivity conservation planning. *Landscape ecology*, 30, 2121-2132.
184. Kühl, H. S., Sop, T., Williamson, E. A., Mundry, R., Brugière, D., Campbell, G., ... & Boesch, C. (2017). The Critically Endangered western chimpanzee declines by 80%. *American Journal of Primatology*, 79(9), e22681.
185. Kümpel, N. F., Milner-Gulland, E. J., Cowlshaw, G., & Rowcliffe, J. M. (2010). Incentives for hunting: the role of bushmeat in the household economy in rural Equatorial Guinea. *Human Ecology*, 38, 251-264.
186. Lahm, S. A., Barnes, R. F. W., Beardsley, K., & Cervinka, P. (1998). A method for censusing the greater white-nosed monkey in northeastern Gabon using the population density gradient in relation to roads. *Journal of Tropical Ecology*, 14(5), 629-643.
187. Lahm, S. A. (2001). Hunting and wildlife in Northeastern Gabon: why conservation should extend beyond protected areas. In Weber, W., White, L. J. T., Vedder, A., & Treves-Naughton, L., (Eds.), *African Rain Forest Ecology and Conservation* (344-354). CT: Yale University Press.
188. Lang, N., Schindler, K., & Wegner, J. D. (2021). High carbon stock mapping at large scale with optical satellite imagery and spaceborne LiDAR. *arXiv preprint arXiv:2107.07431*.

189. Laurance, W. F., Sayer, J., & Cassman, K. G. (2014). Agricultural expansion and its impacts on tropical nature. *Trends in ecology & evolution*, 29(2), 107-116.
190. Laurance, W. F. (1999). Reflections on the tropical deforestation crisis. *Biological conservation*, 91(2-3), 109-117.
191. Laurance, W. F., Alonso, A., Lee, M., & Campbell, P. (2006). Challenges for forest conservation in Gabon, Central Africa. *Futures*, 38(4), 454-470.
192. Laurance, W. F. (2001a). Tropical logging and human invasions. *Conservation Biology*, 15(1), 4-5.
193. Laurance, W. F., Goosem, M., & Laurance, S. G. (2009). Impacts of roads and linear clearings on tropical forests. *Trends in ecology & evolution*, 24(12), 659-669.
194. Laurance, W. F. (2001b). Immense logging deal to sustain war in the Congo. *Trends in Ecology & Evolution*, 16(12), 670.
195. Laurance, W. F., & Laurance, S. G. (1996). Responses of five arboreal marsupials to recent selective logging in tropical Australia. *Biotropica*, 310-322.
196. Lehmann, C. E., Archibald, S. A., Hoffmann, W. A., & Bond, W. J. (2011). Deciphering the distribution of the savanna biome. *New Phytologist*, 191(1), 197-209.
197. Leroy, E. M., Rouquet, P., Formenty, P., Souquiere, S., Kilbourne, A., Froment, J. M., ... & Rollin, P. E. (2004). Multiple Ebola virus transmission events and rapid decline of central African wildlife. *Science*, 303(5656), 387-390.
198. Lewis, S. L. (2006). Tropical forests and the changing earth system. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 361(1465), 195-210.
199. Lewis, S. L., Lloyd, J., Sith, S., Mitchard, E. T., & Laurance, W. F. (2009). Changing ecology of tropical forests: evidence and drivers. *Annual Review of Ecology, Evolution, and Systematics*, 40(1), 529-549.
200. Lewis, J. (2000). The Batwa pygmies of the great lakes region (Vol. 209). Minority Rights Group International.
201. Lewis, S. L., Edwards, D. P., & Galbraith, D. (2015). Increasing human dominance of tropical forests. *Science*, 349(6250), 827-832.
202. Lewis, S. L., Sonké, B., Sunderland, T., Begne, S. K., Lopez-Gonzalez, G., Van Der Heijden, G. M., ... & Zemagho, L. (2013). Above-ground biomass and structure of 260 African tropical forests. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1625), 20120295.
203. Lézine, A. M., Zheng, W., Braconnot, P., & Krinner, G. (2011). Late Holocene plant and climate evolution at Lake Yoa, northern Chad: pollen data and climate simulations. *Climate of the Past*, 7(4), 1351-1362.
204. Limburg, K. E., O'Neill, R. V., Costanza, R., & Farber, S. (2002). Complex systems and valuation. *Ecological economics*, 41(3), 409-420.
205. Lindenmayer, D. B., Laurance, W. F., & Franklin, J. F. (2012). Global decline in large old trees. *Science*, 338(6112), 1305-1306.
206. Livingstone, D. A. (1967). Postglacial vegetation of the Ruwenzori Mountains in equatorial Africa. *Ecological Monographs*, 37(1), 25-52.
207. Lloyd, J., & Farquhar, G. D. (2008). Effects of rising temperatures and [CO₂] on the physiology of tropical forest trees. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1498), 1811-1817.
208. Locatelli, B., Kanninen, M., Brockhaus, M., Colfer, C. P., Murdiyarso, D., & Santoso, H. (2008). Facing an uncertain future: How forests and people can adapt to climate change. Bogor Barat, Indonesia: Center for International Forestry Research.
209. Lucht, W., Schaphoff, S., Erbrecht, T., Heyder, U., & Cramer, W. (2006). Terrestrial vegetation redistribution and carbon balance under climate change. *Carbon balance and management*, 1, 1-7.
210. Lyon, C., Saupe, E. E., Smith, C. J., Hill, D. J., Beckerman, A. P., Stringer, L. C., ... & Aze, T. (2022). Climate change research and action must look beyond 2100. *Global Change Biology*, 28 (2), 349-361.
211. MacArthur, R. H., & Wilson, E. O. (2001). *The theory of island biogeography*. Princeton University Press.
212. Mailund, T., Halager, A. E., Westergaard, M., Duthiel, J. Y., Munch, K., Andersen, L. N., ... & Schierup, M. H. (2012). A new isolation with migration model along complete genomes infers very different divergence processes among closely related great ape species. *PLoS genetics*, 8(12), e1003125.
213. Maisels, F., Bergl, R. A., & Williamson, E. A. (2018). Gorilla gorilla (amended version of 2016 assessment). The IUCN Red List of Threatened Species 2018: e. T9404A136250858.
214. Maisels, F., Strindberg, S., Blake, S., Wittemyer, G., Hart, J., Williamson, E. A., ... & Warren, Y. (2013). Devastating decline of forest elephants in Central Africa. *PloS one*, 8(3), e59469.
215. Malcolm, J. R., & Ray, J. C. (2000). Influence of timber extraction routes on central African small-mammal communities, forest structure, and tree diversity. *Conservation Biology*, 14(6), 1623-1638.
216. Maley, J. (1996). The African rain forest—main characteristics of changes in vegetation and climate from the Upper Cretaceous to the Quaternary. *Proceedings of the Royal Society of Edinburgh, Section B: Biological Sciences*, 104, 31-73.
217. Maley, J. (2001). The impact of arid phases on the African rain forest through geological history. In Weber, W., White, L. J., Naughton-Traves, L., & Vedder, A. (Eds.), *African rain forest ecology and conservation: an interdisciplinary perspective*. Yale University Press, 68-87.
218. Maley, J. (1989). Late Quaternary climatic changes in the African rain forest: forest refugia and the major role of sea surface temperature variations. In Leinen, M., & Sarnthein, M. (Eds.), *Paleoclimatology and paleometeorology: modern and past patterns of global atmospheric transport* (585-616). Springer Dordrecht.
219. Maley, J. (1991). The African rain forest vegetation and palaeoenvironments during late Quaternary. *Climatic change*, 19(1), 79-98.
220. Maley, J., & Brenac, P. (1998). Vegetation dynamics, palaeoenvironments and climatic changes in the forests of western Cameroon during the last 28,000 years BP. Review of Palaeobotany and Palynology, 99(2), 157-187.
221. Maley J. (2002). A catastrophic destruction of African forests about 2500 years ago still exerts a major influence on present vegetation formations. *IDS Bulletin*. 33, 13 –30.
222. Malhi, Y., & Wright, J. (2004). Spatial patterns and recent trends in the climate of tropical rainforest regions. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 359(1443), 311-329.
223. Malhi, Y., Roberts, J. T., Betts, R. A., Killeen, T. J., Li, W., & Nobre, C. A. (2008). Climate change, deforestation, and the fate of the Amazon. *science*, 319(5860), 169-172.
224. Mancini, F., Hodgson, J. A., & Isaac, N. J. (2022). Co-designing an indicator of habitat connectivity for England. *Frontiers in Ecology and Evolution*, 10, 892987.
225. Margules, C. R., & Pressey, R. L. (2000). Systematic conservation planning. *Nature*, 405(6783), 243-253.
226. Martens, P., McEvoy, D., & Chang, C. (2009). The climate change challenge: linking vulnerability, adaptation, and mitigation. Current opinion in environmental sustainability, 1(1), 14-18.
227. Masi, S., Austerlitz, F., Chabaud, C., Lafosse, S., Marchi, N., Georges, M., ... & Ségurel, L. (2021). No evidence for female kin association, indications for extragroup paternity, and sex-biased dispersal patterns in wild western gorillas. *Ecology and Evolution*, 11(12), 7634-7646.
228. Mayaux, P., Bartholomé, E., Fritz, S., & Belward, A. (2004). A new land-cover map of Africa for the year 2000. *Journal of biogeography*, 31(6), 861-877.
229. Mayaux, P., Pekel, J. F., Desclée, B., Donnay, F., Lupi, A., Achard, F., ... & Belward, A. (2013). State and evolution of the African rainforests between 1990 and 2010. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1625), 20120300.
230. Mbenda, H. G. N., Awasthi, G., Singh, P. K., Gouado, I., & Das, A. (2014). Does malaria epidemiology project Cameroon as 'Africa in miniature'? *Journal of biosciences*, 39, 727-738.
231. McClean, C. J., Lovett, J. C., Küper, W., Hannah, L., Sommer, J. H., Barthlott, W., ... & Taplin, J. R. (2005). African plant diversity and climate change. *Annals of the Missouri Botanical Garden*, 139-152.
232. McEvoy, J. F., Hall, G. P., & McDonald, P. G. (2016). Evaluation of unmanned aerial vehicle shape, flight path and camera type for waterfowl surveys: disturbance effects and species recognition. *PeerJ*, 4, e1831.
233. McGlynn, G., Mooney, S., & Taylor, D. (2013). Palaeoecological evidence for Holocene environmental change from the Virunga volcanoes in the Albertine Rift, central Africa. *Quaternary Science Reviews*, 61, 32-46.
234. McManus, K. F., Kelley, J. L., Song, S., Veeramah, K. R., Woerner, A. E., Stevison, L. S., ... & Hammer, M. F. (2015). Inference of gorilla demographic and selective history from whole-genome sequence data. *Molecular biology and evolution*, 32(3), 600-612.
235. McNeillage, A., Robbins, M. M., Gray, M., Olupot, W., Babaasa, D., Bitariho, R., ... & Baker, J. (2006). Census of the mountain gorilla *Gorilla beringei beringei* population in Bwindi Impenetrable National Park, Uganda. *Oryx*, 40(4), 419-427.
236. Megevand, C. (2013). *Deforestation trends in the Congo Basin: reconciling economic growth and forest protection*. Washington, DC: World Bank.
237. Mehlman, P. T. (2008). Current status of wild gorilla populations and strategies for their conservation. *Conservation in the 21st century: Gorillas as a case study*, 3-54.
238. Midgley, G. F., & Thuiller, W. (2005). Global environmental change and the uncertain fate of biodiversity. *New Phytologist*, 638-641.
239. Mills, M. G. L., Biggs, H. C., & Whyte, I. J. (1995). The relationship between rainfall, lion predation and population trends in African herbivores. *Wildlife Research*, 22(1), 75-87.

240. Milner-Gulland, E. J., & Bennett, E. L. (2003). Wild meat: the bigger picture. *Trends in Ecology & Evolution*, 18(7), 351-357.
241. Moise, R. (2011). If Pygmies could talk: creating indigenous development in equatorial Africa. *Before Farming*, 2011, 4, 1-23.
242. Molinaro, G., Hansen, M. C., Potapov, P. V., Tyukavina, A., Stehman, S., Barker, B., & Humber, M. (2017). Quantification of land cover and land use within the rural complex of the Democratic Republic of Congo. *Environmental Research Letters*, 12(10), 104001.
243. Molua, E. L. (2007). The economic impact of climate change on agriculture in Cameroon. *World Bank Policy Research Working Paper*, (4364).
244. Monfreda, C., Ramankutty, N., & Foley, J. A. (2008). Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000. *Global biogeochemical cycles*, 22(1).
245. Mora, C., Metzger, R., Rollo, A., & Myers, R. A. (2007). Experimental simulations about the effects of overexploitation and habitat fragmentation on populations facing environmental warming. *Proceedings of the Royal Society B: Biological Sciences*, 274(1613), 1023-1028.
246. Morton, D. C., Nagol, J., Carabajal, C. C., Rosette, J., Palace, M., Cook, B. D., ... & North, P. R. (2014). Amazon forests maintain consistent canopy structure and greenness during the dry season. *Nature*, 506(7487), 221-224.
247. Motlagh, F., Hamideh, S., Gallagher, M., Yan, G., & van de Lindt, J. W. (2024). Bonds for disaster resilience: A review of literature and practice. *International Journal of Disaster Risk Reduction*, 104, Article 104318.
248. Montgomery, C. A., & Adams, D. M. (1994). The marginal cost of species preservation: the northern spotted owl. *Journal of Environmental Economics and Management*, 26(2), 111-128.
249. Morcatty, T. Q., El Bizri, H. R., Carneiro, H. C. S., Biasizzo, R. L., Alméri, C. R. D. O., Silva, E. S. D., ... & Figueira, J. E. C. (2013). Habitat loss and mammalian extinction patterns: are the reserves in the Quadriátero Ferrífero, southeastern Brazil, effective in conserving mammals?. *Ecological research*, 28, 935-947.
250. Murphy, P., Pritchard, M. P., & Smith, B. (2000). The destination product and its impact on traveller perceptions. *Tourism management*, 21(1), 43-52.
251. Naidoo, R., Balmford, A., Ferraro, P. J., Polasky, S., Ricketts, T. H., & Rouget, M. (2006). Integrating economic costs into conservation planning. *Trends in ecology & evolution*, 21(12), 681-687.
252. Naidoo, R., & Adamowicz, W. L. (2005). Economic benefits of biodiversity exceed costs of conservation at an African rainforest reserve. *Proceedings of the National Academy of Sciences*, 102(46), 16712-16716.
253. Nasi, R., Taber, A., & Van Vliet, N. (2011). Empty forests, empty stomachs? Bushmeat and livelihoods in the Congo and Amazon Basins. *International Forestry Review*, 13(3), 355-368.
254. Naughton-Treves, L., & Weber, W. (2001). Human dimensions of the African rain forest. In Weber, W., White L. J. T., Vedder, A., Naughton-Treves, L. (Eds.), *African Rain Forest Ecology and Conservation*. Yale University Press, 20-46.
255. Ndoye, O., Ruiz-Pérez, M., & Eyebe, A. (1998). The markets of non-timber forest products in the humid forest zone of Cameroon. *Rural Development Forestry Network Paper* 22c, 1-25.
256. Nemani, R. R., Keeling, C. D., Hashimoto, H., Jolly, W. M., Piper, S. C., Tucker, C. J., ... & Running, S. W. (2003). Climate-driven increases in global terrestrial net primary production from 1982 to 1999. *science*, 300(5625), 1560-1563.
257. Nepstad, D., Lefebvre, P., Lopes da Silva, U., Tomasella, J., Schlesinger, P., Solórzano, L., ... & Guerreira Benito, J. (2004). Amazon drought and its implications for forest flammability and tree growth: A basin-wide analysis. *Global change biology*, 10(5), 704-717.
258. Nepstad, D. C., Tohver, I. M., Ray, D., Moutinho, P., & Cardiot, G. (2007). Mortality of large trees and lianas following experimental drought in an Amazon forest. *Ecology*, 88(9), 2259-2269.
259. Nepstad, D. C., Verssimo, A., Alencar, A., Nobre, C., Lima, E., Lefebvre, P., ... & Brooks, V. (1999). Large-scale impoverishment of Amazonian forests by logging and fire. *Nature*, 398(6727), 505-508.
260. Neigh, C. S., Nelson, R. F., Ranson, K. J., Margolis, H. A., Montesano, P. M., Sun, G., ... & Andersen, H. E. (2013). Taking stock of circumboreal forest carbon with ground measurements, airborne and spaceborne LiDAR. *Remote Sensing of Environment*, 137, 274-287.
261. Neumann, K., Bostoen, K., Höhn, A., Kahlheber, S., Ngomanda, A., & Tchiengue, B. (2012). First farmers in the Central African rainforest: A view from southern Cameroon. *Quaternary International*, 249, 53-62.
262. Ngomanda, A., Jolly, D., Bentaleb, I., Chepstow-Lusty, A., Maley, J., Fontugne, M., ... & Rabenkogo, N. (2007). Lowland rainforest response to hydrological changes during the last 1500 years in Gabon, Western Equatorial Africa. *Quaternary research*, 67(3), 411-425.
263. Nguetsop, V. F., Servant-Vildary, S., & Servant, M. (2004). Late Holocene climatic changes in west Africa, a high resolution diatom record from equatorial Cameroon. *Quaternary Science Reviews*, 23(5-6), 591-609.
264. Nkem, J. N., Somorin, O. A., Jum, C., Idinoba, M. E., Bele, Y. M., & Sonwa, D. J. (2013). Profiling climate change vulnerability of forest indigenous communities in the Congo Basin. *Mitigation and Adaptation Strategies for Global Change*, 18(5), 513-533.
265. Nkem, J., Kalame, F. B., Idinoba, M., Somorin, O. A., Ndoye, O., & Awono, A. (2010). Shaping forest safety nets with markets: Adaptation to climate change under changing roles of tropical forests in Congo Basin. *environmental science & policy*, 13(6), 498-508.
266. Nogherotto, R., Coppola, E., Giorgi, F., & Mariotti, L. (2013). Impact of Congo Basin deforestation on the African monsoon. *Atmospheric Science Letters*, 14(1), 45-51.
267. Norby, R. J., DeLucia, E. H., Gielen, B., Calfapietra, C., Giardina, C. P., King, J. S., ... & Oren, R. (2005). Forest response to elevated CO₂ is conserved across a broad range of productivity. *Proceedings of the National Academy of Sciences*, 102(50), 18052-18056.
268. Oates, J. F., Abedi-Lartey, M., McGraw, W. S., Struhsaker, T. T., & Whitesides, G. H. (2000). Extinction of a West African red colobus monkey. *Conservation Biology*, 14(5), 1526-1532.
269. Oliver, T. H., & Morecroft, M. D. (2014). Interactions between climate change and land use change on biodiversity: attribution problems, risks, and opportunities. *Wiley Interdisciplinary Reviews: Climate Change*, 5(3), 317-335.
270. Olschewski, R., Tschardt, T., Benítez, P. C., Schwarze, S., & Klein, A. M. (2006). Economic evaluation of pollination services comparing coffee landscapes in Ecuador and Indonesia. *Ecology and Society*, 11(1).
271. Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V., Underwood, E. C., ... & Kassem, K. R. (2001). Terrestrial Ecoregions of the World: A New Map of Life on Earth: A new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity. *BioScience*, 51(11), 933-938.
272. Osborne, C. P. (2008). Atmosphere, ecology and evolution: what drove the Miocene expansion of C4 grasslands?. *Journal of Ecology*, 96(1), 35-45.
273. O'Sullivan, R. (2009). Reducing Emissions from Deforestation in Developing Countries: An Introduction. In Streck, C., O'Sullivan, R., Janson-Smith, T., & Tarasofsky, R. (Eds.), *Climate change and forests: emerging policy and market opportunities*, 179-190. Brookings Institution Press.
274. Owen-Smith, N. (1990). Demography of a large herbivore, the greater kudu *Tragelaphus strepsiceros*, in relation to rainfall. *The Journal of Animal Ecology*, 59, 893-913.
275. Pagani, M., Caldeira, K., Berner, R., & Beerling, D. J. (2009). The role of terrestrial plants in limiting atmospheric CO₂ decline over the past 24 million years. *Nature*, 460(7251), 85-88.
276. Pahl-Wostl, C. (2009). A conceptual framework for analysing adaptive capacity and multi-level learning processes in resource governance regimes. *Global environmental change*, 19(3), 354-365.
277. Pauw, W. P. (2025). The three missed opportunities of the UNFCCC's New Collective Quantified Goal on climate finance. *Earth System Governance*, 24, 100255.
278. Pearce, F. (2012). *Turning Point: What Future for Forest Peoples and Resources in the Emerging World Order*. Washington, DC: Rights and Resources Group.
279. Pelletier, J., Horning, N., Laporte, N., Samndong, R. A., & Goetz, S. (2018). Anticipating social equity impacts in REDD+ policy design: An example from the Democratic Republic of Congo. *Land Use Policy*, 75, 102-115.
280. Piel, A. K., Cruncheon, A., Knot, I. E., Chalmers, C., Fergus, P., Mulero-Pázmány, M., & Wich, S. A. (2021). Noninvasive technologies for primate conservation in the 21st century. *International Journal of Primatology*, 1-35.
281. Pimm, S. L., Alibhai, S., Bergl, R., Dehgan, A., Giri, C., Jewell, Z., ... & Loarie, S. (2015). Emerging technologies to conserve biodiversity. *Trends in ecology & evolution*, 30(11), 685-696.
282. Plumptre, A. J., Kirkby, A., Spira, C., Kivono, J., Mitamba, G., Ngoy, E., ... & Kujirakwinja, D. (2021). Changes in Grauer's gorilla (*Gorilla beringei graueri*) and other primate populations in the Kahuzi-Biega National Park and Oku Community Reserve, the heart of Grauer's gorilla global range. *American Journal of Primatology*, 83(7), e23288.
283. Plumptre, A. J., Nixon, S., Kujirakwinja, D. K., Vieilledent, G., Critchlow, R., Williamson, E. A., ... & Hall, J. S. (2016). Catastrophic decline of world's largest primate: 80% loss of Grauer's Gorilla (*Gorilla beringei graueri*) population justifies critically endangered status. *PloS one*, 11(10), e0162697.
284. Plumptre, A. J., Robbins, M. M., & Williamson, E. A. (2019). Gorilla beringei. *The IUCN Red List of Threatened Species* 2019: e.T39994A115576640.
285. Polasky, S., Camm, J. D., & Garber-Yonts, B. (2001). Selecting biological reserves cost-effectively: an application to terrestrial

- vertebrate conservation in Oregon. *Land Economics*, 77(1), 68-78.
286. Polley, H. W., Mayeux, H. S., Johnson, H. B., & Tischler, C. R. (1997). atmospheric CO₂, soil water, and shrub/grass ratios on rangelands. *Rangeland Ecology & Management/Journal of Range Management Archives*, 50(3), 278-284.
 287. Post, E., Peterson, R. O., Stenseth, N. C., & McLaren, B. E. (1999). Ecosystem consequences of wolf behavioural response to climate. *Nature*, 401(6756), 905-907.
 288. Poulter, B., Aragao, L., Heyder, U., Gumpenberger, M., Heinke, J., Langerwisch, F., ... & Cramer, W. (2010). Net biome production of the Amazon Basin in the 21st century. *Global Change Biology*, 16(7), 2062-2075.
 289. Powell, B., Hall, J., & Johns, T. (2011). Forest cover, use and dietary intake in the East Usambara Mountains, Tanzania. *International Forestry Review*, 13(3), 305-317.
 290. Putz, F. E., Dykstra, D. P., & Heinrich, R. (2000). Why poor logging practices persist in the tropics. *Conservation Biology*, 14(4), 951-956.
 291. Putz, F. E., Zuidema, P. A., Synnott, T., Peña-Claros, M., Pinard, M. A., Sheil, D., ... & Zagt, R. (2012). Sustaining conservation values in selectively logged tropical forests: the attained and the attainable. *Conservation Letters*, 5(4), 296-303.
 292. Prado-Martinez, J., Sudmant, P. H., Kidd, J. M., Li, H., Kelley, J. L., Lorente-Galdos, B., ... & Marques-Bonet, T. (2013). Great ape genetic diversity and population history. *Nature*, 499(7459), 471-475.
 293. Priess, J. A., Mimler, M., Klein, A. M., Schwarze, S., Tschamtkte, T., & Steffan-Dewenter, I. (2007). Linking deforestation scenarios to pollination services and economic returns in coffee agroforestry systems. *Ecological Applications*, 17(2), 407-417.
 294. Ramankutty, N., Evan, A. T., Monfreda, C., & Foley, J. A. (2008). Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global biogeochemical cycles*, 22(1).
 295. Randerson, J. T., Van der Werf, G. R., Collatz, G. J., Giglio, L., Still, C. J., Kasibhatla, P., ... & Kasischke, E. S. (2005). Fire emissions from C3 and C4 vegetation and their influence on interannual variability of atmospheric CO₂ and $\delta^{13}\text{C}_{\text{CO}_2}$. *Global Biogeochemical Cycles*, 19(2).
 296. Ratnam, J., Bond, W. J., Fensham, R. J., Hoffmann, W. A., Archibald, S., Lehmann, C. E., ... & Sankaran, M. (2011). When is a 'forest' savanna, and why does it matter?. *Global Ecology and Biogeography*, 20(5), 653-660.
 297. Ravindranath, N. H. (2007). Mitigation and adaptation synergy in forest sector. *Mitigation and Adaptation Strategies for Global Change*, 12(5), 843-853.
 298. Reager, J. T., & Famiglietti, J. S. (2013). Characteristic mega-basin water storage behavior using GRACE. *Water resources research*, 49(6), 3314-3329.
 299. Reynaud-Farrera, I., Maley, J., & Wirrmann, D. (1996). Vegetation and climate in the forests of South-West Cameroon since 4770 years BP: pollen analysis of sediments from Lake Ossa. *CR Acad. Sci. Paris*, 322 (series II a), 749-755.
 300. Ricketts, T. H., Daily, G. C., Ehrlich, P. R., & Michener, C. D. (2004). Economic value of tropical forest to coffee production. *Proceedings of the National Academy of Sciences*, 101(34), 12579-12582.
 301. Ricketts, T. H., Regetz, J., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., Bogdanski, A., ... & Viana, B. F. (2008). Landscape effects on crop pollination services: are there general patterns?. *Ecology Letters*, 11(5), 499-515.
 302. Ribot, J. C., Najam, A., & Watson, G. (1996). Climate variation, vulnerability and sustainable development in the semi-arid tropics. In Ribot, J., Magalhães, A. R., & Panagides, S. (Eds.), *Climate Variability, Climate Change and Social Vulnerability in the Semi-arid Tropics*, 13-54. Cambridge University Press.
 303. Riggio, J., Baillie, J. E., Brumby, S., Ellis, E., Kennedy, C. M., Oakleaf, J. R., ... & Jacobson, A. P. (2020). Global human influence maps reveal clear opportunities in conserving Earth's remaining intact terrestrial ecosystems. *Global Change Biology*, 26(8), 4344-4356.
 304. Riley, L., & Riley, W. (2005). *Nature's strongholds: the world's great wildlife reserves*. Princeton University Press.
 305. Rizkalla, C., Blanco-Silva, F., & Gruver, S. (2007). Modeling the impact of Ebola and bushmeat hunting on Western Lowland Gorillas. *EcoHealth*, 4, 151-155.
 306. Robbins, M. M., & McNeillage, A. (2003). Home range and frugivory patterns of mountain gorillas in Bwindi Impenetrable National Park, Uganda. *International Journal of Primatology*, 24, 467-491.
 307. Robbins, M. M., Gray, M., Fawcett, K. A., Nutter, F. B., Uwingeli, P., Mburanumwe, I., ... & Robbins, A. M. (2011). Extreme conservation leads to recovery of the Virunga mountain gorillas. *PLoS one*, 6(6), e19788.
 308. Robbins, M. M. (1995). A demographic analysis of male life history and social structure of mountain gorillas. *Behaviour*, 132(1-2), 21-47.
 309. Robinson, J., & Bennett, E. L. (2000). Carrying capacity limits to sustainable hunting in tropical forests. In Robinson, J., & Bennett, E. L. (Eds.), *Hunting for sustainability in tropical forests*. Columbia University Press, 13-30.
 310. Root, T. L., Price, J. T., Hall, K. R., Schneider, S. H., Rosenzweig, C., & Pounds, J. A. (2003). Fingerprints of global warming on wild animals and plants. *Nature*, 421(6918), 57-60.
 311. Rowcliffe, J. M., Jansen, P. A., Kays, R., Kranstauber, B., & Carbone, C. (2016). Wildlife speed cameras: measuring animal travel speed and day range using camera traps. *Remote Sensing in Ecology and Conservation*, 2(2), 84-94.
 312. Roy, J., Arandjelovic, M., Bradley, B. J., Guschanski, K., Stephens, C. R., Bucknell, D., ... & Vigilant, L. (2014a). Recent divergences and size decreases of eastern gorilla populations. *Biology Letters*, 10(11), 20140811.
 313. Roy, J., Gray, M., Stoinski, T., Robbins, M. M., & Vigilant, L. (2014b). Fine-scale genetic structure analyses suggest further male than female dispersal in mountain gorillas. *BMC ecology*, 14.
 314. Rupp, S. K. (2003). Interethnic Relations in Southeastern Cameroon: Challenging the "Hunter-Gatherer"-"Farmer" Dichotomy. *African study monographs*, 28, 37-56.
 315. Rupp, S. K. (2011). *Forests of Belonging: identities, ethnicities, and stereotypes in the Congo River Basin*. Seattle, WA: University of Washington Press.
 316. Ruvoilo, M., Pan, D., Zehr, S., Goldberg, T., Disotell, T. R., & Von Dornum, M. (1994). Gene trees and hominoid phylogeny. *Proceedings of the National Academy of Sciences*, 91(19), 8900-8904.
 317. Ruvoilo, M. (1997). Genetic diversity in hominoid primates. *Annual review of anthropology*, 26(1), 515-540.
 318. Saatchi, S., Asefi-Najafabady, S., Malhi, Y., Aragão, L. E., Anderson, L. O., Myneni, R. B., & Nemani, R. (2013). Persistent effects of a severe drought on Amazonian forest canopy. *Proceedings of the National Academy of Sciences*, 110(2), 565-570.
 319. Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T., Salas, W., ... & Morel, A. (2011a). Benchmark map of forest carbon stocks in tropical regions across three continents. *Proceedings of the national academy of sciences*, 108(24), 9899-9904.
 320. Saatchi, S. S., Houghton, R. A., Dos Santos Alvares, R. C., Soares, J. V., & Yu, Y. (2007). Distribution of aboveground live biomass in the Amazon basin. *Global change biology*, 13(4), 816-837.
 321. Saatchi, S., Marlier, M., Chazdon, R. L., Clark, D. B., & Russell, A. E. (2011b). Impact of spatial variability of tropical forest structure on radar estimation of aboveground biomass. *Remote sensing of environment*, 115(11), 2836-2849.
 322. Sage, R. F. (2004). The evolution of C4 photosynthesis. *New phytologist*, 161(2), 341-370.
 323. Salick, J., & Ross, N. (2018). Traditional peoples and climate change. In Infield, E. M. H., Abunnsar, Y., & Ryan R. L. (Eds.), *Planning for Climate Change*, 160-166. Routledge.
 324. Saltonstall, K., Amato, G., & Powell, J. (1998). Mitochondrial DNA variability in Grauer's gorillas of Kahuzi-Biega National Park. *Journal of Heredity*, 89(2), 129-135.
 325. Samanta, A., Ganguly, S., Vermote, E., Nemani, R. R., & Myneni, R. B. (2012). Why is remote sensing of amazon forest greenness so challenging?. *Earth Interactions*, 16(7), 1-14.
 326. Samset, I. (2002). Conflict of interests or interests in conflict? diamonds & war in the DRC. *Review of African Political Economy*, 29(93-94), 463-480.
 327. Sandler, T. (1998). Global and regional public goods: a prognosis for collective action. *Fiscal studies*, 19(3), 221-247.
 328. Sankaran, M., Hanan, N. P., Scholes, R. J., Ratnam, J., Augustine, D. J., Cade, B. S., ... & Zambatis, N. (2005). Determinants of woody cover in African savannas. *Nature*, 438(7069), 846-849.
 329. Sassen, M., & Jum, C. (2007). Assessing local perspectives in a forested landscape of central Cameroon. *Forests, Trees and Livelihoods*, 17(1), 23-42.
 330. Scally, A., Dutheil, J. Y., Hillier, L. W., Jordan, G. E., Goodhead, I., Herrero, J., ... & Durbin, R. (2012). Insights into hominid evolution from the gorilla genome sequence. *Nature*, 483(7388), 169-175.
 331. Saura, S., Bertzy, B., Bastin, L., Battistella, L., Mandrici, A., & Dubois, G. (2018). Protected area connectivity: Shortfalls in global targets and country-level priorities. *Biological conservation*, 219, 53-67.
 332. Scally, A., Yngvadottir, B., Xue, Y., Ayub, Q., Durbin, R., & Tyler-Smith, C. (2013). A genome-wide survey of genetic variation in gorillas using reduced representation sequencing. *PLoS One*, 8(6), e65066.
 333. Schaller, G. E. (1963). *The mountain gorilla: Ecology and behavior*. The University of Chicago Press.
 334. Schefuß, E., Schouten, S., & Schneider, R. R. (2005). Climatic controls on central African hydrology during the past 20,000 years. *Nature*, 437(7061), 1003-1006.

334. Shackley, M. (2007). *Atlas of travel and tourism development*. Routledge.
335. Schloss, C. A., Nuñez, T. A., & Lawler, J. J. (2012). Dispersal will limit ability of mammals to track climate change in the Western Hemisphere. *Proceedings of the national academy of sciences*, 109(22), 8606-8611.
336. Schmitz, O. J., Post, E., Burns, C. E., & Johnston, K. M. (2003). Ecosystem responses to global climate change: moving beyond color mapping. *BioScience*, 53(12), 1199-1205.
337. Shoo, L. P., Storlie, C., Vanderwal, J., Little, J., & Williams, S. E. (2011). Targeted protection and restoration to conserve tropical biodiversity in a warming world. *Global Change Biology*, 17(1), 186-193.
338. Sloat, L. L., Davis, S. J., Gerber, J. S., Moore, F. C., Ray, D. K., West, P. C., & Mueller, N. D. (2020). Climate adaptation by crop migration. *Nature communications*, 11(1), 1243.
339. Smith, J., & Scherr, S. J. (2003). Capturing the value of forest carbon for local livelihoods. *World development*, 31(12), 2143-2160.
340. Smit, B., & Wandel, J. (2006). Adaptation, adaptive capacity and vulnerability. *Global environmental change*, 16(3), 282-292.
341. Somorin, O. A., Brown, H. C. P., Visseren-Hamakers, I. J., Sonwa, D. J., Arts, B., & Nkem, J. (2012). The Congo Basin forests in a changing climate: policy discourses on adaptation and mitigation (REDD+). *Global Environmental Change*, 22(1), 288-298.
342. Somorin, O. A. (2010). Climate impacts, forest-dependent rural livelihoods and adaptation strategies in Africa: A review. *African Journal of Environmental Science and Technology*, 4(13), 903-912.
343. Sonwa, D. J., Walker, S., Nasi, R., & Kanninen, M. (2011). Potential synergies of the main current forestry efforts and climate change mitigation in Central Africa. *Sustainability Science*, 6, 59-67.
344. Sonwa, D. J., Nkem, J. N., Idinoba, M. E., Bele, M. Y., & Jum, C. (2012). Building regional priorities in forests for development and adaptation to climate change in the Congo Basin. *Mitigation and Adaptation Strategies for Global Change*, 17, 441-450.
345. Spillmann, B., van Noordwijk, M. A., Willems, E. P., Mitra Setia, T., Wipfli, U., & van Schaik, C. P. (2015). Validation of an acoustic location system to monitor Bornean orangutan (*Pongo pygmaeus wurmbii*) long calls. *American Journal of Primatology*, 77(7), 767-776.
346. Spillmann, B., van Schaik, C. P., Setia, T. M., & Sadjadi, S. O. (2017). Who shall I say is calling? Validation of a caller recognition procedure in Bornean flanged male orangutan (*Pongo pygmaeus wurmbii*) long calls. *Bioacoustics*, 26(2), 109-120.
347. Städele, V., Arandjelovic, M., Nixon, S., Bergl, R. A., Bradley, B. J., Breuer, T., ... & Vigilant, L. (2022). The complex Y-chromosomal history of gorillas. *American Journal of Primatology*, 84 (3), e23363.
348. Staver, A. C., Archibald, S., & Levin, S. (2011). Tree cover in sub-Saharan Africa: rainfall and fire constrain forest and savanna as alternative stable states. *Ecology*, 92(5), 1063-1072.
349. Strassburg, B. B., Rodrigues, A. S., Gusti, M., Balmford, A., Fritz, S., Obersteiner, M., ... & Brooks, T. M. (2012). Impacts of incentives to reduce emissions from deforestation on global species extinctions. *Nature Climate Change*, 2(5), 350-355.
350. Steffensen, J. P., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Fischer, H., ... & White, J. W. (2008). High-resolution Greenland ice core data show abrupt climate change happens in few years. *Science*, 321(5889), 680-684.
351. Strindberg, S., Maisels, F., Williamson, E. A., Blake, S., Stokes, E. J., Aba'a, R., ... & Wilkie, D. S. (2018). Guns, germs, and trees determine density and distribution of gorillas and chimpanzees in Western Equatorial Africa. *Science advances*, 4(4), eaar2964.
352. Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P. M., (Eds.). (2013). *Climate Change 2013: The Physical Science Basis: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
353. Sunderlin, W. D., Angelsen, A., Belcher, B., Burgers, P., Nasi, R., Santoso, L., & Wunder, S. (2005). Livelihoods, forests, and conservation in developing countries: an overview. *World development*, 33(9), 1383-1402.
354. Synes, N. W., Ponchon, A., Palmer, S. C., Osborne, P. E., Bodedi, G., Travis, J. M., & Watts, K. (2020). Prioritising conservation actions for biodiversity: Lessening the impact from habitat fragmentation and climate change. *Biological Conservation*, 252, 108819.
355. Taylor, D. M. (1990). Late quaternary pollen records from two Ugandan mires: evidence for environmental changes in the Rukiga highlands of southwest Uganda. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 80(3-4), 283-300.
356. Tchindjang, M., Banga, C. R., Nankam, A., & Makak, J. S. (2005, July). Mapping of protected areas evolution in Cameroon from the beginning to 2000: Lesson to learn and perspectives. In *Proceedings of the 22nd international cartographic conference* (pp. 9-16).
357. Thalmann, O., Fischer, A., Lankester, F., Pääbo, S., & Vigilant, L. (2007). The complex evolutionary history of gorillas: insights from genomic data. *Molecular biology and evolution*, 24(1), 146-158.
358. Thalmann, O., Wegmann, D., Spitzner, M., Arandjelovic, M., Guschanski, K., Leuenberger, C., ... & Vigilant, L. (2011). Historical sampling reveals dramatic demographic changes in western gorilla populations. *BMC evolutionary biology*, 11.
359. Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y. C., ... & Williams, S. E. (2004). Extinction risk from climate change. *Nature*, 427(6970), 145-148.
360. Thomas, D. S., & Twyman, C. (2005). Equity and justice in climate change adaptation amongst natural-resource-dependent societies. *Global environmental change*, 15(2), 115-124.
361. Thompson, I., Mackey, B., McNulty, S., & Mosseler, A. (2009). Forest resilience, biodiversity, and climate change. In *A synthesis of the biodiversity/resilience/stability relationship in forest ecosystems*. Secretariat of the Convention on Biological Diversity, Montreal. Technical Series, 43(1), 1-67.
362. Thuiller, W., Lavorel, S., Araújo, M. B., Sykes, M. T., & Prentice, I. C. (2005). Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences*, 102(23), 8245-8250.
363. Thuiller, W., Broennimann, O., Hughes, G., Alkamade, J. R. M., Midgley, G. F., & Corsi, F. (2006). Vulnerability of African mammals to anthropogenic climate change under conservative land transformation assumptions. *Global Change Biology*, 12(3), 424-440.
364. Thuiller, W., Guéguen, M., Renaud, J., Karger, D. N., & Zimmermann, N. E. (2019). Uncertainty in ensembles of global biodiversity scenarios. *Nature Communications*, 10(1), 1446.
365. Tichaawa, T. M. (2017). Business tourism in Africa: The case of Cameroon. *Tourism Review International*, 21(2), 181-192.
366. Tieguhong, J. C., Ndoye, O. (2006). Transforming subsistence products to propellers of sustainable rural development: non-timber forest products (NTFPs) production and trade in Cameroon. In Wohlmuth, K., Burger, P., Gutowski, A., Hussain, M. N., Kneduk, T., Meyn, M (Eds.), *Africa - escaping the primary commodities dilemma*, 107-138. Lit Verlag.
367. Tierney, J. E., Russell, J. M., Huang, Y., Damsté, J. S. S., Hopmans, E. C., & Cohen, A. S. (2008). Northern hemisphere controls on tropical southeast African climate during the past 60,000 years. *science*, 322(5899), 252-255.
368. Tierney, J. E., Russell, J. M., & Huang, Y. (2010). A molecular perspective on Late Quaternary climate and vegetation change in the Lake Tanganyika basin, East Africa. *Quaternary Science Reviews*, 29(5-6), 787-800.
369. Tierney, J. E., Lewis, S. C., Cook, B. I., LeGrande, A. N., & Schmidt, G. A. (2011). Model, proxy and isotopic perspectives on the East African Humid Period. *Earth and Planetary Science Letters*, 307(1-2), 103-112.
370. Tinley, K. L. (1982). The influence of soil moisture balance on ecosystem patterns in southern Africa. In Huntley, J. B., Walker, H. B. (Eds.), *Ecology of tropical savannas*. Springer Berlin Heidelberg, 175-192.
371. Tocheri, M. W., Dommain, R., McFarlin, S. C., Burnett, S. E., Troy Case, D., Orr, C. M., ... & Jungers, W. L. (2016). The evolutionary origin and population history of the grauer gorilla. *American Journal of Physical Anthropology*, 159, 4-18.
372. Tol, R. S., Downing, T. E., Kuik, O. J., & Smith, J. B. (2004). Distributional aspects of climate change impacts. *Global Environmental Change*, 14(3), 259-272.
373. Tompkins, E. L., & Adger, W. N. (2004). Does adaptive management of natural resources enhance resilience to climate change?. *Ecology and society*, 9(2).
374. Travers, T. J., Alison, J., Taylor, S. D., Crick, H. Q., & Hodgson, J. A. (2021). Habitat patches providing south-north connectivity are under-protected in a fragmented landscape. *Proceedings of the Royal Society B*, 288(1957), 20211010.
375. Tuia, D., Kellenberger, B., Beery, S., Costelloe, B. R., Zuffi, S., Risse, B., ... & Berger-Wolf, T. (2022). Perspectives in machine learning for wildlife conservation. *Nature communications*, 13 (1), 792.
376. Uchida, A. (1996). What we don't know about great ape variation. *Trends in ecology & evolution*, 11(4), 163-168.
377. United Nations Framework Convention on Climate Change (UNFCCC). (2015). Adoption of the Paris agreement. Proposal by the President, 282(2).
378. UNEP, C., & IUCN, T. (2013). Elephants in the dust-the African elephant crisis. *A rapid response assessment*. Norway: United Nations Environment Programme, GRID-Arendal. Retrieved from <https://www.unep.org/resources/report/elephants-dust-african-elephant-crisis>
379. United Nations. (2015). Framework Convention on Climate Change. *Adoption of the Paris agreement*. Retrieved from <https://unfccc.int/documents/37107>
380. Valentini, R., Arneth, A., Bombelli, A., Castaldi, S., Cazzolla Gatti, R., Chevallier, F., ... & Scholes, R. J. (2014). A full greenhouse gases budget of Africa: synthesis, uncertainties, and vulnerabilities. *Biogeosciences*, 11(2), 381-407.
381. van den Berg, J., & Biesbrouck, K. (2000). The social dimension of rainforest management in Cameroon: Issues for co-management. *The Tropenbos-Cameroon Programme*.
382. Van Breemen, N. (1993). Soils as biotic constructs favouring net primary productivity. *Geoderma*, 57(3), 183-211.

383. Van der Valk, T., Díez-del-Molino, D., Marques-Bonet, T., Guschanski, K., & Dalén, L. (2019). Historical genomes reveal the genomic consequences of recent population decline in eastern gorillas. *Current Biology*, 29(1), 165-170.
384. Van Vliet, N., Milner-Gulland, E. J., Bousquet, F., Saqalli, M., & Nasi, R. (2010). Effect of small-scale heterogeneity of prey and hunter distributions on the sustainability of bushmeat hunting. *Conservation Biology*, 24(5), 1327-1337.
385. Van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., ... & Rose, S. K. (2011). The representative concentration pathways: an overview. *Climatic change*, 109, 5-31.
386. Vigilant, L., Roy, J., Bradley, B. J., Stoneking, C. J., Robbins, M. M., & Stoinski, T. S. (2015). Reproductive competition and inbreeding avoidance in a primate species with habitual female dispersal. *Behavioral Ecology and Sociobiology*, 69, 1163-1172.
387. Vincens, A., Schwartz, D., Bertaux, J., Elenga, H., & de Namur, C. (1998). Late Holocene climatic changes in western equatorial Africa inferred from pollen from Lake Sinnnda, southern Congo. *Quaternary Research*, 50(1), 34-45.
388. Vincens, A., Schwartz, D., Elenga, H., Reynaud-Farrera, I., Alexandre, A., Bertaux, J., ... & Servant, M. (1999). Forest response to climate changes in Atlantic Equatorial Africa during the last 4000 years BP and inheritance on the modern landscapes. *Journal of Biogeography*, 26(4), 879-885.
389. Vincens, A. (1993). New pollen sequence from Lake Tanganyika: 30,000 years of botanical and climatic history of the northern basin. *Review of Palaeobotany and Palynology*, 78 (3-4), 381-394.
390. Vinding, D. (Ed.). (2019). Middle East. In Berger, D. N. (Ed), *The Indigenous World 2019*. International Work Group for Indigenous Affairs (IWGIA), 375-384.
391. Walker, B., Gunderson, L., Kinzig, A., Folke, C., Carpenter, S., & Schultz, L. (2006). A handful of heuristics and some propositions for understanding resilience in social-ecological systems. *Ecology and society*, 11(1).
392. Walsh, P. D., Abernethy, K. A., Bermejo, M., Beyers, R., De Wachter, P., Akou, M. E., ... & Wilkie, D. S. (2003). Catastrophic ape decline in western equatorial Africa. *Nature*, 422(6932), 611-614.
393. Walsh, P. D. (2006). Ebola and Commercial Hunting: Dim Prospects for African Apes. In Laurance, W. F., & Peres, C. A. (Eds.), *Emerging Threats to Tropical Forests*. IL: The University of Chicago Press, 175-198.
394. Walter, A., & Segerstedt, A. (2012). International trade of biofuels: current trends and the potential role of Africa. *Bioenergy for sustainable development in Africa*, 147-162.
395. Walther, G. R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J., ... & Bairlein, F. (2002). Ecological responses to recent climate change. *Nature*, 416(6879), 389-395.
396. Wan, M., Colfer, C. J., & Powell, B. (2011). Forests, women and health: opportunities and challenges for conservation. *International Forestry Review*, 13(3), 369-387.
397. Ward, M., Saura, S., Williams, B., Ramírez-Delgado, J. P., Arafeh-Dalmau, N., Allan, J. R., ... & Watson, J. E. (2020). Just ten percent of the global terrestrial protected area network is structurally connected via intact land. *Nature communications*, 11(1), 4563.
398. Warman, L., & Moles, A. T. (2009). Alternative stable states in Australia's Wet Tropics: a theoretical framework for the field data and a field-case for the theory. *Landscape Ecology*, 24, 1-13.
399. Washington, R., James, R., Pearce, H., Pokam, W. M., & Moufouma-Okia, W. (2013). Congo Basin rainfall climatology: can we believe the climate models? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1625), 20120296.
400. Watrin, J., Lézine, A. M., & Hély, C. (2009). Plant migration and plant communities at the time of the "green Sahara". *Comptes Rendus. Géoscience*, 341(8-9), 656-670.
401. Watts, D. P. (1996). Comparative socio-ecology of gorillas. In McGrew, W. C., Marchant, L. F., Nishida T. (Eds.), *Great ape societies*. Cambridge University Press, 16-28.
402. White, L. J. T. (2001a). The African rain forest: Climate and vegetation. In Weber, W., White L. J. T., Vedder, A., Naughton-Treves, L. (Eds.), *African Rain Forest Ecology and Conservation*. Yale University Press, 3-29.
403. White L. J. T. (2001b). Forest-savanna dynamics and the origins of Marantaceae forest in central Gabon. In Weber, W., White, L. J., Naughton-Treves, L., & Vedder, A. (Eds.), *African rain forest ecology and conservation: an interdisciplinary perspective*. Yale University Press, 165-182.
404. White F. J. (1985). *The Vegetation of Africa. A Descriptive Memoir to Accompany the UNESCO/AETFAT/UNSO Vegetation Map of Africa*. Unesco, 1983.
405. Wiben, M., Kulesza, J., & Kulesza, P. (Eds.). (2019). Central Africa. In Berger, D. N. (Ed), *The Indigenous World 2019*. International Work Group for Indigenous Affairs (IWGIA), 479-517.
406. Wilkie, D. S., & Laporte, N. (2001). Forest area and deforestation in central Africa: current knowledge and future directions. In Weber, W., White L. J. T., Vedder, A., Naughton-Treves, L. (Eds.), *African Rain Forest Ecology and Conservation*. Yale University Press, 119-139.
407. Wilkie, D. S., Shaw, E., Rotberg, F., Morelli, G., & Auzel, P. (2000). Roads, development, and conservation in the Congo Basin. *Conservation biology*, 14(6), 1614-1622.
408. Wilkie, D. S., Sidle, J. G., Boundzanga, G. C., Auzel, P., & Blake, S. (2001). Defaunation, not deforestation: Commercial logging and market hunting in northern Congo. In Fimbel, R., Grajal, A., & Robinson, J. (Eds), *The cutting edge: conserving wildlife in logged tropical forests*. Columbia University Press, 375-400.
409. Wilkie, D. S., Sidle, J. G., & Boundzanga, G. C. (1992). Mechanized logging, market hunting, and a bank loan in Congo. *Conservation biology*, 6(4), 570-580.
410. Wilkie, D. S., & Carpenter, J. F. (1999). Can nature tourism help finance protected areas in the Congo Basin?. *Oryx*, 33(4), 332-338.
411. Williams, R. J., Carter, J., Duff, G. A., Woinarski, J. Z., Cook, G. D., & Farrer, S. L. (2005). Carbon accounting, land management, science and policy uncertainty in Australian savanna landscapes: introduction and overview. *Australian Journal of Botany*, 53(7), 583-588.
412. Willis, K. J., & McElwain, J. C. (2014). *The evolution of plants*. Oxford University Press.
413. Willis, K. J., Bennett, K. D., Burrough, S. L., Macias-Fauria, M., & Tovar, C. (2013). Determining the response of African biota to climate change: using the past to model the future. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1625), 20120491.
414. Willis, K. J., & Bhagwat, S. A. (2009). Biodiversity and climate change. *Science*, 326(5954), 806-807.
415. Willis, K. J., & MacDonald, G. M. (2011). Long-term ecological records and their relevance to climate change predictions for a warmer world. *Annual Review of Ecology, Evolution, and Systematics*, 42(1), 267-287.
416. Wirmann, D., Bertaux, J., & Kossoni, A. (2001). Late Holocene paleoclimatic changes in Western Central Africa inferred from mineral abundance in dated sediments from Lake Ossa (Southwest Cameroon). *Quaternary Research*, 56(2), 275-287.
417. Witmer, G. W. (2005). Wildlife population monitoring: some practical considerations. *Wildlife Research*, 32(3), 259-263.
418. Woodburn, J. (1997). Indigenous discrimination: The ideological basis for local discrimination against hunter-gatherer minorities in sub-Saharan Africa. *Ethnic and Racial Studies*, 20(2), 345-361.
419. Woodroffe, R., & Ginsberg, J. R. (1998). Edge effects and the extinction of populations inside protected areas. *Science*, 280(5372), 2126-2128.
420. Woodward, F. I. (2008). Forest air conditioning. *Nature*, 454(7203), 422-423.
421. Wooller, M. J., Street-Perrott, F. A., & Agnew, A. D. Q. (2000). Late Quaternary fires and grassland palaeoecology of Mount Kenya, East Africa: evidence from charred grass cuticles in lake sediments. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 164(1-4), 207-230.
422. World Bank. (2004). *Sustaining forests: a development strategy*. The World Bank.
423. Wu, J. (2000). Slippage effects of the conservation reserve program. *American Journal of Agricultural Economics*, 82(4), 979-992.
424. Xu, L., Samanta, A., Costa, M. H., Ganguly, S., Nemani, R. R., & Myneni, R. B. (2011). Widespread decline in greenness of Amazonian vegetation due to the 2010 drought. *Geophysical Research Letters*, 38(7).
425. Xue, Y., Prado-Martinez, J., Sudmant, P. H., Narasimhan, V., Ayub, Q., Szpak, M., ... & Scally, A. (2015). Mountain gorilla genomes reveal the impact of long-term population decline and inbreeding. *Science*, 348(6231), 242-245.
426. Yamagiwa, J., Mwanza, N., Yumoto, T., & Maruhashi T. (1992). Travel distances and food habits of eastern lowland gorillas: a comparative analysis. In Itoigawa, N., Sugiyama, Y., Sackett, G. P., & Thompson, R. K. R. (Eds.), *Topics in Primatology Vol. 2: Behavior, Ecology, and Conservation*. University of Tokyo Press, 267-281.
427. Yamagiwa, J., & Mwanza, N. (1994). Day-journey length and daily diet of solitary male gorillas in lowland and highland habitats. *International Journal of Primatology*, 15, 207-224.
428. Yamagiwa, J., Maruhashi, T., Yumoto, T. & Mwanza, N. (1996). Dietary and ranging overlap in sympatric gorillas and chimpanzees in Kahuzi-Biega National Park, Zaire. In McGrew, W. C., Marchant, L. F., & Nishida, T. (Eds.), *Great Ape Societies*. Cambridge University Press, 82-98.
429. Yamagiwa, J. (2003). Bushmeat poaching and the conservation crisis in Kahuzi-Biega National Park, Democratic Republic of the Congo. *Journal of Sustainable Forestry*, 16(3-4), 111-130.

430. Yu, N., Jensen-Seaman, M. I., Chemnick, L., Ryder, O., & Li, W. H. (2004). Nucleotide diversity in gorillas. *Genetics*, 166(3), 1375-1383.
431. Zelazowski, P., Malhi, Y., Huntingford, C., Sitch, S., & Fisher, J. B. (2011). Changes in the potential distribution of humid tropical forests on a warmer planet. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369(1934), 137-160.
432. Zimmermann, E., & Lerch, C. (1993). The complex acoustic design of an advertisement call in male mouse lemurs (*Microcebus murinus*, Prosimii, Primates) and sources of its variation. *Ethology*, 93(3), 211-224.
433. Zinke, P. J. (1961). The pattern of influence of individual forest trees on soil properties. *Ecology*, 43(1), 130-133.
434. Zhang, X., Scholz, C. A., Hecky, R. E., Wood, D. A., Zal, H. J., & Ebinger, C. J. (2014). Climatic control of the late Quaternary turbidite sedimentology of Lake Kivu, East Africa: Implications for deep mixing and geologic hazards. *Geology*, 42(9), 811-814.
435. Zhang, Y., Wang, Y., & Niu, H. (2017). Spatio-temporal variations in the areas suitable for the cultivation of rice and maize in China under future climate scenarios. *Science of the Total Environment*, 601, 518-531.
436. Zhou, L., Tian, Y., Myneni, R. B., Ciais, P., Saatchi, S., Liu, Y. Y., ... & Hwang, T. (2014). Widespread decline of Congo rainforest greenness in the past decade. *Nature*, 509(7498), 86-90.



Photo 10: Leaf-footed bug (Subfamily Coreinae) in logged primary rainforest along the Mpié river.