

Optional Column E Explanation Form

This form is intended as an aid to completing the Column E explanation. It is not an official form and its use is voluntary. Names, addresses, protocols, veterinary care programs, and the like, are not required as part of an explanation. A Column E explanation must be written so as to be understood by lay persons as well as scientists.

1. Registration Number: 51-F-0016

2. Number 30 of animals used in this study.

3. Species (common name) Guinea Pig of animals used in this study.

4. Explain the procedure producing pain and/or distress.

Herpes simplex virus 1 and 2 (HSV 1 and HSV 2) are pathogens of human oral or genital herpes. Infection of humans by HSV 1 and HSV 2 results in local virus replication at the inoculation site (eye, oral or genital); then it spreads through sensory nerves to the regional ganglia (trigeminal or dorsal root ganglia) where the virus establishes a latent infection. Those viruses in ganglia can reactivate to cause recurrent outbreaks of disease over the course of a person's life. Both viral and host factors are thought to play a role in the maintenance of HSV 1 or HSV 2 in the latent state, and these factors may also influence rates of reactivation.

The mechanism of HSV latent infection and reactivation are poorly understood. Specific viral genes appear to be involved in the establishment, maintenance and reactivation of latency.

The animal model is essential for the study of HSV pathogenesis and latency due to the poorly understood interaction between virus and host that allow latency to occur, and is necessary for the study of vaccines, antiviral drugs or passive protection of antibody before clinical testing. No computer simulation or cell culture model has successfully mimicked the above processes.

This study is designed to evaluate the effects of mutations in different regions of the virus genome on the course of latent infection. The guinea pig is the only model for HSV2 acute infection and spontaneous recurrent disease. Spontaneous recurrent disease must occur in order to assess the effects of the mutations made in the virus genome. Smaller animals such as the mouse can be latently infected with HSV2, but they do not experience spontaneous recurrent disease.

5. Provide scientific justification why pain and/or distress could not be relieved. State methods or means used to determine that pain and/or distress relief would interfere with test results.

There are six possible routes of infection used to study herpes simplex virus pathogenesis in animal models. Virus can be given via the skin, ear, vagina, eye, footpad or systemically. Herpes infection at any of these sites can spread to central nervous system tissue and lead to death of the

The investigator must inform the veterinary staff of the commencement of studies which include animals that will experience pain or distress. When animals are placed on studies in which morbidity is anticipated, the experimental endpoint must be clearly indicated on the cage card and contact information for investigators **MUST** be provided to the veterinary staff. This will allow the veterinary staff to ensure appropriate monitoring of animals which are expected to exhibit more than minimal or transient morbidity prior to the experimental endpoint. In animals that are showing signs of paresis or paralysis, the urinary bladder will be monitored for urinary retention. The urinary bladder will be expressed at least once a day in animals experiencing