# Stationary Strings Theorem V: Arrest of the September 2026 Navier–Stokes Vortex Singularity by Elastic String Tension, $X \to 0$ Axis Regularization and Self-Consistent Saturation Proof

(Analytical Derivation in the $(X, \eta, q)$ Similarity Coordinates of the Millennium Vortex Solution, Cartesian Axis Regularization and High-Precision Radau/DOP853 Numerical Simulation Results)

Fahri Meral

October 15, 2026

**Abstract**

The September 2026 study *Finite Time Blowup for Navier–Stokes* [1] proves that a smooth fluid starting from rest in the three-dimensional incompressible Navier–Stokes equations develops, in finite time ($t \uparrow 1$, $\tau = 1 - t \to 0$), an axisymmetric vortex core reaching infinite velocity ($\|u_\theta^{(0)}\|_{L^\infty} \asymp \tau^{-1/2-h} \to \infty$). The singularity is driven, in similarity coordinates $(X, \eta, q)$, by the radial pressure drop $\Pi_X^{(0)} = F(X, \eta)^2$ produced by centripetal acceleration and by the cylindrical shear stress $\Sigma_{r\theta} = -q^{-1-h} F(X, \eta) a(X, \eta)$ [1, pp. 8, 26, 27].

In this work, the "dimensional shear and singularity-free core" mechanism predicted by the *Stationary Strings Theorem* [2–5] is applied directly to the September 2026 Navier–Stokes inner-core equations, the $\sim 1/X$ coordinate singularity on the $X \to 0$ symmetry axis is removed by a Cartesian regularity factor ($\ell_\theta = \sqrt{X/X_*}\,\tilde{\ell}$), and the resulting coupled differential system is solved in the parameter space $h \in \{0.002, 0.005, 0.008, 0.010\}$ and $\mu_s \in \{0.02, 0.05, 0.10, 0.20\}$ with high-precision solvers (`Radau`, `DOP853`, `brentq`, $10^{-14}$ tolerance):

1. **Definite Singularity Arrest in the Explicit Model:** Under the negative hoop tension $-\frac{T_{\theta\theta}^{\mathrm{sicim}}}{r}$ produced by one-dimensional elastic strings, the effective pressure gradient vanishes at $s_{\mathrm{kritik}} = -\ln \tau_{\mathrm{kritik}} \in [16.1196793715, 58.9321282335]$ ($\Pi_{\mathrm{eff},X} = 0$), and the diverging Navier–Stokes velocity is arrested at a finite upper bound $u_{\max} \in [5.92 \times 10^2, 1.02 \times 10^{12}]$.

2. **Self-Quenched Saturation in the Fully Self-Consistent Model:** When the nonlinear backreaction of the string tension on the effective shear ($F_{\text{eff}}$) is included, the system is found to choke itself (*self-quenching*) and reach a stable asymptotic saturation at $\Pi_{\text{eff}} \approx 0.181 - 0.187 > 0$ ($s = 40$) instead of a sharp zero crossing.

3. $X \to 0$ **Axis Regularization (Theorem 2 and Test T6):** With the transformation $\ell_\theta(X, s) = \sqrt{X/X_*}\,\tilde{\ell}(X, s)$ and the driving $G(X) = F\sqrt{a^2 + \beta_0^2 l^2}$ ($\beta_0 = 0.5$), the $\frac{1}{2X}$ pole in the pressure gradient cancels; it is verified that at $X = 10^{-12}$ the old formula gives $1.68 \times 10^6$ while the regularized formula gives 0.0478 (contrast $\sim 3.5 \times 10^7$), with $\Pi_{\text{reg}}(0) = 0.952236996205853$ on the exact axis and a finite inner-core central pressure integral $\Pi_{\text{eff,reg}}(0, s) = -0.876471225586346$.

4. **Acoustic Event Horizon and Topological Information Invariant:** Under the radial inflow profile $V_0(X, \eta)$, the wave frequency is verified to vanish ($\omega' \to 0$) at the scale $q_{\text{ufuk}}(X, \eta) \in [9.76 \times 10^{-4}, 1.20 \times 10^{-2}]$, and the topological information invariant is verified to be conserved at machine precision, $\max|\mathcal{I}_n - nc_s| \le 7.11 \times 10^{-15}$, for the string modes $n \in \{1, 2, 3, 5\}$.

5. **Cutoff-Free Global Convergence on $[0, \infty)$ and Proposition 4.2 Stress Quenching (Theorem 3 / Tests T7–T8):** In the balance-subtracted ($\tilde{\ell}^2 - \ell_0^2$) global model with balance-preserving relaxation, the far-field tail is shown to decay as $O(X^{-1-h})$ and the infinite $[0, \infty)$ central pressure integral to converge with $\sim 10^{-14}$ `quad` error without any cutoff; moreover, under the Proposition 4.2 transport operators of [1, pp. 26–27], the string backreaction is proven to quench the tangential stress residual by %5.31 in the $\|\mathcal{T}_0\|_{L^2}$ norm and by %14.34 in the $\|\mathcal{T}_0\|_{L^\infty}$ norm.

6. **Full $(X, s)$ FENE-P PDE Solution and Rational Curvature Factor (Theorem 4 / Tests T9–T10):** In the 2D PDE solution including radial advection $W(X, 0)X\partial_X\tilde{\ell}$ and the FENE-P finite-extensibility bound ($b_{\max} = 10^{16}$), core saturation ($\max\tilde{\ell}^2/b_{\max} = 0.9943 < 1$) and finite velocity ($u_{\max} \approx 3.83$–$4.00 \times 10^8$) are shown to persist; under the rational cylindrical curvature factor $\mathcal{K}_{\text{curv}}(X) = \frac{X_0 + X_*}{2X_*(X_0 + X)}$, the PDE far-field wake exponent rises from $q = 0.9098$ to $q = 1.9010 > 1$, so that the global PDE integral ($I_\infty \approx 0.867336$) also converges without cutoff.

7. **Two-Field Dynamic PDE and Skew-Symmetric Lyapunov Power Cancellation (Theorem 5 / Test T11):** Raising the fluid velocity from an algebraic to a fully dynamic field, with deviation advection $W[X\partial_X(F - F_{\text{base}}) + (1 + h)(F - F_{\text{base}})]$ and $F$-proportional skew-symmetric FENE-P coupling, the reactive powers cancel pointwise, $P_{\text{fluid}} + P_{\text{string}} \equiv 0$ (error $\sim 6 \times 10^{-16}$); positivity ($0 < F \le F_{\text{base}}$, bitwise base preservation at $\mu_s = 0$) and convergent $I_\infty \approx 0.903$–$0.951$ with $q \approx 2.00$ are preserved; the core velocity is suppressed to $F(X_*, 40) \approx 0.483$–$0.491 < 0.498$.

8. **$n$-Mode Spectral Transfer in the Co-Moving Lagrange Frame (Test T12):** On a uniform $N_\xi = 128$ label grid (per-parcel ODEs, no upwind diffusion and no convolution kernel), the $n = 1, 2, 3, 4$ quantum modes are verified to transfer into the fluid velocity field with exact 4/4 $k_{\max} = n$ matching and %90.6–%99.2 spectral purity ($|r| = 0.8143$–$0.9453$); the Euler-window $[\,0.004, 0.040\,]$ escape is recorded as the justification of the Lagrange frame.

# 1 Introduction and Problem Statement

In the first four studies of the *Stationary Strings Theorem* [2–5], it was theoretically modeled that matter entering vortex singularities splits into two opposite dimensions, **String (Frequency)** and **Light (Flow)**: while the extrinsic flow dimension stretches to infinity ($\mathcal{L} \to \infty$), the intrinsic oscillation frequency collapses to zero ($\omega_n \to 0$), and the negative-pressure string core forming at the center removes the singularity.

The 166-page September 2026 study *Finite Time Blowup for Navier–Stokes* [1] proves the formation of a velocity singularity $\limsup_{t\uparrow 1} \|u(t)\|_{L^\infty(\mathbb{R}^3)} = \infty$ under finite kinetic energy in the three-dimensional incompressible Navier–Stokes equations ($\partial_t u + (u\cdot\nabla)u - \nu\Delta u + \nabla p = f,\ \nabla\cdot u = 0$) [1, Theorem 1.1, p. 1]. That solution, however, is derived for a pure (pointwise) continuum fluid containing no intrinsic elastic/topological microstructure.

In this work, we add one-dimensional elastic string (Oldroyd-B / Polyakov filament) tension dynamics to the exact inner-core equations of [1], Sections 3.1 (pp. 7–8) and 4.1 (pp. 25–27), deriving both the analytical theorems and the $X \to 0$ axis regularization, and present the verification results (`Test T1-T12`) obtained with the `duran_sicimler_sim` numerical solver package.

# 2 September 2026 Navier–Stokes Inner-Core Equations and Analytical Reference Profiles

With viscosity scaled to $\nu = 1$, time remaining to singularity $\tau = 1 - t \to 0$ and fixed parameter $0 < h < \frac{1}{100}$, the azimuthal velocity growth exponent is defined as $A = \frac{1}{2} + h$ and the axial length exponent as $D = \frac{1}{2} - h$ ($A + D = 1$) [1, p. 7]. In cylindrical coordinates $(r, \theta, z)$, with concentration scale $q = q(z, \tau) > 0$, the dimensionless similarity coordinates $(X, \eta)$ are [1, Equation (3.2), p. 7; p. 25]:

$$\tau = q(1 - \eta^2), \qquad z = q^D \eta, \qquad X = \frac{r^2}{2q}, \qquad L(\eta) = 1 - 2h\eta^2, \qquad d(\eta) = 1 - \eta^2 \quad (1)$$

The leading velocity components and pressure of the vortex core are [1, Equations

(4.3)–(4.4), p. 25]:

$$\begin{aligned} u_\theta^{(0)} &= q^{-A}E(X,\eta), \qquad u_z^{(0)} = q^{-A}U(X,\eta), \\ ru_r^{(0)} &= V_0(X,\eta), \qquad p^{(0)} = q^{-2A}\Pi^{(0)}(X,\eta) \end{aligned} \tag{2}$$

with the on-axis ($X = 0$) Cartesian regularity condition $E(X,\eta) = \sqrt{2X}F(X,\eta)$. Incompressibility and the centripetal-pressure balance (`navier-stokes.pdf` p. 26, **Equation 4.7**) give the fundamental relations [1, p. 26]:

$$\begin{aligned} V_0(X,\eta) &= \frac{X}{L(\eta)}\left(2\eta U - 2D\eta\mathcal{A}_X(U) - d(\eta)\partial_\eta\mathcal{A}_X(U)\right), \\ \Pi_X^{(0)}(X,\eta) &= \frac{E(X,\eta)^2}{2X} = F(X,\eta)^2 \end{aligned} \tag{3}$$

where $\mathcal{A}_X(U)(X,\eta) = \frac{1}{X}\int_0^X U(x,\eta)dx$ is the radial averaging operator. Per **Equations (4.8) and (4.11)** on pp. 26–27 [1, pp. 26–27], the logarithmic derivative $l(X,\eta) = \mathcal{D}_X \log(\sqrt{2X}E)$ and the cylindrical radial shear field are:

$$\begin{aligned} \Sigma_{r\theta}(X,\eta,q) &= \partial_r u_\theta^{(0)} - \frac{u_\theta^{(0)}}{r} = -q^{-1-h}F(X,\eta)\,a(X,\eta), \\ a(X,\eta) &= 2 - 2l(X,\eta) = -2\frac{X\partial_X F}{F} \end{aligned} \tag{4}$$

Our solver uses analytical reference profiles satisfying both the $X \to 0$ on-axis regularity ($E/\sqrt{2X} \to F_0$) and the $X \to \infty$ asymptotic decay ($E \sim c_\infty X^{-1/2-h}$) of [1, p. 8] exactly:

$$\begin{aligned} F(X,\eta) &= \frac{F_0(1-\alpha_\eta\eta^2)}{(1+X/X_0)^{1+h}}, \qquad U(X,\eta) = \frac{U_0\,\eta}{(1+X/X_0)^{1+h}}, \\ a(X,\eta) &= \frac{2(1+h)X}{X_0+X}, \qquad l(X,\eta) = \frac{X_0 - hX}{X_0+X} \end{aligned} \tag{5}$$

For this profile the algebraic identity $l(X,0) = 1 - \frac{1}{2}a(X,0)$ holds exactly (numerical verification error $1.11 \times 10^{-16}$).

# 3 Elastic String Hoop Tension ($-\frac{T_{\theta\theta}}{r}$), Singularity Arrest and $X \to 0$ Axis Regularization

## 3.1 String Evolution and Arrest on the Reference Core Ring ($X = X_*$, Theorem 1)

Defining the logarithmic singularity time $s = -\ln q = -\ln \tau \in [0, \infty)$ ($q = e^{-s}$), the azimuthal elongation component $\ell(s) \equiv \ell_\theta(X_*, s)$ of the one-dimensional elastic string on the vortex core ring $X_* > 0$ evolves under the $\Sigma_{r\theta}$ shear of Equation (4) and the $\gamma_{\rm rel}$ relaxation term as:

$$\frac{d\ell(s)}{ds} = \kappa_r \, F_{\rm eff}(X_*, s) |a(X_*, 0)| \, e^{hs} \, \ell(s) - \gamma_{\rm rel} \, e^{-s} \, \ell(s), \qquad \ell(0) = \ell_0 \tag{6}$$

The negative hoop tension $-\frac{T^{\rm sicim}_{\theta\theta}}{r}$ produced by the stretched string ring in cylindrical coordinates ($T^{\rm sicim}_{\theta\theta} = \mu_s \ell(s)^2$) modifies the radial pressure gradient of Equation (3) and the effective velocity profile with feedback as:

$$\begin{aligned} \Pi_{{\rm eff},X}(X_*, s) &= F(X_*, 0)^2 - \frac{\mu_s \, e^{-(1+2h)s} \, \ell(s)^2}{2X_*}, \\ u^{\rm eff}_\theta(X_*, s) &= e^{(1/2+h)s} \sqrt{\max\left(0, \, 2X_* \Pi_{{\rm eff},X}(X_*, s)\right)} \end{aligned} \tag{7}$$

In the explicit model neglecting backreaction ($F_{\rm eff} = F(X_*, 0)$), Equation (6) is solved exactly in closed form:

$$\ell_{\rm expl}(s) = \ell_0 \exp\left( \frac{\kappa_r F(X_*, 0) |a(X_*, 0)|}{h} \left(e^{hs} - 1\right) - \gamma_{\rm rel} \left(1 - e^{-s}\right) \right) \tag{8}$$

While $\ell_{\rm expl}(s)^2$ grows at a double-exponential (stretched-exponential) rate, the factor $e^{-(1+2h)s}$ decays only exponentially; hence for every $\mu_s > 0$ and $h > 0$ there exists a finite $s_{\rm kritik} < \infty$ ($\tau_{\rm kritik} = e^{-s_{\rm kritik}} > 0$) with $\Pi_{{\rm eff},X}(X_*, s_{\rm kritik}) = 0$, and the velocity is arrested at $u_{\max} = \max_{s \in [0, s_{\rm kritik}]} u^{\rm eff}_\theta(X_*, s) < \infty$.

## 3.2 Theorem 2: $X \to 0$ Cartesian Axis Regularization (Axis-Regularization)

If the elongation $\ell(X, s)$ in Equation (7) is kept constant ($\ell_0 > 0$) in the $X \to 0$ limit, the term $-\frac{\mu_s \ell^2}{2X}$ produces a $\sim 1/X$ pole on the $X \to 0$ symmetry axis.

**Theorem 2 (Cartesian Axis Regularization):** *By cylindrical on-axis regularity (just as in the $E(X, \eta) = \sqrt{2X} F(X, \eta)$ [1, Equation (4.4)] structure), the azimuthal string elongation is defined as:*

$$\ell_\theta(X,s) = \sqrt{\frac{X}{X_*}}\,\tilde{\ell}(X,s), \qquad \tilde{\ell}(X,0) = \ell_0 \tag{9}$$

*Inserting Equation (9) into the original feedback term* $\frac{\mu_s e^{-(1+2h)s}\ell_\theta^2}{2X}$ *cancels the* $2X$ *factors identically over the entire* $X \in [0,\infty)$ *domain:*

$$\Pi^{\text{reg}}_{\text{eff},X}(X,s) = F(X,0)^2 - \left(\frac{\mu_s}{2X_*}\right) e^{-(1+2h)s}\,\tilde{\ell}(X,s)^2 \tag{10}$$

*and the equation contains no* $1/X$ *singularity. The reduced string amplitude* $\tilde{\ell}(X,s)$ *is driven by the core strain rate* $G(X)$*, combining both the differential shear* $a(X,0)$ *and the core rotation/compression invariant* $l(X,0) = \frac{X_0 - hX}{X_0 + X}$*:*

$$\begin{aligned} \frac{d\tilde{\ell}(X,s)}{ds} &= \kappa_r\, G(X)\, e^{hs}\,\tilde{\ell}(X,s) - \gamma_{\text{rel}}\, e^{-s}\,\tilde{\ell}(X,s), \\ G(X) &= F(X,0)\sqrt{a(X,0)^2 + \beta_0^2\, l(X,0)^2} \end{aligned} \tag{11}$$

*On the* $X = 0$ *axis,* $a(0,0) = 0$ *and* $l(0,0) = 1$*, so* $G(0) = F_0\beta_0 > 0$ *is exactly finite; the zero-driving problem on the axis* $(F|a|(0) = 0)$ *thus disappears,* $\Pi^{\text{reg}}_{\text{eff},X}(0,s)$ *stays finite on the exact axis, and the cumulative central pressure integral* $\Pi_{\text{eff,reg}}(0,s) = -\int_0^{X_{\max}} \Pi^{\text{reg}}_{\text{eff},X}(X,s)\,dX$ *over the inner-core domain* $[0, X_{\max}]$ *is absolutely finite.* □

## 3.3 Theorem 3: Balance-Subtracted Global Model and Cutoff-Free Convergence on $[0,\infty)$ (Test T7)

In Oldroyd-B and Polyakov string mechanics the base equilibrium length $\ell_0$ produces no net force; the net elastic hoop tension acting on the fluid is $\Delta T_{\theta\theta} = \mu_s(\ell_\theta^2(X,s) - \ell_\theta^2(X,0))$, i.e. the difference $(\tilde{\ell}(X,s)^2 - \ell_0^2)$ in the reduced amplitude. Writing the relaxation term in the **balance-preserving** form $-\gamma_{\text{rel}} e^{-s}(\tilde{\ell} - \ell_0)$ gives:

$$\frac{d\tilde{\ell}(X,s)}{ds} = \kappa_r\, G(X)\, e^{hs}\,\tilde{\ell}(X,s) - \gamma_{\text{rel}}\, e^{-s}\left(\tilde{\ell}(X,s) - \ell_0\right), \qquad \tilde{\ell}(X,0) = \ell_0 \tag{12}$$

$$\Pi^{\text{global}}_{\text{eff},X}(X,s) = F(X,0)^2 - \left(\frac{\mu_s}{2X_*}\right) e^{-(1+2h)s}\left(\tilde{\ell}(X,s)^2 - \ell_0^2\right) \tag{13}$$

**Theorem 3:** *Since in the far field* $(X \to \infty)$ $G(X) \sim C_G X^{-1-h}$ $(C_G = F_0\sqrt{4(1+h)^2 + \beta_0^2 h^2})$ *vanishes, and since balance-preserving relaxation preserves the* $\tilde{\ell} = \ell_0$ *equilibrium in undriven regions,* $(\tilde{\ell}(X,s)^2 - \ell_0^2) \sim 2\ell_0^2\kappa_r K(s) G(X) \sim O(X^{-1-h})$ *decays. Hence* $\Pi^{\text{global}}_{\text{eff},X}(X,s)$ *is both pole-free on the* $X = 0$ *axis and absolutely integrable over the infinite* $[0,\infty)$ *interval with no cutoff (*$\left|\int_0^\infty \Pi^{\text{global}}_{\text{eff},X}(X,s)dX\right| < \infty$*).* □

## 3.4 Proposition 4.2 Transport Operators and Tangential Stress Residual $\mathcal{T}_{0,r\theta}$ (Test T8)

Per `navier-stokes.pdf` p. 8 (Proposition B.2) and pp. 26–27 (**Proposition 4.2, Equations 4.8–4.11**), the general axial velocity profile with nonzero even component on the axis ($U_{\text{even}} = 0.5$) and the transport operators are:

$$U_{\text{full}}(X,\eta) = \frac{U_{\text{even}}(1-\alpha_\eta\eta^2) + U_0\,\eta}{(1+X/X_0)^{1+h}} \tag{14}$$

$$\begin{cases} W(X,\eta) = 1 - 2D\eta\mathcal{A}_X(U_{\text{full}}) - d(\eta)\partial_\eta\mathcal{A}_X(U_{\text{full}}) \\ H_c(X,\eta) = D\eta + d(\eta)U_{\text{full}}(X,\eta) \end{cases} \tag{15}$$

The inviscid angular momentum source $S_q(X,\eta)$, the on-axis-regular ($C_Q = 0$) radial integral $Q_s(X,\eta)$ and the tangential stress residual $\mathcal{T}_{0,r\theta}(X,\eta)$ [1, Equations (4.9)–(4.11), pp. 26–27]:

$$\begin{aligned} S_q(X,\eta) &= -Wl - h(1-2\eta U_{\text{full}}) - H_c\frac{\partial_\eta F}{F}, \\ Q_s(X,\eta) &= \frac{1}{F(X,\eta)}\int_0^1 \sigma F(\sigma X,\eta)S_q(\sigma X,\eta)\,d\sigma \end{aligned} \tag{16}$$

$$\mathcal{T}_{0,r\theta}(X,\eta) = F(X,\eta)\left(\frac{XQ_s(X,\eta)}{L(\eta)} - a(X,\eta)\right), \qquad Q_s(0,\eta) = \frac{S_q(0,\eta)}{2} \tag{17}$$

take the above form. On the midplane ($\eta = 0, X = 0$), $W(0,0) = 0, l(0,0) = 1, \partial_\eta F(0,0) = 0$, so the analytic identities $S_q(0,0) = -h$ and $Q_s(0,0) = -h/2$ hold exactly.

## 3.5 Full $(X,s)$ Advective PDE + FENE-P Model (Test T9)

Adding the radial advection operator $+W(X,0)X\partial_X\tilde{\ell}$ of Equation (4.14) on p. 27 of `navier-stokes.pdf` and the FENE-P finite-extensibility bound ($b_{\max} = \ell_{\max}^2$ ($10^{16}$)), the full partial differential equation (PDE) system in $(X,s)$ space reads:

$$\begin{aligned} \frac{\partial\tilde{\ell}(X,s)}{\partial s} + W(X,0)X\frac{\partial\tilde{\ell}(X,s)}{\partial X} &= \kappa_r\,G_{\text{eff}}(X,s)\,e^{hs}\,\tilde{\ell}(X,s)\left(1 - \frac{\tilde{\ell}(X,s)^2}{b_{\max}}\right) \\ &\quad - \gamma_{\text{rel}}\,e^{-s}\left(\tilde{\ell}(X,s) - \ell_0\right) \end{aligned} \tag{18}$$

$$\Pi^{\mathrm{PDE}}_{\mathrm{eff},X}(X,s) = F(X,0)^2 - \left(\frac{\mu_s}{2X_*}\right) e^{-(1+2h)s} \frac{\tilde{\ell}(X,s)^2 - \ell_0^2}{1 - \tilde{\ell}(X,s)^2/b_{\max}},$$
$$F_{\mathrm{eff}}(X,s) = \sqrt{\max\left(0,\, \Pi^{\mathrm{PDE}}_{\mathrm{eff},X}(X,s)\right)} \tag{19}$$

## 3.6 Theorem 4: Rational Cylindrical Curvature Factor and Global PDE Convergence (Test T10)

In the advective 2D PDE model, positive radial advection ($dX/ds = W(X,0)X > 0$) carries the FENE-saturated core strain to the outer region ($X \to \infty$), forming a slowly decaying wake with exponent $q_{\mathrm{wake}} \approx 0.9098 < 1$. Under the constant $\frac{1}{2X_*}$ coefficient this wake diverges the $[0,\infty)$ integral, because the $\ell_\theta = \sqrt{X/X_*}\,\tilde{\ell}$ transformation removes the physical $\frac{1}{2X}$ cylindrical curvature damping at the $X \to \infty$ boundary.

**Theorem 4 (Rational Curvature Factor):** *When the tangential string elongation is defined with a Padé-type rational factor preserving both Cartesian regularity on the $X \to 0$ axis ($\ell_\theta \sim \sqrt{X}$) and physical finite extensibility in the $X \to \infty$ far field ($\ell_\theta \sim \tilde{\ell}$):*

$$\ell_\theta(X,s) = \sqrt{\frac{(X_0 + X_*)X}{X_*(X_0 + X)}}\,\tilde{\ell}(X,s),$$
$$\mathcal{K}_{\mathrm{curv}}(X) = \frac{\ell_\theta(X,s)^2}{2X\,\tilde{\ell}(X,s)^2} = \frac{X_0 + X_*}{2X_*(X_0 + X)} \tag{20}$$

*the effective PDE pressure gradient takes the following fully global form:*

$$\Pi^{\mathrm{PDE,curv}}_{\mathrm{eff},X}(X,s) = F(X,0)^2 - \mathcal{K}_{\mathrm{curv}}(X)\,\mu_s\, e^{-(1+2h)s} \frac{\tilde{\ell}(X,s)^2 - \ell_0^2}{1 - \tilde{\ell}(X,s)^2/b_{\max}} \tag{21}$$

*This factor satisfies $\mathcal{K}_{\mathrm{curv}}(0) = \frac{X_0+X_*}{2X_*X_0} = 1.0$ (pole-free on axis), $\mathcal{K}_{\mathrm{curv}}(X_*) = \frac{1}{2X_*} = 0.5$ (exact match on the reference ring) and $X \cdot \mathcal{K}_{\mathrm{curv}}(X) \to 1.0$ as $X \to \infty$ ($\mathcal{K}_{\mathrm{curv}} \sim X^{-1}$). The PDE far-field wake exponent thus rises from $q_{\mathrm{wake}} \approx 0.91$ to $q = 1 + q_{\mathrm{wake}} \approx 1.9010 > 1$, and the 2D PDE global pressure integral $I_\infty = \int_0^\infty \Pi^{\mathrm{PDE,curv}}_{\mathrm{eff},X}(X,s)dX$ converges absolutely.* □

## 3.7 Theorem 5: Two-Field Dynamic PDE, Cylindrical Dilution and Skew-Symmetric Lyapunov Power Cancellation (Test T11)

In the previous stages the fluid velocity was computed algebraically ($\sqrt{\Pi_{\mathrm{eff}}}$); it is now raised to a fully dynamic field: $F(X,s)$ and $\tilde{\ell}(X,s)$ evolve as a coupled $2N_X$ system on

an $N_X = 80$ logarithmic grid ($X \in [0, 300]$, Method of Lines + `Radau`, $s \in [0, 40]$). Since the self-advection of the base flow is cancelled by the background balance, $F$-advection acts only on the deviation $\delta F = F - F_{\text{base}}$; cylindrical similarity carries $X^{1+h}\delta F$ together, hence the dilution term $+(1+h)W(X,0)\,\delta F$. Physical string-fluid drag is proportional to the local velocity ($F \to 0^+$ kills the force):

$$\begin{aligned}
&\frac{\partial F}{\partial s} + W(X,0)\left[X\frac{\partial (F - F_{\text{base}})}{\partial X} + (1+h)(F - F_{\text{base}})\right] \\
&\quad = -\gamma_F e^{-s}(F - F_{\text{base}}) - \kappa_r \mathcal{M}(X,s) F(X,s) \hat{G}(X)(\tilde{\ell}^2 - \ell_0^2)\phi_{\text{FENE}}, \\
&\frac{\partial \tilde{\ell}}{\partial s} + W(X,0) X \frac{\partial \tilde{\ell}}{\partial X} \\
&\quad = +\kappa_r F(X,s)^2 \hat{G}(X) e^{hs} \tilde{\ell} \phi_{\text{FENE}} - \gamma_{\text{rel}} e^{-s}(\tilde{\ell} - \ell_0),
\end{aligned} \tag{22}$$

with $\hat{G}(X) = \sqrt{a(X,0)^2 + \beta_0^2\, l(X,0)^2}$, $\phi_{\text{FENE}} = 1 - \tilde{\ell}^2/b_{\max}$ (unsigned) and $\mathcal{M}(X,s) = \mu_s \mathcal{K}_{\text{curv}}(X) e^{-(1+h)s}$ ($F(X,0) = F_{\text{base}}(X)$, $\tilde{\ell}(X,0) = \ell_0 = 1.0$, $\gamma_F = \gamma_{\text{rel}} = 0.1$, $b_{\max} = 10^{16}$).

**Theorem 5 (Skew-Symmetric Lyapunov Power Cancellation):** *Defining the reactive powers as* $P_{\text{fluid}} = F(\partial_s F)_{\text{react}}$ *and* $P_{\text{string}} = \mathcal{M} e^{-hs}(\tilde{\ell}^2 - \ell_0^2)(\partial_s \tilde{\ell})_{\text{react}}/\tilde{\ell}$, *all factors* ($\phi_{\text{FENE}}$, $\mathcal{M}$, $e^{\pm hs}$) *cancel pairwise and*

$$P_{\text{fluid}}(X,s) + P_{\text{string}}(X,s) \equiv 0 \tag{23}$$

*holds over the entire* $(X,s)$ *grid at machine precision (*$\max|P_{\text{fluid}} + P_{\text{string}}|/\max|P_{\text{fluid}}| \sim 6 \times 10^{-16}$*). Moreover* $F(X,s) \equiv F_{\text{base}}(X)$ *is preserved bitwise at* $\mu_s = 0$*; for* $\mu_s > 0$, $0 < F(X,s) \le F_{\text{base}}(X)$ *and* $I_\infty$ *converges with* $q \approx 2.00 > 1.9$. □

*Scaling note:* applying the literal $m = 2h$ (undecayed) weight drives $F$ negative at $s \approx 4$ ($F(X_*, s)$: $+0.32 \to -0.37$) and collapses the string ($\min \tilde{\ell} \sim 10^{-8}$); hence the consistent variant $m = 1 + h$ is used. The cancellation algebra is $m$-independent, so Eq. (23) is exact under either scaling.

# 4 Acoustic Event Horizon ($\omega' \to 0$) and Topological Information Invariant ($\mathcal{I}_n = nc_s$)

The local frequency of a wave propagating at speed $c_s$ inside the vortex is Doppler-shifted by the radial inflow velocity $u_r^{(0)}(X,\eta,q) = \frac{V_0(X,\eta)}{\sqrt{2Xq}}$ of Equations (2) and (3):

$$\frac{\omega'(X,\eta,q)}{\omega_0} = 1 - \frac{|u_r^{(0)}(X,\eta,q)|}{c_s} = 1 - \sqrt{\frac{q_{\text{ufuk}}(X,\eta)}{q}}, \qquad q_{\text{ufuk}}(X,\eta) = \frac{V_0(X,\eta)^2}{2Xc_s^2} \tag{24}$$

At the $q \to q_{\text{ufuk}}(X, \eta)$ limit the wave freezes completely with $\omega' \to 0$ (Acoustic Event Horizon). Meanwhile the $n$-th harmonic string oscillator in logarithmic time ($s = -\ln \tau$, $Q(s) \equiv q_n(\tau)$) satisfies:

$$\frac{d^2Q}{ds^2} + \frac{dQ}{ds} + \omega_{n,0}^2 \, e^{-2(1+h)s} \, Q(s) = 0, \qquad \omega_{n,0} = \frac{nc_s}{\ell_0} \tag{25}$$

Under vortex shear the intrinsic frequency vanishes as $\omega_n(\tau) = \omega_{n,0}\tau^{+h} \to 0$, while the light/flow size stretches as $\mathcal{L}_{\text{isik}}(\tau) = \ell_0 \tau^{-h} \to \infty$, and their product, the topological information invariant $\mathcal{I}_n(\tau) = \omega_n(\tau) \cdot \mathcal{L}_{\text{isik}}(\tau) = nc_s$, stays constant.

# 5 Numerical Simulation Results and Verification Analysis

The theoretical equations are solved with the `SciPy Radau` (stiff ODE solver), `DOP853` (8th-order Runge–Kutta), `quad` (numerical integration) and `brentq` ($10^{-14}$ root finder) algorithms in `solver_engine.py`. Default simulation parameters are $F_0 = 1.0, X_0 = 1.0, \alpha_\eta = 0.3, U_0 = 1.0, \kappa_r = 1.0, \gamma_{\text{rel}} = 0.1, \ell_0 = 1.0, X_* = 1.0, c_s = 5.0, \beta_0 = 0.5$.

## 5.1 Task 1.1 and 1.2 Results: Explicit Arrest and Self-Consistent Saturation

Tables 1–3 report, for the 16 scanned $(h, \mu_s)$ pairs in the explicit model, the critical arrest scale $s_{\text{kritik}}$, the critical time remaining to singularity $\tau_{\text{kritik}}$ and the attained maximum finite velocity $u_{\max}$; Table 4 compares the explicit model with the fully self-consistent backreaction model.

Table 1: Critical arrest scale $s_{\text{kritik}} = -\ln \tau_{\text{kritik}}$ ($X_* = 1.0$, explicit model, `brentq` tolerance $10^{-14}$).

| $\mu_s \backslash h$ | 0.002 | 0.005 | 0.008 | 0.010 |
|---|---|---|---|---|
| 0.02 | 58.9321282335 | 37.3891341406 | 29.6074962230 | 26.5006102235 |
| 0.05 | 50.7999113156 | 32.3644252892 | 25.7034898409 | 23.0434132611 |
| 0.10 | 43.5515385707 | 27.8727534167 | 22.2064958804 | 19.9431484391 |
| 0.20 | 34.6692875415 | 22.3517534700 | 17.8988763400 | 16.1196793715 |

As seen in Tables 1–3, arrest occurs earlier ($s_{\text{kritik}}$ and $u_{\max}$ decrease) as the string elastic modulus $\mu_s$ or the exponent parameter $h$ increases; yet even at the smallest parameters ($\mu_s = 0.02, h = 0.002$) the singularity is definitely arrested at $\tau_{\text{kritik}} = 2.55 \times 10^{-26}$, before reaching $t = 1$.

Table 2: Critical time threshold $\tau_{\text{kritik}} = e^{-s_{\text{kritik}}} > 0$ at which arrest occurs before the singularity.

| $\mu_s \backslash h$ | 0.002 | 0.005 | 0.008 | 0.010 |
|---|---|---|---|---|
| 0.02 | $2.547428 \times 10^{-26}$ | $5.782363 \times 10^{-17}$ | $1.385568 \times 10^{-13}$ | $3.096929 \times 10^{-12}$ |
| 0.05 | $8.667200 \times 10^{-23}$ | $8.796475 \times 10^{-15}$ | $6.872523 \times 10^{-12}$ | $9.825910 \times 10^{-11}$ |
| 0.10 | $1.218448 \times 10^{-19}$ | $7.852664 \times 10^{-13}$ | $2.269036 \times 10^{-10}$ | $2.181728 \times 10^{-9}$ |
| 0.20 | $8.776467 \times 10^{-16}$ | $1.962261 \times 10^{-10}$ | $1.685065 \times 10^{-8}$ | $9.984175 \times 10^{-8}$ |

Table 3: Maximum finite velocity $u_{\max}$ (upper value) and the optimum scale $s_{\text{opt}}$ at which it is attained (below, in parentheses). While $u_\theta^{(0)}$ grows without bound in pure Navier–Stokes, the velocity in the string system is entirely finite.

| $\mu_s \backslash h$ | 0.002 | 0.005 | 0.008 | 0.010 |
|---|---|---|---|---|
| 0.02 | $1.017055\times10^{12}$ (58.00) | $2.842997\times10^{7}$ (36.50) | $6.802640\times10^{5}$ (28.74) | $1.556152\times10^{5}$ (25.65) |
| 0.05 | $1.587953\times10^{10}$ (49.86) | $2.081658\times10^{6}$ (31.46) | $8.672912\times10^{4}$ (24.83) | $2.472853\times10^{4}$ (22.18) |
| 0.10 | $3.850729\times10^{8}$ (42.61) | $1.987847\times10^{5}$ (26.96) | $1.354633\times10^{4}$ (21.32) | $4.696131\times10^{3}$ (19.07) |
| 0.20 | $3.950142\times10^{6}$ (33.72) | $1.084105\times10^{4}$ (21.43) | $1.345939\times10^{3}$ (17.00) | $5.921677\times10^{2}$ (15.24) |

## 5.2 Task 1.2 Results: Self-Quenched Saturation in the Fully Self-Consistent Model

Solving the nonlinear ODE system with the active $F_{\text{eff}}(X_*, s) = \sqrt{\max(0, \Pi_{\text{eff},X})}$ feedback of Equation (6) with `Radau` ($\text{rtol} = 10^{-10}, \text{atol} = 10^{-13}$) reveals a key physical mechanism (Table 4):

- As the string tension grows and lowers $\Pi_{\text{eff},X}$, the effective shear source $F_{\text{eff}}(X_*, s)$ driving the string also decreases. For example, at $\mu_s = 0.20, h = 0.005$ and $s = 30$, the self-consistent string length is $\ell_{\text{self}} = 2.59 \times 10^6$ versus $\ell_{\text{expl}} = 9.90 \times 10^6$ in the explicit model ($\sim 3.8\times$ suppression).

- As a result of this mutual balancing, instead of crashing to zero with a sharp collapse, the pressure gradient in the self-consistent system reaches a **smooth asymptotic saturation (self-quenching saturation)** at $\Pi_{\text{eff}}(40) \in [0.180554, 0.186909] > 0$ for all $\mu_s$ values at $s = 40$ (no zero crossing occurs even in tests extended to $s = 200$; $\Pi_{\text{eff}} > 0$ is preserved). This finding proves that the $s_{\text{kritik}}$ values of the explicit model are a conservative lower bound, and that the fully backreacting system locks itself into a stable core equilibrium.

## 5.3 Task 2 and Theorem 2 (Test T6) Numerical Verification Results

The Acoustic Event Horizon scale $q_{\text{ufuk}}(X, \eta)$ computed with Equation (24) is given in Table 5; the topological information invariant errors obtained with the `DOP853` solutions

Table 4: Comparison of the Explicit model with the Fully Self-Consistent backreaction model ($h = 0.005, X_* = 1.0$). In the self-consistent model, backreaction quenches growth, stabilizing saturation at $\Pi_{\text{eff}} \approx 0.181 - 0.187$.

| $\mu_s$ | $s_{\text{krit}}^{\text{expl}}$ | $u_{\max}^{\text{expl}}$ | $\Pi_{\text{eff}}^{\text{self}}(s=40)$ | $u_\theta^{\text{self}}(s=40)$ | Zero crossing ($s \le 40$) |
|---|---|---|---|---|---|
| 0.02 | 37.3891341406 | $2.8430 \times 10^7$ | 0.186909 | $3.6231 \times 10^8$ | None (asymptotic saturation) |
| 0.05 | 32.3644252892 | $2.0817 \times 10^6$ | 0.182713 | $3.5822 \times 10^8$ | None (asymptotic saturation) |
| 0.10 | 27.8727534167 | $1.9878 \times 10^5$ | 0.181261 | $3.5679 \times 10^8$ | None (asymptotic saturation) |
| 0.20 | 22.3517534700 | $1.0841 \times 10^4$ | 0.180554 | $3.5610 \times 10^8$ | None (asymptotic saturation) |

of Equation (25) and the independent numerical verification tests (`T1-T6`) in Table 6; and the **Test T6** results ($h = 0.005, \mu_s = 0.1, s = 10, \beta_0 = 0.5$) conducted for Theorem 2 ($X \to 0$ Cartesian axis regularization) in detail in Table 7.

Table 5: Acoustic Event Horizon scale $q_{\text{ufuk}}(X, \eta)$ at which the wave frequency vanishes ($\omega' = 0$) in the Navier–Stokes vortex ($h = 0.005, c_s = 5.0$). The midplane ($\eta = 0$) gives the earliest horizon formation.

| $X \backslash \eta$ | $\eta = 0.0$ (midplane) | $\eta = 0.5$ |
|---|---|---|
| $X = 0.2$ | 0.0033210863 | 0.0009755990 |
| $X = 0.5$ | 0.0065627621 | 0.0022716356 |
| $X = 1.0$ | 0.0095758250 | 0.0039106040 |
| $X = 2.0$ | 0.0120034031 | 0.0058256927 |

## 5.4 Task 3 / Test T7 Results: Cutoff-Free Global Convergence on $[0, \infty)$

In the balance-subtracted global model (where $\tilde{\ell}^2 - \ell_0^2$ is computed directly via $2\ell_0\delta + \delta^2$ to prevent tail cancellation loss), the cutoff-free $[0, \infty)$ central pressure results obtained with the $[0, 50]$ inner integral and the $x = 50u^{-1/h}$ tail transformation are given in the table below. Constancy of the far-tail product $X^{1+h}(\tilde{\ell}^2 - \ell_0^2)$ is verified as $40.8996 \to 40.8287$ (relative difference $1.73 \times 10^{-3}$) at $s = 10$ and $84.4855 \to 84.1665$ ($3.78 \times 10^{-3}$) at $s = 20$; `delta_global` vs. `Radau` cross-check agreement is at the $2.20 \times 10^{-14}$ level.

## 5.5 Task 4 / Test T8 Results: Proposition 4.2 Operators and $\mathcal{T}_0$ Stress Residual Quenching

For $h = 0.005, \eta = 0$, the on-axis value $S_q(0, 0) = -0.005 = -h$ is exact, and $|Q_s(10^{-15}, 0) - S_q(0, 0)/2| = 1.48 \times 10^{-16} < 10^{-14}$ is verified ($W \in [0, 0.7616]$ is monotonically increasing over $X \in [0, 10]$). The reduction of the tangential stress residual $\mathcal{T}_0$ norms under string backreaction at $s = 10, \mu_s = 0.1$ is presented below:

Table 6: Topological information invariant $\mathcal{I}_n(\tau) = \omega_n(\tau) \cdot \mathcal{L}_{\text{isik}}(\tau) = nc_s$ conservation errors and solver verification tests (`solver_engine.run_verification()`).

| Test / Mode ($n$) | Initial / Reference value | Maximum numerical Error / Status |
|---|---|---|
| Mode $n = 1$ ($\max \lvert\mathcal{I}_1 - 1c_s\rvert$) | $\omega_{1,0} = 5.0$, $nc_s = 5.0$ | $8.88 \times 10^{-16}$ |
| Mode $n = 2$ ($\max \lvert\mathcal{I}_2 - 2c_s\rvert$) | $\omega_{2,0} = 10.0$, $nc_s = 10.0$ | $1.78 \times 10^{-15}$ |
| Mode $n = 3$ ($\max \lvert\mathcal{I}_3 - 3c_s\rvert$) | $\omega_{3,0} = 15.0$, $nc_s = 15.0$ | $3.55 \times 10^{-15}$ |
| Mode $n = 5$ ($\max \lvert\mathcal{I}_5 - 5c_s\rvert$) | $\omega_{5,0} = 25.0$, $nc_s = 25.0$ | $7.11 \times 10^{-15}$ |
| **T1:** `Radau` ODE vs. analytic $\ell(s)$ ($s \in [0, 12]$) | $X_* = 1, h = 0.005$ | $1.17 \times 10^{-13}$ (relative) |
| **T2:** Conjugate exponent identity ($A + D = 1$) | $h \in \{0.002, 0.005, 0.008, 0.01\}$ | 0.00 (1.0 exact) |
| **T3:** Pressure derivative $d\Pi^{(0)}/dX = F(X, \eta)^2$ | 9 grid points | $4.16 \times 10^{-11}$ (absolute) |
| **T5:** Axis regularity $E(X, 0)/\sqrt{2X} \to F_0$ | $X = 10^{-8}$, $F_0 = 1.0$ | $1.00 \times 10^{-8}$ (0.9999999900) |
| **T6:** Theorem 2 axis regularization ($X \to 0$) | $h = 0.005, \mu_s = 0.1$, $s = 10$ | Finite (`finite=true`), see Table 7 |
| **T7:** Theorem 3 cutoff-free globality ($[0, \infty)$) | $h = 0.005, \mu_s = 0.1$ | $2.40 \times 10^{-14}$ (quad), `pass=true` |
| **T8:** Proposition 4.2 ($Q_s$, $\mathcal{T}_{0,r\theta}$) | $h = 0.005, \eta = 0$ | $1.48 \times 10^{-16}$ (identity), $L^2$ %5.31 quench |
| **T9:** Full PDE + FENE-P | $h = 0.005, \mu_s = 0.1$ | Grid 2.99%, FENE $< 1$ |
| **T10:** Rational curvature ($\mathcal{K}_{\text{curv}}$) | $h = 0.005, \mu_s = 0.1$ | $q = 1.9010 > 1$, $I_\infty \approx 0.867336$ |
| **T11:** Theorem 5 two-field PDE | $h = 0.005$ | $P_f + P_s \sim 6 \times 10^{-16}$, $q \approx 2.00$, see Table 12 |
| **T12:** Lagrange $n$-mode | $h = 0.005, \mu_s = 0.1$ | $k_{\max} == n$ 4/4, purity $\geq 0.9055$, see Table 13 |

Table 7: Theorem 2 ($X \to 0$ Cartesian axis regularization) **Test T6** numerical verification results ($h = 0.005, \mu_s = 0.1, s = 10, \beta_0 = 0.5$). While the old formula blows up on the axis ($\sim 1.68 \times 10^6$), the regularized formula produces a finite $\sim 0.0478$ correction ($\sim 3.5 \times 10^7$ contrast) and a convergent central pressure.

| Physical / Mathematical Quantity Tested | Numerical Value and Verification Result |
|---|---|
| $l(X, 0) = 1 - a(X, 0)/2$ identity error (6 points) | $1.11 \times 10^{-16}$ |
| Axis driving speed $G(0.0)$ / expected $F_0\beta_0$ | 0.5 / 0.5 (exact) |
| Axis driving continuity $\lvert G(0.0) - G(10^{-12})\rvert$ | $1.01 \times 10^{-12}$ |
| On-axis pressure gradient $\Pi^{\text{reg}}_{\text{eff},X}(0.0, s)$ | 0.952236996205853 |
| Axis pressure continuity $\lvert\Pi_{\text{reg}}(0.0) - \Pi_{\text{reg}}(10^{-12})\rvert$ | $1.03 \times 10^{-12}$ |
| Old string correction term ($X = 10^{-12}$) | 1681670.028826187 ($+\infty$ at $X = 0$) |
| New (Theorem 2) string correction term ($X = 10^{-12}$) | 0.04776300379316238 (contrast $\sim 3.5 \times 10^7$) |
| Central pressure $\Pi_{\text{eff,reg}}(0, s) = -\int_0^{50} \Pi^{\text{reg}}_{\text{eff},X} dX$ | $-0.876471225586346$ (`quad` error $4.7 \times 10^{-12}$) |
| $F$-part + string splitting cross-check error | $3.33 \times 10^{-16}$ |

Table 8: **Test T7:** Theorem 3 cutoff-free global central pressure $\Pi^{\text{global}}_{\text{center}} = -\int_0^\infty \Pi^{\text{global}}_{\text{eff},X} dX$ ($h = 0.005, \mu_s = 0.1$).

| $s$ | $\Pi^{\text{global}}_{\text{center}}$ | `quad` error |
|---|---|---|
| 5 | 0.41435589805519 | $2.40 \times 10^{-14}$ |
| 10 | $-0.8649405028647309$ | $1.02 \times 10^{-14}$ |
| 15 | $-0.8593943070652157$ | $1.14 \times 10^{-14}$ |
| 20 | $-0.7915661350122237$ | $8.47 \times 10^{-15}$ |





Figure 1: **Task 1.1 and 1.2 Numerical Solution Plot** ($h = 0.005, X_* = 1.0$)**:** In the explicit model the string tension zeroes the pressure at $s_{\text{krit}} = 22.35$ for $\mu_s = 0.20$ and at $s_{\text{krit}} = 32.36$ for $\mu_s = 0.05$, arresting the velocity at a finite $u_{\max}$ peak. In the fully backreacting (*self-consistent*) model, the decreasing $F_{\text{eff}}$ self-quenches string growth, carrying the system to a stable saturation at $\Pi_{\text{eff}}(40) \approx 0.181$.

Table 9: **Test T8:** Proposition 4.2 tangential stress residual quenching ($h = 0.005, \eta = 0, s = 10, \mu_s = 0.1$).

| Norm | $\|\mathcal{T}_0^{(0)}\|$ (pure NS) | $\|\mathcal{T}_0^{\text{eff}}\|$ (string) | Reduction |
|---|---|---|---|
| $L^2([0,10])$ | 1.21534414 | 1.15085199 | %5.31 |
| $L^\infty([0,10])$ | 0.54468726 | 0.46655594 | %14.34 |





Figure 2: **Theorem 2 and Test T6 Proof** $(h = 0.005, \mu_s = 0.1, s = 10)$**:** While the old unregularized formula blows up to $-\infty$ with a $1.68 \times 10^6$ correction at $X = 10^{-12}$ (red dashed curve), the Theorem-2-regularized formula converges smoothly ($|\Pi_{\mathrm{reg}}(0) - \Pi_{\mathrm{reg}}(10^{-12})| = 1.03 \times 10^{-12}$) to $\Pi_{\mathrm{reg}}(0) = 0.952236996205853$ on the $X = 0$ symmetry axis (teal curve).

## 5.6 Task 5 and Task 6 / Test T9 and Test T10 Results: Full $(X, s)$ FENE-P PDE and Rational Curvature Convergence

In the 2D FENE-P PDE model ($b_{\max} = 10^{16}, s \in [0, 40]$) solved with an $N_X = 80$ logarithmic grid ($[0, 50]$), upwind discretization and `Radau`, the FENE bound reaches saturation without overflow at $\max(\tilde{\ell}^2)/b_{\max} = 0.99428 < 1$, the axis stays finite with $\Pi(0, 40) = 0.7524362544443486$, and grid convergence ($N_X = 160$ vs. 80) is verified at %2.99.

In the constant-$\frac{1}{2X_*}$-coefficient PDE model, positive advection ($dX/ds = WX > 0$) carries the FENE-saturated core strain outward, leaving a slowly decaying negative pressure wake with exponent $q \approx 0.9098 \leq 1$ ($\Pi(300, 40) < 0$), and the constant-coefficient $I_\infty$ diverges. In contrast, with the rational curvature factor $\mathcal{K}_{\mathrm{curv}}(X) = \frac{X_0+X_*}{2X_*(X_0+X)}$ of **Theorem 4 (Test T10)** activated, the wake exponent jumps to $q = 1.9010418395208726 > 1$, the negative wake is suppressed ($\Pi(300, 40) = +7.68 \times 10^{-6}$), and the global PDE integral converges as $I_\infty \approx 0.8673356418769254$.

With the rational curvature factor $\mathcal{K}_{\mathrm{curv}}(X) = \frac{X_0+X_*}{2X_*(X_0+X)}$ of Equation (20) (`curv=True`) instead of the constant $\frac{1}{2X_*}$ coefficient, the **Test T10** results ($h = 0.005, \mu_s = 0.10, s = 40$) below are obtained. The far-field wake exponent rises from 0.9098 to $q = 1.9010418395208726 > 1$, the negative pressure wake at $X = 300$ is sup-

Table 10: **Test T9:** Advection-free ODE vs. full PDE comparison ($h = 0.005, s = 40, b_{\max} = 10^{16}$). With constant coefficients the PDE $I_\infty$ diverges ($q \approx 0.91 < 1$); the core integral $I_{[0,50]}$ and the ODE $I_\infty$ are finite.

| $\mu_s$ | Model | $\Pi(X_*, 40)$ | $\Pi(0, 40)$ | $I_{[0,50]}$ | $I_\infty$ | $u_{\max}$ | $\max \tilde{\ell}^2 / b_{\max}$ |
|---|---|---|---|---|---|---|---|
| 0.05 | ODE | 0.161422 | 0.775325 | 0.869612 | 0.888277 | $3.85 \times 10^8$ | 0.9968 |
| 0.05 | PDE | 0.213457 | 0.775325 | 0.754186 | divergent† | $4.00 \times 10^8$ | 0.9968 |
| 0.10 | ODE | 0.160873 | 0.752436 | 0.862006 | 0.880672 | $3.83 \times 10^8$ | 0.9943 |
| 0.10 | PDE | 0.201045 | 0.752436 | 0.673947 | divergent† | $3.91 \times 10^8$ | 0.9943 |
| 0.20 | ODE | 0.160610 | 0.739626 | 0.856007 | 0.874672 | $3.82 \times 10^8$ | 0.9892 |
| 0.20 | PDE | 0.188022 | 0.739626 | 0.603802 | divergent† | $3.83 \times 10^8$ | 0.9892 |

† In the constant-coefficient PDE model $q \approx 0.91 < 1$, so the $[0, \infty)$ tail integral diverges.

pressed into positive sign ($\Pi(300, 40) = +7.68 \times 10^{-6}$), and the $[0, \infty)$ global PDE integral converges as $I_\infty \approx 0.8673356418769254$. Moreover, since $\mathcal{K}_{\text{curv}}(0) = 1.0 = 2 \times \frac{1}{2X_*}$, the axis dynamics of the $\mu_s = 0.10$ curved model ($\Pi(0, 40) = 0.739626$, $\max \tilde{\ell}^2 / b_{\max} = 0.9892$) matches the constant-coefficient $\mu_s = 0.20$ axis dynamics to 6 significant digits, providing independent cross-verification.

Table 11: **Test T10:** 2D PDE results under the Theorem 4 rational curvature factor $\mathcal{K}_{\text{curv}}(X)$ ($h = 0.005, \mu_s = 0.1, s = 40$). The wake exponent goes $0.91 \to 1.90 > 1$ and the PDE $I_\infty$ integral converges.

| Quantity | Numerical Value and Verification |
|---|---|
| $\mathcal{K}_{\text{curv}}(0)$ / $\mathcal{K}_{\text{curv}}(X_*)$ | 1.0 (exact) / $0.5 = 1/(2X_*)$ (exact) |
| On-axis pressure | 0.739625806970796 / 0.98918 |
| $\Pi_{\text{eff},X}^{\text{PDE,curv}}(0, 40)$ / $\max \tilde{\ell}^2 / b_{\max}$ | (identical to $\mu_s = 0.20$ constant axis) |
| Extended-domain ($[50, 300]$) PDE wake exponent $q$ | **1.9010418395208726 > 1** (constant model: 0.9098) |
| Far-field pressure $\Pi_{\text{eff},X}^{\text{PDE,curv}}(300, 40)$ | $+7.68 \times 10^{-6}$ (negative wake suppressed) |
| Cumulative PDE integral $I_{[0,300]}$ / tail / $\mathbf{I}_\infty$ | 0.865136911748 / −0.000908164080 / **0.867335641877** (finite) |

## 5.7 Test T11 Results: Two-Field Dynamic PDE and Skew-Symmetric Lyapunov Cancellation

Equation (22) is solved on an $N_X = 80$ logarithmic grid ($X \in [0, 300]$, $2N_X$ coupled system, `Radau`, $s \in [0, 40]$, $b_{\max} = 10^{16}$). The reactive powers satisfy $P_{\text{fluid}} + P_{\text{string}} \equiv 0$ over the whole $(X, s)$ grid ($\max |P_f + P_s| / \max |P_f| \sim 6 \times 10^{-16}$); at $\mu_s = 0$, $F \equiv F_{\text{base}}$ is preserved bitwise; for all $\mu_s > 0$, $F \leq F_{\text{base}}$ holds everywhere (violation 0.00) with $\min F > 0$. The far-field exponent is $q \approx 1.99$–$2.00 > 1.9$, so $I_\infty$ converges; $u_{\max}$ sits in the core ($X \approx 0.95$) below the pure-NS bound ($4.17 \times 10^8$).

Table 12: **Test T11:** Two-field dynamic PDE results ($h = 0.005, s = 40$). Core suppression + positivity + convergent $I_\infty$.

| $\mu_s$ | $F(X_*, 40)$ | $F(0, 40)$ | $I_\infty$ | $u_{\max}$ |
|---|---|---|---|---|
| 0.05 | 0.49138327866878023 | 0.8420958862989822 | 0.9509260063237199 | $4.11 \times 10^8$ |
| 0.10 | 0.4876170862575625 | 0.7902275133121127 | 0.9302277288388315 | $4.07 \times 10^8$ |
| 0.20 | 0.4825439357138414 | 0.7188563439046025 | 0.902719968684912 | $4.03 \times 10^8$ |

### 5.8 Test T12 Results: $n$-Mode Transfer in the Co-Moving Lagrange Frame

Since the Euler-window material of $[0, 10]$ is measured to collapse into $\xi_0 \in [0.004, 0.040]$ at $s = 25$ (96% escape), the spectral measurement is performed in the co-moving Lagrange label frame ($\xi = \ln(1 + X_0)/\ln(11) \in [0, 1]$, uniform $N_\xi = 128$ grid, per-parcel ODEs with no upwind diffusion and no convolution kernel): symmetric readout $\Delta F_n = \frac{1}{2}(F_n^{+\varepsilon} - F_n^{-\varepsilon})$ ($\varepsilon = 0.02$), $R_n = \Delta F_n / \max(10^{-30}, |\Delta F_{\text{unif}}|)$ (envelope floor $1.17 \times 10^{-5}$; the $10^{-30}$ floor triggers at 0/128 points), Fourier over uniform $\xi$. All four modes match exactly with $k_{\max} == n$; purity 0.9055–0.9917, correlation $|r| = 0.8143$–$0.9453$ (mild decay at high modes from differential-amplification dispersion of the inhomogeneous core; clipping 0/128 and symmetric-vs-one-sided difference $\leq 0.004$ rule out the alternatives).

Table 13: **Test T12:** Symmetric Lagrange spectrum ($s = 25, h = 0.005, \mu_s = 0.1$). $k_{\max} == n$ matches 4/4.

| $n$ | $k_{\max}$ | purity $\lvert c_n\rvert/\lVert c\rVert$ | $\lvert r\rvert$ |
|---|---|---|---|
| 1 | 1 | 0.9917 | 0.9453 |
| 2 | 2 | 0.9688 | 0.9364 |
| 3 | 3 | 0.9372 | 0.8685 |
| 4 | 4 | 0.9055 | 0.8143 |

## 6 Discussion, Model Limitations and Reproducibility

1. **Explicit Arrest vs. Self-Consistent Saturation:** The explicit model proves arrest with finite $s_{\text{kritik}}$ and $u_{\max}$; the self-consistent model self-quenches growth via backreaction, turning the finite-$s$ zero into asymptotic saturation (no zero crossing at $s \leq 40$). The $s_{\text{kritik}}$ values of the explicit model are hence a conservative lower-bound estimate.

2. $X \to 0$ **Axis Regularization and Far-Field Boundary Condition (Honesty Record):** The $X \to 0$ axis singularity is fully removed by Theorem 2: with $\ell_\theta = \sqrt{X/X_*}\,\tilde{\ell}$, $2X$ cancels, and $\Pi^{\text{reg}}_{\text{eff},X}$ and $G(X)$ are finite on the exact axis ($X = 0.0$ and $X = 10^{-12}$) (Table 7). Since the reduced string term tends to a constant in the far field ($X \to \infty$, $\tilde{\ell} \to$ constant), the pure $\int_0^\infty$ integral contains a divergent far-field shift; the cumulative central pressure integral is therefore reported over the





Figure 3: **Full PDE + FENE-P solution** ($h = 0.005, s \in [0, 40]$)**:** $\tilde{\ell}(X, s)$ profiles, $\Pi(X, 40)$ and $u(X, 40)$ ODE/PDE comparison, $\Pi(X_*, s)$ evolution, cumulative integral $I(s)$ and FENE saturation ($\max \tilde{\ell}^2/b_{\max} < 1$). Figure generated with `generate_figures.py`.





Figure 4: **Two-field PDE + Lyapunov + spectral fingerprint** ($h = 0.005, \mu_s = 0.1$)**:** $F(X, s)$ and $\tilde{\ell}(X, s)$ evolution, Lyapunov exact-zero power ratio ($\sim 10^{-16}$), $H_{\text{total}}(s)$ energy curve with Lagrange $\mathcal{R}_n(\xi)$ traces and purity/correlation bars. Figure generated with `generate_figures.py`.

inner-core domain of influence ($[0, X_{\max} = 50]$, where the $F$-part has converged to %98) as $\Pi_{\text{eff,reg}}(0, s) = -0.876471225586346$. The remaining constant shift belongs to the boundary pressure matching between the inner core and the outer annulus in [1] and does not affect the physics near the axis.

3. **Phase-2 integrity (T7–T10):** Theorem 3 makes the global pressure integral converge without cutoff (T7), Proposition 4.2 stress residual is quenched by string backreaction (T8), the full FENE-P PDE preserves core saturation and finite velocity (T9), and Theorem 4's rational curvature factor also converges the PDE wake (T10, $I_\infty \approx 0.867336$).

4. **Phase-3 integrity (T11–T12):** Theorem 5 verifies skew-symmetric power cancellation in the two-field dynamic PDE ($P_{\text{fluid}} + P_{\text{string}} \equiv 0$, $\sim 6 \times 10^{-16}$), positivity ($0 < F \leq F_{\text{base}}$) and the convergent $I_\infty$ integral ($q \approx 2.00$) (T11); the co-moving Lagrange frame proves spectral transfer with exact 4/4 $k_{\max} = n$ matching (T12, purity %90.6–%99.2). All tests (T1–T12) pass with `solver_engine.py`.

5. **Reproducibility:** All tables and figures (`fig1`–`fig6`) are reproduced exactly with the `duran_sicimler_sim/` directory commands `python solver_engine.py` and `python generate_figures.py`.

# References


[1] OpenAI. (2026, September). *Finite Time Blowup for Navier–Stokes.* 166 pp. (Theorem 1.1 [p. 1], Section 3.1 [pp. 7–8] and Section 4.1 Inner-Core Profile Equations (4.2)–(4.14) [pp. 25–27]). Available at: `https://openai.com/tr-TR/index/navier-stokes-solution/`

[2] Meral, F. (2025a). *Duran Sicimler Teoremi: Zamanın ve Entropinin Yokluğunda Mükemmel Düzenin Yeniden Doğuşu.*

[3] Meral, F. (2025b). *Duran Sicimler Teoremi II: Görelilik, Faz Dönüşümü ve Bilginin Kozmolojik Döngüsü.*

[4] Meral, F. (2026a). *Duran Sicimler Teoremi III: Maddenin İki Boyutlu İkizliği, Girdap Tekilliği ve Bilgiye Faz Ayrışması.*

[5] Meral, F. (2026b). *Duran Sicimler Teoremi IV: Kozmik Fisyon, Girdap Yakıtı ve Tekilliksiz Bilgi Çekirdeği.*

[6] Unruh, W. G. (1981). *Experimental black-hole evaporation?* Physical Review Letters, 46(21), 1351.

[7] Oldroyd, J. G. (1950). *On the formulation of rheological equations of state*. Proceedings of the Royal Society of London A, 200(1063), 523–541.