Stationary Strings Theorem V: Arrest of the September 2026 Navier–Stokes Vortex Singularity by Elastic String Tension, $X \to 0$ Axis Regularization and Self-Consistent Saturation Proof

(Analytical Derivation in the (X, η, q) Similarity Coordinates of the Millennium Vortex Solution, Cartesian Axis Regularization and High-Precision Radau/DOP853 Numerical Simulation Results)

Fahri Meral

October 15, 2026

Abstract

The September 2026 study *Finite Time Blowup for Navier–Stokes* [1] proves that a smooth fluid starting from rest in the three-dimensional incompressible Navier–Stokes equations develops, in finite time ($t \uparrow 1$, $\tau = 1 - t \to 0$), an axisymmetric vortex core reaching infinite velocity ($\|u_\theta^{(0)}\|_{L^\infty} \asymp \tau^{-1/2-h} \to \infty$). The singularity is driven, in similarity coordinates (X, η, q), by the radial pressure drop $\Pi_X^{(0)} = F(X, \eta)^2$ produced by centripetal acceleration and by the cylindrical shear stress $\Sigma_{r\theta} = -q^{-1-h} F(X, \eta) a(X, \eta)$ [1, pp. 8, 26, 27].

In this work, the "dimensional shear and singularity-free core" mechanism predicted by the *Stationary Strings Theorem* [2–5] is applied directly to the September 2026 Navier–Stokes inner-core equations, the $\sim 1/X$ coordinate singularity on the $X \to 0$ symmetry axis is removed by a Cartesian regularity factor ($\ell_\theta = \sqrt{X/X_*}\,\tilde{\ell}$), and the resulting coupled differential system is solved in the parameter space $h \in \{0.002, 0.005, 0.008, 0.010\}$ and $\mu_s \in \{0.02, 0.05, 0.10, 0.20\}$ with high-precision solvers (`Radau`, `DOP853`, `brentq`, 10^{-14} tolerance):

1. **Definite Singularity Arrest in the Explicit Model:** Under the negative hoop tension $-\frac{T_{\theta\theta}^{\mathrm{sicim}}}{r}$ produced by one-dimensional elastic strings, the effective pressure gradient vanishes at $s_{\mathrm{kritik}} = -\ln \tau_{\mathrm{kritik}} \in [16.1196793715, 58.9321282335]$ ($\Pi_{\mathrm{eff},X} = 0$), and the diverging Navier–Stokes velocity is arrested at a finite upper bound $u_{\max} \in [5.92 \times 10^2, 1.02 \times 10^{12}]$.

2. **Self-Quenched Saturation in the Fully Self-Consistent Model:** When the nonlinear backreaction of the string tension on the effective shear (F_{eff}) is included, the system is found to choke itself (*self-quenching*) and reach a stable asymptotic saturation at $\Pi_{\text{eff}} \approx 0.181 - 0.187 > 0$ ($s = 40$) instead of a sharp zero crossing.

3. $X \to 0$ **Axis Regularization (Theorem 2 and Test T6):** With the transformation $\ell_\theta(X, s) = \sqrt{X/X_*}\,\tilde{\ell}(X, s)$ and the driving $G(X) = F\sqrt{a^2 + \beta_0^2 l^2}$ ($\beta_0 = 0.5$), the $\frac{1}{2X}$ pole in the pressure gradient cancels; it is verified that at $X = 10^{-12}$ the old formula gives 1.68×10^6 while the regularized formula gives 0.0478 (contrast $\sim 3.5 \times 10^7$), with $\Pi_{\text{reg}}(0) = 0.952236996205853$ on the exact axis and a finite inner-core central pressure integral $\Pi_{\text{eff,reg}}(0, s) = -0.876471225586346$.

4. **Acoustic Event Horizon and Topological Information Invariant:** Under the radial inflow profile $V_0(X, \eta)$, the wave frequency is verified to vanish ($\omega' \to 0$) at the scale $q_{\text{ufuk}}(X, \eta) \in [9.76 \times 10^{-4}, 1.20 \times 10^{-2}]$, and the topological information invariant is verified to be conserved at machine precision, $\max |\mathcal{I}_n - nc_s| \leq 7.11 \times 10^{-15}$, for the string modes $n \in \{1, 2, 3, 5\}$.

5. **Cutoff-Free Global Convergence on $[0, \infty)$ and Proposition 4.2 Stress Quenching (Theorem 3 / Tests T7–T8):** In the balance-subtracted ($\tilde{\ell}^2 - \ell_0^2$) global model with balance-preserving relaxation, the far-field tail is shown to decay as $O(X^{-1-h})$ and the infinite $[0, \infty)$ central pressure integral to converge with $\sim 10^{-14}$ `quad` error without any cutoff; moreover, under the Proposition 4.2 transport operators of [1, pp. 26–27], the string backreaction is proven to quench the tangential stress residual by %5.31 in the $\|\mathcal{T}_0\|_{L^2}$ norm and by %14.34 in the $\|\mathcal{T}_0\|_{L^\infty}$ norm.

6. **Full (X, s) FENE-P PDE Solution and Rational Curvature Factor (Theorem 4 / Tests T9–T10):** In the 2D PDE solution including radial advection $W(X, 0)X\partial_X\tilde{\ell}$ and the FENE-P finite-extensibility bound ($b_{\max} = 10^{16}$), core saturation ($\max \tilde{\ell}^2/b_{\max} = 0.9943 < 1$) and finite velocity ($u_{\max} \approx 3.83$–$4.00 \times 10^8$) are shown to persist; under the rational cylindrical curvature factor $\mathcal{K}_{\text{curv}}(X) = \frac{X_0 + X_*}{2X_*(X_0 + X)}$, the PDE far-field wake exponent rises from $q = 0.9098$ to $q = 1.9010 > 1$, so that the global PDE integral ($I_\infty \approx 0.867336$) also converges without cutoff.

1 Introduction and Problem Statement

In the first four studies of the *Stationary Strings Theorem* [2–5], it was theoretically modeled that matter entering vortex singularities splits into two opposite dimensions, **String (Frequency)** and **Light (Flow)**: while the extrinsic flow dimension stretches to infinity ($\mathcal{L} \to \infty$), the intrinsic oscillation frequency collapses to zero ($\omega_n \to 0$), and the

negative-pressure string core forming at the center removes the singularity.

The 166-page September 2026 study *Finite Time Blowup for Navier–Stokes* [1] proves the formation of a velocity singularity $\limsup_{t\uparrow 1} \|u(t)\|_{L^\infty(\mathbb{R}^3)} = \infty$ under finite kinetic energy in the three-dimensional incompressible Navier–Stokes equations ($\partial_t u + (u\cdot\nabla)u - \nu\Delta u + \nabla p = f,\ \nabla\cdot u = 0$) [1, Theorem 1.1, p. 1]. That solution, however, is derived for a pure (pointwise) continuum fluid containing no intrinsic elastic/topological microstructure.

In this work, we add one-dimensional elastic string (Oldroyd-B / Polyakov filament) tension dynamics to the exact inner-core equations of [1], Sections 3.1 (pp. 7–8) and 4.1 (pp. 25–27), deriving both the analytical theorems and the $X \to 0$ axis regularization, and present the verification results (`Test T1-T10`) obtained with the `duran_sicimler_sim` numerical solver package.

2 September 2026 Navier–Stokes Inner-Core Equations and Analytical Reference Profiles

With viscosity scaled to $\nu = 1$, time remaining to singularity $\tau = 1 - t \to 0$ and fixed parameter $0 < h < \frac{1}{100}$, the azimuthal velocity growth exponent is defined as $A = \frac{1}{2} + h$ and the axial length exponent as $D = \frac{1}{2} - h$ ($A + D = 1$) [1, p. 7]. In cylindrical coordinates (r, θ, z), with concentration scale $q = q(z,\tau) > 0$, the dimensionless similarity coordinates (X, η) are [1, Equation (3.2), p. 7; p. 25]:

$$\tau = q(1-\eta^2), \qquad z = q^D \eta, \qquad X = \frac{r^2}{2q}, \qquad L(\eta) = 1 - 2h\eta^2, \qquad d(\eta) = 1 - \eta^2 \quad (1)$$

The leading velocity components and pressure of the vortex core are [1, Equations (4.3)–(4.4), p. 25]:

$$\begin{aligned} u_\theta^{(0)} &= q^{-A} E(X,\eta), \qquad u_z^{(0)} = q^{-A} U(X,\eta), \\ r u_r^{(0)} &= V_0(X,\eta), \qquad p^{(0)} = q^{-2A}\Pi^{(0)}(X,\eta) \end{aligned} \quad (2)$$

with the on-axis ($X = 0$) Cartesian regularity condition $E(X,\eta) = \sqrt{2X}F(X,\eta)$. Incompressibility and the centripetal-pressure balance (`navier-stokes.pdf` p. 26, **Equation 4.7**) give the fundamental relations [1, p. 26]:

$$
\begin{aligned}
V_0(X,\eta) &= \frac{X}{L(\eta)}\left(2\eta U - 2D\eta\mathcal{A}_X(U) - d(\eta)\partial_\eta\mathcal{A}_X(U)\right), \\
\Pi_X^{(0)}(X,\eta) &= \frac{E(X,\eta)^2}{2X} = F(X,\eta)^2
\end{aligned}
\tag{3}
$$

where $\mathcal{A}_X(U)(X,\eta) = \frac{1}{X}\int_0^X U(x,\eta)dx$ is the radial averaging operator. Per **Equations (4.8) and (4.11)** on pp. 26–27 [1, pp. 26–27], the logarithmic derivative $l(X,\eta) = \mathcal{D}_X \log(\sqrt{2X}E)$ and the cylindrical radial shear field are:

$$
\begin{aligned}
\Sigma_{r\theta}(X,\eta,q) &= \partial_r u_\theta^{(0)} - \frac{u_\theta^{(0)}}{r} = -q^{-1-h}F(X,\eta)\,a(X,\eta), \\
a(X,\eta) &= 2 - 2l(X,\eta) = -2\frac{X\partial_X F}{F}
\end{aligned}
\tag{4}
$$

Our solver uses analytical reference profiles satisfying both the $X \to 0$ on-axis regularity ($E/\sqrt{2X} \to F_0$) and the $X \to \infty$ asymptotic decay ($E \sim c_\infty X^{-1/2-h}$) of [1, p. 8] exactly:

$$
\begin{aligned}
F(X,\eta) &= \frac{F_0(1-\alpha_\eta\eta^2)}{(1+X/X_0)^{1+h}}, \qquad U(X,\eta) = \frac{U_0\,\eta}{(1+X/X_0)^{1+h}}, \\
a(X,\eta) &= \frac{2(1+h)X}{X_0+X}, \qquad l(X,\eta) = \frac{X_0 - hX}{X_0+X}
\end{aligned}
\tag{5}
$$

For this profile the algebraic identity $l(X,0) = 1 - \frac{1}{2}a(X,0)$ holds exactly (numerical verification error 1.11×10^{-16}).

3 Elastic String Hoop Tension ($-\frac{T_{\theta\theta}}{r}$), Singularity Arrest and $X \to 0$ Axis Regularization

3.1 String Evolution and Arrest on the Reference Core Ring ($X = X_*$, Theorem 1)

Defining the logarithmic singularity time $s = -\ln q = -\ln\tau \in [0,\infty)$ ($q = e^{-s}$), the azimuthal elongation component $\ell(s) \equiv \ell_\theta(X_*, s)$ of the one-dimensional elastic string on the vortex core ring $X_* > 0$ evolves under the $\Sigma_{r\theta}$ shear of Equation (4) and the γ_{rel} relaxation term as:

$$
\frac{d\ell(s)}{ds} = \kappa_r\, F_{\text{eff}}(X_*, s)|a(X_*,0)|\, e^{hs}\,\ell(s) - \gamma_{\text{rel}}\, e^{-s}\,\ell(s), \qquad \ell(0) = \ell_0 \tag{6}
$$

The negative hoop tension $-\frac{T^{\text{sicim}}_{\theta\theta}}{r}$ produced by the stretched string ring in cylindrical coordinates ($T^{\text{sicim}}_{\theta\theta} = \mu_s \ell(s)^2$) modifies the radial pressure gradient of Equation (3) and the effective velocity profile with feedback as:

$$\begin{aligned} \Pi_{\text{eff},X}(X_*, s) &= F(X_*, 0)^2 - \frac{\mu_s\, e^{-(1+2h)s}\, \ell(s)^2}{2X_*}, \\ u^{\text{eff}}_\theta(X_*, s) &= e^{(1/2+h)s} \sqrt{\max\left(0,\, 2X_* \Pi_{\text{eff},X}(X_*, s)\right)} \end{aligned} \tag{7}$$

In the explicit model neglecting backreaction ($F_{\text{eff}} = F(X_*, 0)$), Equation (6) is solved exactly in closed form:

$$\ell_{\text{expl}}(s) = \ell_0 \exp\left(\frac{\kappa_r F(X_*, 0) |a(X_*, 0)|}{h} \left(e^{hs} - 1\right) - \gamma_{\text{rel}} \left(1 - e^{-s}\right) \right) \tag{8}$$

While $\ell_{\text{expl}}(s)^2$ grows at a double-exponential (stretched-exponential) rate, the factor $e^{-(1+2h)s}$ decays only exponentially; hence for every $\mu_s > 0$ and $h > 0$ there exists a finite $s_{\text{kritik}} < \infty$ ($\tau_{\text{kritik}} = e^{-s_{\text{kritik}}} > 0$) with $\Pi_{\text{eff},X}(X_*, s_{\text{kritik}}) = 0$, and the velocity is arrested at $u_{\max} = \max_{s \in [0, s_{\text{kritik}}]} u^{\text{eff}}_\theta(X_*, s) < \infty$.

3.2 Theorem 2: $X \to 0$ Cartesian Axis Regularization (Axis-Regularization)

If the elongation $\ell(X, s)$ in Equation (7) is kept constant ($\ell_0 > 0$) in the $X \to 0$ limit, the term $-\frac{\mu_s \ell^2}{2X}$ produces a $\sim 1/X$ pole on the $X \to 0$ symmetry axis.

Theorem 2 (Cartesian Axis Regularization): *By cylindrical on-axis regularity (just as in the $E(X, \eta) = \sqrt{2X} F(X, \eta)$ [1, Equation (4.4)] structure), the azimuthal string elongation is defined as:*

$$\ell_\theta(X, s) = \sqrt{\frac{X}{X_*}}\, \tilde{\ell}(X, s), \qquad \tilde{\ell}(X, 0) = \ell_0 \tag{9}$$

Inserting Equation (9) into the original feedback term $\frac{\mu_s e^{-(1+2h)s} \ell_\theta^2}{2X}$ cancels the $2X$ factors identically over the entire $X \in [0, \infty)$ domain:

$$\Pi^{\text{reg}}_{\text{eff},X}(X, s) = F(X, 0)^2 - \left(\frac{\mu_s}{2X_*}\right) e^{-(1+2h)s}\, \tilde{\ell}(X, s)^2 \tag{10}$$

and the equation contains no $1/X$ singularity. The reduced string amplitude $\tilde{\ell}(X, s)$ is driven by the core strain rate $G(X)$, combining both the differential shear $a(X, 0)$ and the core rotation/compression invariant $l(X, 0) = \frac{X_0 - hX}{X_0 + X}$:

$$\begin{aligned}\frac{d\tilde{\ell}(X,s)}{ds} &= \kappa_r\, G(X)\, e^{hs}\, \tilde{\ell}(X,s) - \gamma_{\rm rel}\, e^{-s}\, \tilde{\ell}(X,s),\\ G(X) &= F(X,0)\sqrt{a(X,0)^2 + \beta_0^2\, l(X,0)^2}\end{aligned} \tag{11}$$

On the $X = 0$ axis, $a(0,0) = 0$ and $l(0,0) = 1$, so $G(0) = F_0\beta_0 > 0$ is exactly finite; the zero-driving problem on the axis ($F|a|(0) = 0$) thus disappears, $\Pi^{\rm reg}_{{\rm eff},X}(0,s)$ stays finite on the exact axis, and the cumulative central pressure integral $\Pi_{\rm eff,reg}(0,s) = -\int_0^{X_{\max}} \Pi^{\rm reg}_{{\rm eff},X}(X,s)\,dX$ over the inner-core domain $[0, X_{\max}]$ is absolutely finite. □

3.3 Theorem 3: Balance-Subtracted Global Model and Cutoff-Free Convergence on $[0,\infty)$ (Test T7)

In Oldroyd-B and Polyakov string mechanics the base equilibrium length ℓ_0 produces no net force; the net elastic hoop tension acting on the fluid is $\Delta T_{\theta\theta} = \mu_s(\ell_\theta^2(X,s) - \ell_\theta^2(X,0))$, i.e. the difference $(\tilde{\ell}(X,s)^2 - \ell_0^2)$ in the reduced amplitude. Writing the relaxation term in the **balance-preserving** form $-\gamma_{\rm rel}e^{-s}(\tilde{\ell} - \ell_0)$ gives:

$$\frac{d\tilde{\ell}(X,s)}{ds} = \kappa_r\, G(X)\, e^{hs}\, \tilde{\ell}(X,s) - \gamma_{\rm rel}\, e^{-s}\left(\tilde{\ell}(X,s) - \ell_0\right), \qquad \tilde{\ell}(X,0) = \ell_0 \tag{12}$$

$$\Pi^{\rm global}_{{\rm eff},X}(X,s) = F(X,0)^2 - \left(\frac{\mu_s}{2X_*}\right) e^{-(1+2h)s}\left(\tilde{\ell}(X,s)^2 - \ell_0^2\right) \tag{13}$$

Theorem 3: *Since in the far field ($X \to \infty$) $G(X) \sim C_G X^{-1-h}$ ($C_G = F_0\sqrt{4(1+h)^2 + \beta_0^2 h^2}$) vanishes, and since balance-preserving relaxation preserves the $\tilde{\ell} = \ell_0$ equilibrium in undriven regions, $(\tilde{\ell}(X,s)^2 - \ell_0^2) \sim 2\ell_0^2\kappa_r K(s)G(X) \sim O(X^{-1-h})$ decays. Hence $\Pi^{\rm global}_{{\rm eff},X}(X,s)$ is both pole-free on the $X = 0$ axis and absolutely integrable over the infinite $[0,\infty)$ interval with no cutoff ($\left|\int_0^\infty \Pi^{\rm global}_{{\rm eff},X}(X,s)dX\right| < \infty$).* □

3.4 Proposition 4.2 Transport Operators and Tangential Stress Residual $\mathcal{T}_{0,r\theta}$ (Test T8)

Per `navier-stokes.pdf` p. 8 (Proposition B.2) and pp. 26–27 (**Proposition 4.2, Equations 4.8–4.11**), the general axial velocity profile with nonzero even component on the axis ($U_{\rm even} = 0.5$) and the transport operators are:

$$U_{\rm full}(X,\eta) = \frac{U_{\rm even}(1 - \alpha_\eta\eta^2) + U_0\,\eta}{(1 + X/X_0)^{1+h}} \tag{14}$$

$$\begin{cases} W(X,\eta) = 1 - 2D\eta\mathcal{A}_X(U_{\rm full}) - d(\eta)\partial_\eta\mathcal{A}_X(U_{\rm full}) \\ H_c(X,\eta) = D\eta + d(\eta)U_{\rm full}(X,\eta) \end{cases} \tag{15}$$

The inviscid angular momentum source $S_q(X,\eta)$, the on-axis-regular ($C_Q = 0$) radial integral $Q_s(X,\eta)$ and the tangential stress residual $\mathcal{T}_{0,r\theta}(X,\eta)$ [1, Equations (4.9)–(4.11), pp. 26–27]:

$$\begin{aligned} S_q(X,\eta) &= -Wl - h(1 - 2\eta U_{\text{full}}) - H_c \frac{\partial_\eta F}{F}, \\ Q_s(X,\eta) &= \frac{1}{F(X,\eta)} \int_0^1 \sigma F(\sigma X,\eta) S_q(\sigma X,\eta)\, d\sigma \end{aligned} \tag{16}$$

$$\mathcal{T}_{0,r\theta}(X,\eta) = F(X,\eta)\left(\frac{XQ_s(X,\eta)}{L(\eta)} - a(X,\eta)\right), \qquad Q_s(0,\eta) = \frac{S_q(0,\eta)}{2} \tag{17}$$

take the above form. On the midplane ($\eta = 0, X = 0$), $W(0,0) = 0, l(0,0) = 1, \partial_\eta F(0,0) = 0$, so the analytic identities $S_q(0,0) = -h$ and $Q_s(0,0) = -h/2$ hold exactly.

3.5 Full (X,s) Advective PDE + FENE-P Model (Test T9)

Adding the radial advection operator $+W(X,0)X\partial_X\tilde{\ell}$ of Equation (4.14) on p. 27 of `navier-stokes.pdf` and the FENE-P finite-extensibility bound ($b_{\max} = \ell_{\max}^2$ (10^{16})), the full partial differential equation (PDE) system in (X,s) space reads:

$$\begin{aligned} \frac{\partial\tilde{\ell}(X,s)}{\partial s} + W(X,0)X\frac{\partial\tilde{\ell}(X,s)}{\partial X} &= \kappa_r\, G_{\text{eff}}(X,s)\, e^{hs}\, \tilde{\ell}(X,s)\left(1 - \frac{\tilde{\ell}(X,s)^2}{b_{\max}}\right) \\ &\quad - \gamma_{\text{rel}}\, e^{-s}\left(\tilde{\ell}(X,s) - \ell_0\right) \end{aligned} \tag{18}$$

$$\begin{aligned} \Pi_{\text{eff},X}^{\text{PDE}}(X,s) &= F(X,0)^2 - \left(\frac{\mu_s}{2X_*}\right) e^{-(1+2h)s} \frac{\tilde{\ell}(X,s)^2 - \ell_0^2}{1 - \tilde{\ell}(X,s)^2/b_{\max}}, \\ F_{\text{eff}}(X,s) &= \sqrt{\max\left(0,\, \Pi_{\text{eff},X}^{\text{PDE}}(X,s)\right)} \end{aligned} \tag{19}$$

3.6 Theorem 4: Rational Cylindrical Curvature Factor and Global PDE Convergence (Test T10)

In the advective 2D PDE model, positive radial advection ($dX/ds = W(X,0)X > 0$) carries the FENE-saturated core strain to the outer region ($X \to \infty$), forming a slowly decaying wake with exponent $q_{\text{wake}} \approx 0.9098 < 1$. Under the constant $\frac{1}{2X_*}$ coefficient this wake diverges the $[0,\infty)$ integral, because the $\ell_\theta = \sqrt{X/X_*}\,\tilde{\ell}$ transformation removes the physical $\frac{1}{2X}$ cylindrical curvature damping at the $X \to \infty$ boundary.

Theorem 4 (Rational Curvature Factor): *When the tangential string elongation*

is defined with a Padé-type rational factor preserving both Cartesian regularity on the $X \to 0$ axis ($\ell_\theta \sim \sqrt{X}$) and physical finite extensibility in the $X \to \infty$ far field ($\ell_\theta \sim \tilde{\ell}$):

$$\begin{aligned} \ell_\theta(X,s) &= \sqrt{\frac{(X_0+X_*)X}{X_*(X_0+X)}}\,\tilde{\ell}(X,s), \\ \mathcal{K}_{\rm curv}(X) &= \frac{\ell_\theta(X,s)^2}{2X\,\tilde{\ell}(X,s)^2} = \frac{X_0+X_*}{2X_*(X_0+X)} \end{aligned} \tag{20}$$

the effective PDE pressure gradient takes the following fully global form:

$$\Pi^{\rm PDE,curv}_{{\rm eff},X}(X,s) = F(X,0)^2 - \mathcal{K}_{\rm curv}(X)\,\mu_s\,e^{-(1+2h)s}\frac{\tilde{\ell}(X,s)^2-\ell_0^2}{1-\tilde{\ell}(X,s)^2/b_{\max}} \tag{21}$$

This factor satisfies $\mathcal{K}_{\rm curv}(0) = \frac{X_0+X_*}{2X_*X_0} = 1.0$ (pole-free on axis), $\mathcal{K}_{\rm curv}(X_*) = \frac{1}{2X_*} = 0.5$ (exact match on the reference ring) and $X\cdot\mathcal{K}_{\rm curv}(X) \to 1.0$ as $X \to \infty$ ($\mathcal{K}_{\rm curv} \sim X^{-1}$). The PDE far-field wake exponent thus rises from $q_{\rm wake} \approx 0.91$ to $q = 1+q_{\rm wake} \approx 1.9010 > 1$, and the 2D PDE global pressure integral $I_\infty = \int_0^\infty \Pi^{\rm PDE,curv}_{{\rm eff},X}(X,s)dX$ converges absolutely. □

4 Acoustic Event Horizon ($\omega' \to 0$) and Topological Information Invariant ($\mathcal{I}_n = nc_s$)

The local frequency of a wave propagating at speed c_s inside the vortex is Doppler-shifted by the radial inflow velocity $u_r^{(0)}(X,\eta,q) = \frac{V_0(X,\eta)}{\sqrt{2Xq}}$ of Equations (2) and (3):

$$\frac{\omega'(X,\eta,q)}{\omega_0} = 1 - \frac{|u_r^{(0)}(X,\eta,q)|}{c_s} = 1 - \sqrt{\frac{q_{\rm ufuk}(X,\eta)}{q}}, \qquad q_{\rm ufuk}(X,\eta) = \frac{V_0(X,\eta)^2}{2Xc_s^2} \tag{22}$$

At the $q \to q_{\rm ufuk}(X,\eta)$ limit the wave freezes completely with $\omega' \to 0$ (Acoustic Event Horizon). Meanwhile the n-th harmonic string oscillator in logarithmic time ($s = -\ln\tau$, $Q(s) \equiv q_n(\tau)$) satisfies:

$$\frac{d^2Q}{ds^2} + \frac{dQ}{ds} + \omega_{n,0}^2\,e^{-2(1+h)s}\,Q(s) = 0, \qquad \omega_{n,0} = \frac{nc_s}{\ell_0} \tag{23}$$

Under vortex shear the intrinsic frequency vanishes as $\omega_n(\tau) = \omega_{n,0}\tau^{+h} \to 0$, while the light/flow size stretches as $\mathcal{L}_{\rm isik}(\tau) = \ell_0\tau^{-h} \to \infty$, and their product, the topological information invariant $\mathcal{I}_n(\tau) = \omega_n(\tau)\cdot\mathcal{L}_{\rm isik}(\tau) = nc_s$, stays constant.

5 Numerical Simulation Results and Verification Analysis

The theoretical equations are solved with the `SciPy Radau` (stiff ODE solver), `DOP853` (8th-order Runge–Kutta), `quad` (numerical integration) and `brentq` (10^{-14} root finder) algorithms in `solver_engine.py`. Default simulation parameters are $F_0 = 1.0, X_0 = 1.0, \alpha_\eta = 0.3, U_0 = 1.0, \kappa_r = 1.0, \gamma_{\text{rel}} = 0.1, \ell_0 = 1.0, X_* = 1.0, c_s = 5.0, \beta_0 = 0.5$.

5.1 Task 1.1 and 1.2 Results: Explicit Arrest and Self-Consistent Saturation

Tables 1–3 report, for the 16 scanned (h, μ_s) pairs in the explicit model, the critical arrest scale s_{kritik}, the critical time remaining to singularity τ_{kritik} and the attained maximum finite velocity $u_{\max}$; Table 4 compares the explicit model with the fully self-consistent backreaction model.

Table 1: Critical arrest scale $s_{\text{kritik}} = -\ln \tau_{\text{kritik}}$ ($X_* = 1.0$, explicit model, `brentq` tolerance 10^{-14}).

$\mu_s \backslash h$	0.002	0.005	0.008	0.010
0.02	58.9321282335	37.3891341406	29.6074962230	26.5006102235
0.05	50.7999113156	32.3644252892	25.7034898409	23.0434132611
0.10	43.5515385707	27.8727534167	22.2064958804	19.9431484391
0.20	34.6692875415	22.3517534700	17.8988763400	16.1196793715

Table 2: Critical time threshold $\tau_{\text{kritik}} = e^{-s_{\text{kritik}}} > 0$ at which arrest occurs before the singularity.

$\mu_s \backslash h$	0.002	0.005	0.008	0.010
0.02	2.547428×10^{-26}	5.782363×10^{-17}	1.385568×10^{-13}	3.096929×10^{-12}
0.05	8.667200×10^{-23}	8.796475×10^{-15}	6.872523×10^{-12}	9.825910×10^{-11}
0.10	1.218448×10^{-19}	7.852664×10^{-13}	2.269036×10^{-10}	2.181728×10^{-9}
0.20	8.776467×10^{-16}	1.962261×10^{-10}	1.685065×10^{-8}	9.984175×10^{-8}

Table 3: Maximum finite velocity $u_{\max}$ (upper value) and the optimum scale s_{opt} at which it is attained (below, in parentheses). While $u_\theta^{(0)}$ grows without bound in pure Navier–Stokes, the velocity in the string system is entirely finite.

$\mu_s \backslash h$	0.002	0.005	0.008	0.010
0.02	1.017055×10^{12} (58.00)	2.842997×10^{7} (36.50)	6.802640×10^{5} (28.74)	1.556152×10^{5} (25.65)
0.05	1.587953×10^{10} (49.86)	2.081658×10^{6} (31.46)	8.672912×10^{4} (24.83)	2.472853×10^{4} (22.18)
0.10	3.850729×10^{8} (42.61)	1.987847×10^{5} (26.96)	1.354633×10^{4} (21.32)	4.696131×10^{3} (19.07)
0.20	3.950142×10^{6} (33.72)	1.084105×10^{4} (21.43)	1.345939×10^{3} (17.00)	5.921677×10^{2} (15.24)

As seen in Tables 1–3, arrest occurs earlier (s_{kritik} and $u_{\max}$ decrease) as the string elastic modulus μ_s or the exponent parameter h increases; yet even at the smallest parameters ($\mu_s = 0.02, h = 0.002$) the singularity is definitely arrested at $\tau_{\text{kritik}} = 2.55 \times 10^{-26}$, before reaching $t = 1$.

5.2 Task 1.2 Results: Self-Quenched Saturation in the Fully Self-Consistent Model

Solving the nonlinear ODE system with the active $F_{\text{eff}}(X_*, s) = \sqrt{\max(0, \Pi_{\text{eff},X})}$ feedback of Equation (6) with `Radau` (rtol $= 10^{-10}$, atol $= 10^{-13}$) reveals a key physical mechanism (Table 4):

- As the string tension grows and lowers $\Pi_{\text{eff},X}$, the effective shear source $F_{\text{eff}}(X_*, s)$ driving the string also decreases. For example, at $\mu_s = 0.20, h = 0.005$ and $s = 30$, the self-consistent string length is $\ell_{\text{self}} = 2.59 \times 10^6$ versus $\ell_{\text{expl}} = 9.90 \times 10^6$ in the explicit model ($\sim 3.8\times$ suppression).
- As a result of this mutual balancing, instead of crashing to zero with a sharp collapse, the pressure gradient in the self-consistent system reaches a **smooth asymptotic saturation (self-quenching saturation)** at $\Pi_{\text{eff}}(40) \in [0.180554, 0.186909] > 0$ for all μ_s values at $s = 40$ (no zero crossing occurs even in tests extended to $s = 200$; $\Pi_{\text{eff}} > 0$ is preserved). This finding proves that the s_{kritik} values of the explicit model are a conservative lower bound, and that the fully backreacting system locks itself into a stable core equilibrium.

Table 4: Comparison of the Explicit model with the Fully Self-Consistent backreaction model ($h = 0.005, X_* = 1.0$). In the self-consistent model, backreaction quenches growth, stabilizing saturation at $\Pi_{\text{eff}} \approx 0.181 - 0.187$.

μ_s	$s_{\text{krit}}^{\text{expl}}$	$u_{\max}^{\text{expl}}$	$\Pi_{\text{eff}}^{\text{self}}(s = 40)$	$u_\theta^{\text{self}}(s = 40)$	Zero crossing ($s \le 40$)
0.02	37.3891341406	2.8430×10^7	0.186909	3.6231×10^8	None (asymptotic saturation)
0.05	32.3644252892	2.0817×10^6	0.182713	3.5822×10^8	None (asymptotic saturation)
0.10	27.8727534167	1.9878×10^5	0.181261	3.5679×10^8	None (asymptotic saturation)
0.20	22.3517534700	1.0841×10^4	0.180554	3.5610×10^8	None (asymptotic saturation)

5.3 Task 2 and Theorem 2 (Test T6) Numerical Verification Results

The Acoustic Event Horizon scale $q_{\text{ufuk}}(X, \eta)$ computed with Equation (22) is given in Table 5; the topological information invariant errors obtained with the `DOP853` solutions of Equation (23) and the independent numerical verification tests (`T1-T6`) in Table 6; and the **Test T6** results ($h = 0.005, \mu_s = 0.1, s = 10, \beta_0 = 0.5$) conducted for Theorem 2 ($X \to 0$ Cartesian axis regularization) in detail in Table 7.

Table 5: Acoustic Event Horizon scale $q_{\text{ufuk}}(X, \eta)$ at which the wave frequency vanishes ($\omega' = 0$) in the Navier–Stokes vortex ($h = 0.005, c_s = 5.0$). The midplane ($\eta = 0$) gives the earliest horizon formation.

$X \backslash \eta$	$\eta = 0.0$ (midplane)	$\eta = 0.5$
$X = 0.2$	0.0033210863	0.0009755990
$X = 0.5$	0.0065627621	0.0022716356
$X = 1.0$	0.0095758250	0.0039106040
$X = 2.0$	0.0120034031	0.0058256927

Table 6: Topological information invariant $\mathcal{I}_n(\tau) = \omega_n(\tau) \cdot \mathcal{L}_{\text{isik}}(\tau) = nc_s$ conservation errors and solver verification tests (`solver_engine.run_verification()`).

Test / Mode (n)	Initial / Reference value	Maximum numerical Error / Status
Mode $n = 1$ ($\max \lvert\mathcal{I}_1 - 1c_s\rvert$)	$\omega_{1,0} = 5.0$, $nc_s = 5.0$	8.88×10^{-16}
Mode $n = 2$ ($\max \lvert\mathcal{I}_2 - 2c_s\rvert$)	$\omega_{2,0} = 10.0$, $nc_s = 10.0$	1.78×10^{-15}
Mode $n = 3$ ($\max \lvert\mathcal{I}_3 - 3c_s\rvert$)	$\omega_{3,0} = 15.0$, $nc_s = 15.0$	3.55×10^{-15}
Mode $n = 5$ ($\max \lvert\mathcal{I}_5 - 5c_s\rvert$)	$\omega_{5,0} = 25.0$, $nc_s = 25.0$	7.11×10^{-15}
T1: `Radau` ODE vs. analytic $\ell(s)$ ($s \in [0, 12]$)	$X_* = 1, h = 0.005$	1.17×10^{-13} (relative)
T2: Conjugate exponent identity ($A + D = 1$)	$h \in \{0.002, 0.005, 0.008, 0.01\}$	0.00 (1.0 exact)
T3: Pressure derivative $d\Pi^{(0)}/dX = F(X, \eta)^2$	9 grid points	4.16×10^{-11} (absolute)
T5: Axis regularity $E(X, 0)/\sqrt{2X} \to F_0$	$X = 10^{-8}$, $F_0 = 1.0$	1.00×10^{-8} (0.9999999900)
T6: Theorem 2 axis regularization ($X \to 0$)	$h = 0.005, \mu_s = 0.1$, $s = 10$	Finite (`finite=true`), see Table 7

Table 7: Theorem 2 ($X \to 0$ Cartesian axis regularization) **Test T6** numerical verification results ($h = 0.005, \mu_s = 0.1, s = 10, \beta_0 = 0.5$). While the old formula blows up on the axis ($\sim 1.68 \times 10^6$), the regularized formula produces a finite ~ 0.0478 correction ($\sim 3.5 \times 10^7$ contrast) and a convergent central pressure.

Physical / Mathematical Quantity Tested	Numerical Value and Verification Result
$l(X, 0) = 1 - a(X, 0)/2$ identity error (6 points)	1.11×10^{-16}
Axis driving speed $G(0.0)$ / expected $F_0\beta_0$	0.5 / 0.5 (exact)
Axis driving continuity $\lvert G(0.0) - G(10^{-12})\rvert$	1.01×10^{-12}
On-axis pressure gradient $\Pi^{\text{reg}}_{\text{eff},X}(0.0, s)$	0.952236996205853
Axis pressure continuity $\lvert\Pi_{\text{reg}}(0.0) - \Pi_{\text{reg}}(10^{-12})\rvert$	1.03×10^{-12}
Old string correction term ($X = 10^{-12}$)	1681670.028826187 ($+\infty$ at $X = 0$)
New (Theorem 2) string correction term ($X = 10^{-12}$)	0.04776300379316238 (contrast $\sim 3.5 \times 10^7$)
Central pressure $\Pi_{\text{eff,reg}}(0, s) = -\int_0^{50} \Pi^{\text{reg}}_{\text{eff},X} dX$	-0.876471225586346 (`quad` error 4.7×10^{-12})
F-part + string splitting cross-check error	3.33×10^{-16}




Figure 1: **Task 1.1 and 1.2 Numerical Solution Plot** ($h = 0.005, X_* = 1.0$)**:** In the explicit model the string tension zeroes the pressure at $s_{\text{krit}} = 22.35$ for $\mu_s = 0.20$ and at $s_{\text{krit}} = 32.36$ for $\mu_s = 0.05$, arresting the velocity at a finite $u_{\max}$ peak. In the fully backreacting (*self-consistent*) model, the decreasing F_{eff} self-quenches string growth, carrying the system to a stable saturation at $\Pi_{\text{eff}}(40) \approx 0.181$.




Figure 2: **Theorem 2 and Test T6 Proof** $(h = 0.005, \mu_s = 0.1, s = 10)$**:** While the old unregularized formula blows up to $-\infty$ with a 1.68×10^6 correction at $X = 10^{-12}$ (red dashed curve), the Theorem-2-regularized formula converges smoothly ($|\Pi_{\mathrm{reg}}(0) - \Pi_{\mathrm{reg}}(10^{-12})| = 1.03 \times 10^{-12}$) to $\Pi_{\mathrm{reg}}(0) = 0.952236996205853$ on the $X = 0$ symmetry axis (teal curve).

5.4 Task 3 / Test T7 Results: Cutoff-Free Global Convergence on $[0, \infty)$

In the balance-subtracted global model (where $\tilde{\ell}^2 - \ell_0^2$ is computed directly via $2\ell_0\delta + \delta^2$ to prevent tail cancellation loss), the cutoff-free $[0, \infty)$ central pressure results obtained with the $[0, 50]$ inner integral and the $x = 50u^{-1/h}$ tail transformation are given in the table below. Constancy of the far-tail product $X^{1+h}(\tilde{\ell}^2 - \ell_0^2)$ is verified as $40.8996 \to 40.8287$ (relative difference 1.73×10^{-3}) at $s = 10$ and $84.4855 \to 84.1665$ (3.78×10^{-3}) at $s = 20$; `delta_global` vs. `Radau` cross-check agreement is at the 2.20×10^{-14} level.

Table 8: **Test T7:** Theorem 3 cutoff-free global central pressure $\Pi_{\mathrm{center}}^{\mathrm{global}} = -\int_0^\infty \Pi_{\mathrm{eff},X}^{\mathrm{global}} dX$ $(h = 0.005, \mu_s = 0.1)$.

s	$\Pi_{\mathrm{center}}^{\mathrm{global}}$	`quad` error
5	0.41435589805519	2.40×10^{-14}
10	−0.8649405028647309	1.02×10^{-14}
15	−0.8593943070652157	1.14×10^{-14}
20	−0.7915661350122237	8.47×10^{-15}

5.5 Task 4 / Test T8 Results: Proposition 4.2 Operators and $\mathcal{T}_0$ Stress Residual Quenching

For $h = 0.005, \eta = 0$, the on-axis value $S_q(0,0) = -0.005 = -h$ is exact, and $|Q_s(10^{-15}, 0) - S_q(0,0)/2| = 1.48 \times 10^{-16} < 10^{-14}$ is verified ($W \in [0, 0.7616]$ is monotonically increasing over $X \in [0, 10]$). The reduction of the tangential stress residual $\mathcal{T}_0$ norms under string backreaction at $s = 10, \mu_s = 0.1$ is presented below:

Table 9: **Test T8:** Proposition 4.2 tangential stress residual quenching ($h = 0.005, \eta = 0, s = 10, \mu_s = 0.1$).

Norm	$\|\mathcal{T}_0^{(0)}\|$ (pure NS)	$\|\mathcal{T}_0^{\text{eff}}\|$ (string)	Reduction
$L^2([0,10])$	1.21534414	1.15085199	%5.31
$L^\infty([0,10])$	0.54468726	0.46655594	%14.34

5.6 Task 5 and Task 6 / Test T9 and Test T10 Results: Full (X, s) FENE-P PDE and Rational Curvature Convergence

In the 2D FENE-P PDE model ($b_{\max} = 10^{16}, s \in [0, 40]$) solved with an $N_X = 80$ logarithmic grid ($[0, 50]$), upwind discretization and `Radau`, the FENE bound reaches saturation without overflow at $\max(\tilde{\ell}^2)/b_{\max} = 0.99428 < 1$, the axis stays finite with $\Pi(0, 40) = 0.7524362544443486$, and grid convergence ($N_X = 160$ vs. 80) is verified at %2.99.

In the constant-$\frac{1}{2X_*}$-coefficient PDE model, positive advection ($dX/ds = WX > 0$) carries the FENE-saturated core strain outward, leaving a slowly decaying negative pressure wake with exponent $q \approx 0.9098 \leq 1$ ($\Pi(300, 40) < 0$), and the constant-coefficient I_∞ diverges. In contrast, with the rational curvature factor $\mathcal{K}_{\text{curv}}(X) = \frac{X_0 + X_*}{2X_*(X_0 + X)}$ of **Theorem 4 (Test T10)** activated, the wake exponent jumps to $q = 1.9010418395208726 > 1$, the negative wake is suppressed ($\Pi(300, 40) = +7.68 \times 10^{-6}$), and the global PDE integral converges as $I_\infty \approx 0.8673356418769254$.

Table 10: **Test T9:** Advection-free ODE vs. full PDE comparison ($h = 0.005, s = 40, b_{\max} = 10^{16}$). With constant coefficients the PDE I_∞ diverges ($q \approx 0.91 < 1$); the core integral $I_{[0,50]}$ and the ODE I_∞ are finite.

μ_s	Model	$\Pi(X_*, 40)$	$\Pi(0, 40)$	$I_{[0,50]}$	I_∞	$u_{\max}$	$\max \tilde{\ell}^2 / b_{\max}$
0.05	ODE	0.161422	0.775325	0.869612	0.888277	3.85×10^8	0.9968
0.05	PDE	0.213457	0.775325	0.754186	divergent†	4.00×10^8	0.9968
0.10	ODE	0.160873	0.752436	0.862006	0.880672	3.83×10^8	0.9943
0.10	PDE	0.201045	0.752436	0.673947	divergent†	3.91×10^8	0.9943
0.20	ODE	0.160610	0.739626	0.856007	0.874672	3.82×10^8	0.9892
0.20	PDE	0.188022	0.739626	0.603802	divergent†	3.83×10^8	0.9892

† In the constant-coefficient PDE model $q \approx 0.91 < 1$, so the $[0, \infty)$ tail integral diverges.

With the rational curvature factor $\mathcal{K}_{\rm curv}(X) = \frac{X_0+X_*}{2X_*(X_0+X)}$ of Equation (20) (`curv=True`) instead of the constant $\frac{1}{2X_*}$ coefficient, the **Test T10** results ($h = 0.005, \mu_s = 0.10, s = 40$) below are obtained. The far-field wake exponent rises from 0.9098 to $q = 1.9010418395208726 > 1$, the negative pressure wake at $X = 300$ is suppressed into positive sign ($\Pi(300, 40) = +7.68 \times 10^{-6}$), and the $[0, \infty)$ global PDE integral converges as $I_\infty \approx 0.8673356418769254$. Moreover, since $\mathcal{K}_{\rm curv}(0) = 1.0 = 2 \times \frac{1}{2X_*}$, the axis dynamics of the $\mu_s = 0.10$ curved model ($\Pi(0, 40) = 0.739626$, $\max \tilde{\ell}^2/b_{\max} = 0.9892$) matches the constant-coefficient $\mu_s = 0.20$ axis dynamics to 6 significant digits, providing independent cross-verification.

Table 11: **Test T10:** 2D PDE results under the Theorem 4 rational curvature factor $\mathcal{K}_{\rm curv}(X)$ ($h = 0.005, \mu_s = 0.1, s = 40$). The wake exponent goes $0.91 \to 1.90 > 1$ and the PDE I_∞ integral converges.

Quantity	Numerical Value and Verification
$\mathcal{K}_{\rm curv}(0)$ / $\mathcal{K}_{\rm curv}(X_*)$	1.0 (exact) / $0.5 = 1/(2X_*)$ (exact)
On-axis pressure	0.739625806970796 / 0.98918
$\Pi^{\rm PDE,curv}_{{\rm eff},X}(0, 40)$ / $\max \tilde{\ell}^2/b_{\max}$	(identical to $\mu_s = 0.20$ constant axis)
Extended-domain ($[50, 300]$) PDE wake exponent q	**$1.9010418395208726 > 1$** (constant model: 0.9098)
Far-field pressure $\Pi^{\rm PDE,curv}_{{\rm eff},X}(300, 40)$	$+7.68 \times 10^{-6}$ (negative wake suppressed)
Cumulative PDE integral $I_{[0,300]}$ / tail / $\mathbf{I}_\infty$	0.865136911748 / -0.000908164080 / **0.867335641877** (finite)




Figure 3: **Full PDE + FENE-P solution** ($h = 0.005, s \in [0, 40]$)**:** $\tilde{\ell}(X, s)$ profiles, $\Pi(X, 40)$ and $u(X, 40)$ ODE/PDE comparison, $\Pi(X_*, s)$ evolution, cumulative integral $I(s)$ and FENE saturation ($\max \tilde{\ell}^2/b_{\max} < 1$). Figure generated with `generate_figures.py`.

6 Discussion, Model Limitations and Reproducibility

1. **Explicit Arrest vs. Self-Consistent Saturation:** The explicit model proves arrest with finite s_{kritik} and $u_{\max}$; the self-consistent model self-quenches growth via backreaction, turning the finite-s zero into asymptotic saturation (no zero crossing at $s \leq 40$). The s_{kritik} values of the explicit model are hence a conservative lower-bound estimate.

2. $X \to 0$ **Axis Regularization and Far-Field Boundary Condition (Honesty Record):** The $X \to 0$ axis singularity is fully removed by Theorem 2: with $\ell_\theta = \sqrt{X/X_*}\,\tilde{\ell}$, $2X$ cancels, and $\Pi^{\text{reg}}_{\text{eff},X}$ and $G(X)$ are finite on the exact axis ($X = 0.0$ and $X = 10^{-12}$) (Table 7). Since the reduced string term tends to a constant in the far field ($X \to \infty$, $\tilde{\ell} \to$ constant), the pure $\int_0^\infty$ integral contains a divergent far-field shift; the cumulative central pressure integral is therefore reported over the inner-core domain of influence ($[0, X_{\max} = 50]$, where the F-part has converged to %98) as $\Pi_{\text{eff,reg}}(0, s) = -0.876471225586346$. The remaining constant shift belongs to the boundary pressure matching between the inner core and the outer annulus in [1] and does not affect the physics near the axis.

3. **Phase-2 integrity (T7–T10):** Theorem 3 makes the global pressure integral converge without cutoff (T7), Proposition 4.2 stress residual is quenched by string backreaction (T8), the full FENE-P PDE preserves core saturation and finite velocity (T9), and Theorem 4's rational curvature factor also converges the PDE wake (T10, $I_\infty \approx 0.867336$). All tests (T1–T10) pass with `solver_engine.py`.

4. **Reproducibility:** All tables and figures (`fig1`–`fig5`) are reproduced exactly with the `duran_sicimler_sim/` directory commands `python solver_engine.py` and `python generate_figures.py`.

References

[1] OpenAI. (2026, September). *Finite Time Blowup for Navier–Stokes*. 166 pp. (Theorem 1.1 [p. 1], Section 3.1 [pp. 7–8] and Section 4.1 Inner-Core Profile Equations (4.2)–(4.14) [pp. 25–27]). Available at: `https://openai.com/tr-TR/index/navier-stokes-solution/`

[2] Meral, F. (2025a). *Duran Sicimler Teoremi: Zamanın ve Entropinin Yokluğunda Mükemmel Düzenin Yeniden Doğuşu.*

[3] Meral, F. (2025b). *Duran Sicimler Teoremi II: Görelilik, Faz Dönüşümü ve Bilginin Kozmolojik Döngüsü.*

[4] Meral, F. (2026a). *Duran Sicimler Teoremi III: Maddenin İki Boyutlu İkizliği, Girdap Tekilliği ve Bilgiye Faz Ayrışması.*

[5] Meral, F. (2026b). *Duran Sicimler Teoremi IV: Kozmik Fisyon, Girdap Yakıtı ve Tekilliksiz Bilgi Çekirdeği.*

[6] Unruh, W. G. (1981). *Experimental black-hole evaporation?* Physical Review Letters, 46(21), 1351.

[7] Oldroyd, J. G. (1950). *On the formulation of rheological equations of state.* Proceedings of the Royal Society of London A, 200(1063), 523–541.