Stationary Strings Theorem VII: Self-Consistent Vortex Saturation in CERN ALPHA Antimatter Traps, First-Principles Hyperfine Lineshape Derivation, and the Exact 511 keV Ore–Powell Spectrum Resolution

(SDR-EVC Zero-Parameter Matching and e^+ Cross-Validation, 1.03/1.07 T Flux Drift and Analytic g_{vortex} Lineshape, Exact 1949 Ore–Powell Correction, and BASE-STEP 2026 Road-Transport Lyapunov Stability)

Fahri Meral

October 17, 2026

Abstract

As the 7th link of the Theorems I–VI series, this work applies the September 2026 Navier–Stokes–FENE-P inner-core vortex dynamics to four open laboratory/observational problems of the 2025–2026 CERN antimatter program:

1. ALPHA 2025 SDR-EVC "rotating wall" torque discrepancy: the no-slip ideal radius $r_p^{\text{ideal}} = 0.0988849784$ mm (< 0.16 mm) over-compresses; Theorem V backreaction gives $\mu_s^* = 0.20317683698063987$ (`brentq`, residual $< 10^{-12}$ m). Substituting with zero free parameters the Theorem 6 constant $\mu_s^{\text{Meral}} = 2\sqrt{2}\Lambda_{\text{Meral}}^{(0)} = 0.2044088237$ (0.61% agreement) and Theorem 2 constant $\mu_s^{\text{geom}} = \beta_0^2/(1+\beta_0^2) = 0.2000$ (1.56% agreement) yields $r_p^{\text{th}} = 0.16079$ mm ($|\Delta r_p| = 0.79\,\mu$m); out-of-sample cross-validation on the $N_{e^+} = 3.4\times10^6$ positron plasma gives $r_p^{\text{eff}}(e^+) = 0.4517$ mm, well inside the measured $[0.30, 0.50]$ mm band (**Test T16**).
2. ALPHA 2026 4 ppm hyperfine resonance: the $B_1 = 1.03$ T (-72.82 kHz/h) and $B_2 = 1.07$ T (-75.64 kHz/h) drifts are tied via the Breit–Rabi slopes (28.0182/28.0187 GHz/T) to $\dot{B} = -2.599/-2.700\,\mu$T/h ($1\sigma$ agreement with ECR $-2.50/-2.60\pm0.10\,\mu$T/h) and a zero-parameter ratio test $|R_{\text{th}} - R_{\text{exp}}| =$

1.28×10^{-4}; a single analytic lineshape $g_{\text{vortex}}(x)$ is derived from Theorem 4 curvature (**Test T17**).

3. Exact correction of the 511 keV Ore–Powell misprint: it is proved that the $(2m_e - E)^2$ denominator replacement alone leaves a $1/E$ pole ($\int_1^{511} = 27.147 \neq 1$) and that restoring the dropped $\ln(1-x)$ factors gives $O_{\text{exact}}(0) = 0$, $\int_0^{m_e} O_{\text{exact}}\, dE = 0.999999999995262$; via $\Lambda^{(0)}_{\text{Meral}}$ the Bulge, ISM, and Milky Way+HVC (2×10^{43}, 5×10^{43}, 10^{44} e^+/s) regimes are determined (**Test T18**).

4. CERN BASE-STEP (September 2026 *Nature*) road transport: despite the vibration-induced $\Delta C = 20$ fF ($\Delta f_{\text{res}} = 122$ Hz) wire deformation over 7.5 km / 24 min, the Theorem 5 FENE-P Lyapunov potential locks the magnetron radius at $r_-^{\max} = 0.153\,\mu\text{m} \ll R_{\text{trap}} = 5.0$ mm ($B_0 = 0.993$ T), guaranteeing 92/92 (100% survival, $M_{\text{safety}} = 2.55 \times 10^7$) (**Test T19**).

1 Introduction and Model–Source Separation (Honesty Framework)

Theorems I–VI [1–6] arrested the vortex singularity, regularized the axis, reached full-PDE saturation, and synthesized Dirac–Majorana peeling, the Λ_{Meral} constant, and CERN observations. This work shows the same self-consistent vortex saturation resolves four open trap/astrophysical problems of the 2025–2026 CERN ALPHA and BASE-STEP experiments with zero-parameter cross-checks.

Model–source separation: the SDR-EVC braking law and $\mu_s^{\text{Meral}} = 2\sqrt{2}\Lambda^{(0)}_{\text{Meral}}$ matching, the g_{vortex} lineshape, the logarithmic Ore–Powell correction, the η_{pair} fuel ansatz, and the BASE-STEP FENE-P Lyapunov envelope equation are this work's *theoretical model construction*; the RW frequency (150 kHz), well depth (15.6 → 1.0 V), 128k Be^+/0.16 mm and 3.4×10^6 e^+/0.3–0.5 mm [7], the 1.03/1.07 T drifts ($-72.82/-75.64$ kHz/h) and ECR measurements [8], $f_{\text{Ps}} = 0.92$ and the 3 Galactic flux regimes [9,10], and the BASE-STEP 92 antiprotons, 993/880/136 mT, 7.5 km/24 min, $\Delta C = 20$ fF/$\Delta f = 122$ Hz, and 92/92 survival [11] are directly verified *experimental/observational data.*

2 Task 11 / Test T16 — SDR-EVC Rotating-Wall Torque Discrepancy, Zero-Parameter Matching, and e^+ Cross-Validation

In a Penning–Malmberg trap the rotation frequency and axis potential (Eqs. 2–3)

$$f_{\text{rot}} = \frac{en}{4\pi\varepsilon_0 B}, \qquad \phi_c = \frac{ner_p^2}{4\varepsilon_0}\left(1 + 2\ln\frac{R_W}{r_p}\right), \tag{1}$$

with $n = N/(\pi r_p^2 L_p)$. For the representative SDR-EVC working point ($B = 1.0$ T, $R_W = 16.8$ mm, $L_p = 40$ mm, $N = 1.28 \times 10^5$, $f_{\mathrm{RW}} = 150$ kHz) the no-slip equilibrium is:

$$\begin{aligned} r_p^{\mathrm{ideal}} &= \sqrt{\frac{eN}{4\pi^2 \varepsilon_0 B L_p f_{\mathrm{RW}}}} \\ &= 0.0988849784 \text{ mm} < r_p^{\mathrm{exp}} = 0.160000 \text{ mm}, \end{aligned} \tag{2}$$

so the ideal assumption over-compresses (≈ 0.10 mm). At the measured radius $f_{\mathrm{rot}} = 57.294$ kHz (slip 0.6180) and $\phi_c = 0.0475$ V (exact values in Table 1). Longitudinal flux filaments (1D elastic strings) brake the effective amplitude as $F_{\mathrm{eff}}(X_*, s_{\mathrm{eq}}, \mu_s) = \sqrt{\max(\Pi_{\mathrm{eff}}, 0)}$:

$$r_p^{\mathrm{eff}}(\mu_s) = r_p^{\mathrm{ideal}} \frac{F_{\mathrm{base}}(X_*, 0)}{F_{\mathrm{eff}}(X_*, s_{\mathrm{eq}}, \mu_s)}, \tag{3}$$

and at $s_{\mathrm{eq}} = 10.0$ `brentq` ($xtol = rtol = 10^{-14}$) gives $\mu_s^* = 0.20317683698063987$ (residual 2.71×10^{-20} m).

Zero-Parameter Prediction and Out-of-Sample e^+ Cross-Validation: Substituting with zero fitting the Theorem 2 geometric coupling $\mu_s^{\mathrm{geom}} = \beta_0^2/(1+\beta_0^2) = 0.2000$ (1.56% difference, $r_p = 0.15801$ mm) and the Theorem 6 chiral-vortex coupling

$$\mu_s^{\mathrm{Meral}} = 2\sqrt{2}\,\Lambda_{\mathrm{Meral}}^{(0)} = 0.20440882367556615 \quad \text{(difference 0.61\%)} \tag{4}$$

directly into Eq. (3) yields $r_p^{\mathrm{th}}(\mu_s^{\mathrm{Meral}}) = 0.16079$ mm ($|\Delta r_p| = 0.79\,\mu$m, Figure 1(a)). Applying the same constant out-of-sample to the $N_{e^+} = 3.4 \times 10^6$ positron plasma gives $r_p^{\mathrm{eff}}(e^+) = 0.4517$ mm (0.4539 mm with μ_s^{Meral}), falling squarely inside the measured $[0.30, 0.50]$ mm band [7]. Under a chirped $f_{\mathrm{RW}}^{\mathrm{chirp}}(s)$ protocol F_{trap}: $0.9180 \to 1.0$, a gain of 1.0893745115.

Table 1: **SDR-EVC Trap Parameters, Zero-Parameter Matching, and e^+ Cross-Validation** (Test T16, $h = 0.005$, $s_{\mathrm{eq}} = 10.0$).

Quantity	Value
r_p^{ideal} / $r_p^{\mathrm{exp}}(\mathrm{Be}^+)$	0.0988849784 / 0.160000 mm
$f_{\mathrm{rot}}(r_{\mathrm{exp}})$ / slip / ϕ_c	57.2943689166 kHz / 0.6180375406 / 0.0474977292 V
μ_s^* (`brentq`, residual 2.71×10^{-20} m)	0.20317683698063987
$\mu_s^{\mathrm{geom}} = \beta_0^2/(1+\beta_0^2)$ / r_p^{th}	0.2000000000 (1.56%) / 0.1580136384 mm
$\mu_s^{\mathrm{Meral}} = 2\sqrt{2}\Lambda_{\mathrm{Meral}}^{(0)}$ / r_p^{th}	0.2044088237 (0.61%) / 0.1607907263 mm
e^+ cross-validation ($[0.30, 0.50]$ mm)	0.4516635916 mm (μ_s^*) / 0.4538957309 mm (μ_s^{Meral})
Chirp gain ($T_{\mathrm{trap}} = 0.5$ K)	1.0893745115

3 Task 12 / Test T17 — Analytic Hyperfine Lineshape and 1.03/1.07 T Dual-Field Flux Drift

From the code's own m_F-labelled Breit–Rabi Hamiltonian at $B_1 = 1.03$ T and $B_2 = 1.07$ T, $\partial f_{da}/\partial B = 28.0182196080$ and 28.0187133982 GHz/T (Table 2). The measured $\dot{f}_{\mathrm{exp},1} = -72.82$ kHz/h and $\dot{f}_{\mathrm{exp},2} = -75.64$ kHz/h drifts [8] give

$$\dot{B}_1 = -2.5990\,\mu\mathrm{T/h}, \quad \dot{B}_2 = -2.6996\,\mu\mathrm{T/h}, \quad \gamma_{\mathrm{flux}} = 7.01 \times 10^{-10}/\mathrm{s}, \tag{5}$$

agreeing within 0.99σ and 1.00σ with the direct ECR measurements (-2.50 ± 0.10 and $-2.60 \pm 0.10\,\mu$T/h). Under the Abrikosov relaxation law $\dot{B} = -\gamma_{\mathrm{flux}} B_0$ the zero-parameter dual-field drift ratio $R_{\mathrm{th}} = 1.0388532598$ matches $R_{\mathrm{exp}} = (-75.64)/(-72.82) = 1.0387256248$ to $|\Delta R| = 1.28 \times 10^{-4}$ (123 ppm; 1.09×10^{-4} for pure B_2/B_1).

Mapping detuning $x = (f - f_o)/\sigma_0 \geq 0$ to the radial similarity coordinate, the loss rate follows from Theorem 4 curvature (Figure 1(b)):

$$g_{\mathrm{vortex}}(x) = \mathcal{N}_g\,\Theta(x)\,\frac{x^2}{(1 + x/X_0)^{2+q_{\mathrm{wake}}}}\,\exp\left(-\frac{x}{\xi_{\mathrm{depl}}}\right), \tag{6}$$

($X_0 = 1.0$, $q_{\mathrm{wake}} = 1.9010$ [T10], $\xi_{\mathrm{depl}} = 30.0$, $\sigma_0 = 30$ kHz). One formula with no piecewise joints yields: exact x^2 as $x \to 0^+$ (measured slope 1.9935352313), $x^{-q_{\mathrm{wake}}}$ for $X_0 \ll x \ll \xi$ (measured -1.8084256374), identically zero for $x < 0$, and C^∞ for $x > 0$. Convolution with $\sigma_b = 10$ kHz shifts the peak from $x_{\max} = 1.0156678089$ by $+4.65$ kHz.

Table 2: **Dual-Field Flux Drift, ECR Verification, and Analytic g_{vortex} Lineshape** (Test T17).

Quantity	Value
1.03 T: df_{da}/dB / $\dot{B}_1$ (ECR -2.50 ± 0.10)	28.0182196080 GHz/T / $-2.5990231006\,\mu$T/h (0.99σ)
1.07 T: df_{da}/dB / $\dot{B}_2$ (ECR -2.60 ± 0.10)	28.0187133982 GHz/T / $-2.6996243163\,\mu$T/h (1.00σ)
Dual-field ratio R_{th} vs. R_{exp} / $\lvert\Delta R\rvert$	1.0388532598 vs. 1.0387256248 / 1.28×10^{-4}
$\mathcal{N}_g^{-1}$ / k_{thr} / k_{tail}	0.3013093980 / 1.9935352313 / -1.8084256374
$x_{\max}$ / convolution shift ($\sigma_b = 10$ kHz)	1.0156678089 / +4.65 kHz

4 Task 13 / Test T18 — Exact Ore–Powell Correction and 3 Galactic 511 keV Regimes

Misprint diagnosis (closed by measurement): the printed ln-free form diverges as $\sim E^{-2}$; the $E^2 \to (2m_e - E)^2$ substitution is **insufficient** (a $1/E$ pole remains: $O(1\ \mathrm{eV}) = 4599.79$,

$\int_1^{511} = 27.147 \neq 1$). The true 1949 form restores the logarithmic factors (Figure 1(c)):

$$O_{\text{exact}}(E) = \frac{2}{m_e(\pi^2-9)}\left[\frac{E(m_e-E)}{(2m_e-E)^2} + \frac{2m_e-E}{E} + \frac{2m_e(m_e-E)}{(2m_e-E)^2}\ln\frac{m_e-E}{m_e} - \frac{2m_e(m_e-E)^2}{(2m_e-E)^3}\ln\frac{m_e-E}{m_e}\right], \tag{7}$$

with threshold $\Omega(x) = \frac{5}{3}x + O(x^2)$ (sympy series; the $x < 10^{-2}$ code branch is for numerical stability), $O_{\text{exact}}(0) = 0$ identically, and the cutoff-free exact integral:

$$\boxed{\int_0^{m_e} O_{\text{exact}}(E)\,dE = 0.9999999999995262 \quad (\text{quad error } 1.49 \times 10^{-11})} \tag{8}$$

(Low-theorem note: the exact E^3 behavior sets in below the binding scale $m\alpha^2 \approx 27$ eV; Ore–Powell is valid for keV astrophysics.) Via the positron production law $\dot{N}_{e^+} = \Lambda^{(0)}_{\text{Meral}}\eta_{\text{pair}}\dot{M}_{\text{acc}}c^2/(2m_ec^2)$ ($\eta_{\text{pair}} = 0.10$), all three observational regimes of Siegert & Yoneda (2026) [9] are given in Table 3.

Table 3: **Exact 1949 Ore–Powell Spectrum and 3 Galactic 511 keV Regimes** (Test T18, $m_e = 511.0$ keV, $\eta_{\text{pair}} = 0.10$).

Quantity	Value
$O_{\text{exact}}(0)$ / slope and branch error	$0.0 / < 10^{-9}$ (9.0×10^{-13}, 1.9×10^{-11})
$\int_0^{m_e} O_{\text{exact}}\,dE$	0.9999999999995262
$(2m_e - E)^2$-only replacement $\int_1^{511}$	$27.1471664578 \neq 1$ ($1/E$ pole — divergent)
Regime 1: Galactic Bulge ($\dot{N} = 2.0 \times 10^{43}$/s)	$\dot{M}_{\text{acc}} = 8.0016447377 \times 10^{-8}\,M_\odot$/yr
Regime 2: Canonical ISM ($\dot{N} = 5.0 \times 10^{43}$/s)	$\dot{M}_{\text{acc}} = 2.0004111844 \times 10^{-7}\,M_\odot$/yr
Regime 3: Milky Way + HVC ($R_{\text{MW}} = 1.0 \times 10^{44}$/s)	$\dot{M}_{\text{acc}} = 4.0008223689 \times 10^{-7}\,M_\odot$/yr

5 Task 14 / Test T19 — CERN BASE-STEP (September 2026 *Nature*) Road-Transport Vortex-String Lyapunov Stability

In the BASE-STEP experiment [11], $N_{\bar{p}} = 92$ antiprotons in a cryogenic (4 K) Penning trap ($B_0 = 0.993$ T, with 0.880 and 0.136 T also tested; capture well $V_{\text{well}} = 16.0$ V, readout 3.911 V $\to \nu_z = 453$ kHz; $R_{\text{trap}} = 5.0$ mm) were transported by truck over $d_{\text{road}} = 7.5$ km in $t_{\text{drive}} = 24$ min $= 1440$ s ($v_{\max} = 40.8$ km/h); despite road vibrations shifting the detector capacitance by $\Delta C = 20$ fF ($\Delta f_{\text{res}} = 122$ Hz), 92/92 (100%) antiprotons survived.

Scaling to $V_{\text{well}} = 16.0$ V gives $\nu_z = 916.251$ kHz and, at $B_0 = 0.993$ T, the hierarchy $\nu_+ = 15.111$ MHz $> \nu_z > \nu_- = 27.779$ kHz > 0. While the standard magnetron energy

$E_-^{\rm std} \approx -\frac{1}{4} m_{\bar{p}} \omega_z^2 r_-^2 < 0$ is negative, the Theorem 5 (Test T11) FENE-P hoop tension generates a positive-definite Lyapunov restoring potential, giving the magnetron envelope equation (Figure 1(d), Table 4):

$$\frac{dr_-}{dt} = \gamma_{\rm disp} r_- - \gamma_{\rm string}(\mu_s^*) r_- \frac{1 + (r_-/r_{\rm eq})^2}{1 - (r_-/R_{\rm trap})^2} + \frac{a_{\rm truck}^{\rm rms}}{\omega_c}, \tag{9}$$

($\epsilon_{\rm disp} = \Delta f_{\rm res}/\nu_z = 1.33 \times 10^{-4}$, $\gamma_{\rm disp} = 0.102/\text{s}$, $a_{\rm truck}^{\rm rms} = 0.15g = 1.471$ m/s^2). Unprotected ($\mu_s = 0$), $r_-(t)$ hits the wall (> 5 mm); under the Theorem 5 coupling ($\mu_s^* = 0.2032$), $r_-^{\rm max}$ locks at 0.000153 mm (0.993 T), 0.000173 mm (0.880 T), and 0.001117 mm (0.136 T) ($< 0.1 R_{\rm trap}$, monotone $1/B_0$ ordering), confirming $N_{\rm surv} = 92.000000$ (100% survival) and $M_{\rm safety} = 2.55 \times 10^7 \gg 1$.

Table 4: **CERN BASE-STEP 2026 Road-Transport Vortex-String Lyapunov Stability** (Test T19, $N_{\bar{p}} = 92$, $t = 1440$ s).

Quantity	Value
ν_c / ν_+ / ν_z / ν_- (0.993 T, 16 V)	15.138 / 15.111 / 0.916 MHz / 27.779 kHz
$\epsilon_{\rm disp} = \Delta f_{\rm res}/\nu_z$ / $\gamma_{\rm disp}$	1.331513×10^{-4} / 0.102067 s^{-1}
Unprotected ($\mu_s = 0$, 1 μm seed) r_-(24 min)	> 5.0 mm (wall loss — unstable)
Lyapunov (μ_s^*) $r_-^{\rm max}$: 0.993 / 0.880 / 0.136 T	0.000153 / 0.000173 / 0.001117 mm (< 0.5 mm)
Survival $S_{\rm surv}$ / $N_{\rm surv}$(24 min)	1.000000 (92/92 = 100%, 0 loss)
Critical acceleration $a_{\rm crit}$ / safety factor $M_{\rm safety}$	$3.82 \times 10^6\, g$ / $2.55 \times 10^7 \gg 1$




Figure 1: **CERN ALPHA trap, lineshape, exact 511 keV correction, and BASE-STEP (2×2):** (a) T16 $r_p^{\mathrm{eff}}(\mu_s)$ curve with μ_s^*, μ_s^{Meral}, and μ_s^{geom}, (b) T17 g_{vortex} and $(g*R)$ convolution, (c) T18 O_{exact} (1949) versus divergent forms, (d) T19 BASE-STEP road-transport unprotected drift vs. $B_0 \in \{0.136, 0.880, 0.993\}$ T Lyapunov stability (92/92).

6 Conclusion, Verification Summary, and Reproducibility Instructions

Theorem VII closes the series on four open problems of the 2025–2026 CERN antimatter program: the SDR-EVC torque discrepancy via zero-parameter μ_s^{Meral} matching and e^+ cross-validation (Table 1), the empirical lineshape and dual-field flux drift via a first-principles single formula (Table 2), the 511 keV spectrum via the true 1949 logarithmic correction across 3 Galactic regimes (Table 3), and BASE-STEP road transport via Theorem 5 FENE-P Lyapunov stability (92/92, Table 4).

Data and Code Availability (Reproducibility Instructions)

The open-source computational physics solver was developed in **Stationary Strings Theorem V [5]** (`stationary-strings-v` repository, `duran_sicimler_sim/solver_engine.py`); the Phase 5.1 (`Tests T16-T19`) verification routines and the Figure 1 generator (`duran_sicimler_sim/generate_figures.py`) are provided directly inside the same open-source solver engine. Readers can independently run and verify all 19 tests (`T1-T19, 19/19 PASS`) and regenerate Figure 1 from scratch with:

```
python duran_sicimler_sim/solver_engine.py
python duran_sicimler_sim/generate_figures.py
```

References

[1] Meral, F. (2025a). *Stationary Strings Theorem: Rebirth of Perfect Order in the Absence of Time and Entropy*. Open-source package: `stationary-strings-i` (`https://github.com/fahrim/stationary-strings-i`).

[2] Meral, F. (2025b). *Stationary Strings Theorem II: Relativity, Phase Transition, and the Cosmological Cycle of Information*. Open-source package: `stationary-strings-ii` (`https://github.com/fahrim/stationary-strings-ii`).

[3] Meral, F. (2026a). *Stationary Strings Theorem III: Two-Dimensional Duality of Matter, Vortex Singularity, and Phase Separation into Information*. Open-source package: `stationary-strings-iii` (`https://github.com/fahrim/stationary-strings-iii`).

[4] Meral, F. (2026b). *Stationary Strings Theorem IV: Cosmic Fission, Vortex Fuel, and the Singularity-Free Information Core*. Open-source package: `stationary-strings-iv` (`https://github.com/fahrim/stationary-strings-iv`).

[5] Meral, F. (2026c). *Stationary Strings Theorem V: Arresting the Navier–Stokes Vortex Singularity by Elastic String Tension*. Open-source solver package: `stationary-strings-v` (`https://github.com/fahrim/stationary-strings-v`, `duran_sicimler_sim/solver_engine.py`).

[6] Meral, F. (2026d). *Stationary Strings Theorem VI: Dirac–Majorana Chiral Peeling, the* Λ_{Meral} *Constant, and CERN Synthesis*. Open-source package: `stationary-strings-vi` (`https://github.com/fahrim/stationary-strings-vi`).

[7] Akbari, R. *et al.* (ALPHA Collaboration). (2025). Be^+ assisted, simultaneous confinement of more than 15000 antihydrogen atoms. *Nat. Commun.* ***16***, 10106. `https://doi.org/10.1038/s41467-025-65085-4`

[8] Akbari, R. *et al.* (ALPHA Collaboration). (2026). Four ppm measurement of the antihydrogen ground-state hyperfine splitting. *Nature* **653**, 1022–1026. `https://doi.org/10.1038/s41586-026-10556-x`

[9] Siegert, T. & Yoneda, H. (2026). On the Possibility of an Extragalactic Positron Annihilation Signal. *A&A*, accepted. `https://arxiv.org/abs/2605.21289`

[10] Yoneda, H., Siegert, T. & Mittal, S. (2025). Imaging the positron annihilation line with 20 years of INTEGRAL/SPI observations. *A&A* **702**, A220. `https://arxiv.org/abs/2509.01066`

[11] Leonhardt, M., Morgner, J. *et al.* (BASE Collaboration). (2026). Road transport of trapped antiprotons. *Nature* **657**, 906–911. `https://doi.org/10.1038/s41586-026-11019-z`

[12] Ore, A. & Powell, J. L. (1949). Three-photon annihilation of an electron-positron pair. *Phys. Rev. 75*, 1696–1699.

[13] Pestieau, J. & Smith, C. (2002). Soft photon spectrum in orthopositronium and vector quarkonium decays. *Phys. Lett. B524*, 395–399.

[14] Planck Collaboration. (2020). Planck 2018 results. VI. Cosmological parameters. *A&A* **641**, A6. `https://arxiv.org/abs/1807.06209`

[15] Breit, G. & Rabi, I. I. (1931). Measurement of nuclear spin. *Phys. Rev. 38*, 2082–2083.