

Lecture 1: Mathematical Foundations for Lattice-Based Cryptography

Groups, Rings, Fields, Modular Arithmetic,
Vector Spaces, Basis, and Lattices

Dr. Faisal Aslam

Lecture Notes — Session 1 (approx. 2 hours)

Purpose of this lecture. Every technical term used later in the thesis proposal — “lattice,” “modulus,” “ring $\mathbb{Z}_q[x]/(x^n + 1)$ ” — rests on a small set of algebraic building blocks. This lecture builds those blocks, one at a time, from the ground up: *Group* $\rightarrow$ *Subgroup* $\rightarrow$ *Ring* $\rightarrow$ *Field* $\rightarrow$ *Modular Arithmetic* $\rightarrow$ *Vector Space* $\rightarrow$ *Basis* $\rightarrow$ *Lattice*. Nothing here requires prior abstract algebra; every definition is followed immediately by concrete, worked examples.

Contents

1	Why Start With Algebra?	1
2	Sets and Binary Operations (Quick Refresher)	1
3	Groups	2
3.1	Subgroups	2
3.2	Cosets and Quotient Groups: Where $\mathbb{Z}_n$ Really Comes From	3
4	Rings	4
4.1	Zero Divisors: A Structural Reason Some Rings Aren’t Fields	5
5	Fields	5
6	The Modulus: Arithmetic That Wraps Around	6
6.1	Finding Inverses Without Guessing: The Extended Euclidean Algorithm	7
7	Vector Spaces, Basis, and Dimension	8
7.1	Inner Products, Norms, and Orthogonality	8
7.2	Gram–Schmidt Orthogonalization	9
8	Lattices, Properly Defined	10
8.1	Discrete Sets	10
8.2	The Definition	10
8.3	Recovering the Basis Picture	11
8.4	The Determinant (Volume) of a Lattice	13
9	Summary Table	14
10	Full List of Exercises (Assign as Homework Before Lecture 2)	15

1 Why Start With Algebra?

Lattice-based cryptography is, at its core, about doing arithmetic on *vectors* whose coordinates live in a *finite, wraparound number system*, arranged according to strict algebraic rules. Before we can even state what a lattice is, we need precise answers to questions like:

- What does it mean for a set of numbers to have a well-behaved notion of “add” and “multiply”? (**Groups, Rings, Fields**)
- What happens when arithmetic “wraps around,” like a clock? (**Modulus**)
- What does it mean to build every point in a space out of a small number of building-block directions? (**Vector spaces, Basis**)
- What happens when we only allow *whole-number* combinations of those building blocks, instead of any real-number combination? (**Lattice**)

This lecture answers each of these in order. By the end, the definition of a lattice (Section 8) should feel like the natural, inevitable next step, not an isolated new idea.

2 Sets and Binary Operations (Quick Refresher)

A **set** is just a collection of objects (numbers, vectors, symbols — anything). A **binary operation** on a set S is a rule that takes two elements of S and combines them into a third element, e.g., ordinary addition $+$ takes two numbers and produces their sum. We write $a * b$ for a general binary operation.

An operation is **closed** on S if combining two elements of S always gives another element of S (never something outside S). This sounds obvious but fails more often than you’d expect — see Example 1 below.

Example: Closure Can Fail

Let $S = \{1, 2, 3, 4, 5\}$ (just these five numbers) and let $*$ be ordinary addition. Then $2 * 4 = 6$, but $6 \notin S$. So ordinary addition is *not* closed on S . This is exactly the kind of problem modular arithmetic (Section 5) is built to fix.

3 Groups

Definition: Group

A set G together with a binary operation $*$ is called a **group**, written $(G, *)$, if all four of the following hold:

(G1) **Closure:** for all $a, b \in G$, $a * b \in G$.

(G2) **Associativity:** for all $a, b, c \in G$, $(a * b) * c = a * (b * c)$.

(G3) **Identity:** there exists an element $e \in G$ such that $a * e = e * a = a$ for all $a \in G$.

(G4) **Inverses:** for every $a \in G$, there exists $a^{-1} \in G$ such that $a * a^{-1} = a^{-1} * a = e$.

If, in addition, $a * b = b * a$ for all $a, b \in G$ (the order doesn’t matter), the group is called **abelian** (commutative).

Example: Groups You Already Know

- $(\mathbb{Z}, +)$ — the integers under addition. Identity is 0; the inverse of a is $-a$. This is an abelian group.
- $(\mathbb{R} \setminus \{0\}, \times)$ — nonzero real numbers under multiplication. Identity is 1; the inverse of a is $1/a$. Also abelian. (We had to remove 0 because 0 has no multiplicative inverse.)
- $(\mathbb{Z}, \times)$ — integers under multiplication — is **not** a group: 2 has no inverse in $\mathbb{Z}$ (we'd need $1/2$, which isn't an integer). This fails axiom (G4).

Remark: Why Cryptography Cares About Groups

Many classical cryptographic schemes (Diffie–Hellman, ElGamal, elliptic-curve cryptography) are built directly on top of group structure — the security relies on certain problems (like “given g and g^a , find a ”) being hard inside a specific group. Lattice-based cryptography instead builds on richer structures (rings and modules, coming next), but the group is always the first layer underneath.

Exercise: Try This Before Next Lecture

Is $(\mathbb{Z}_5 \setminus \{0\}, \times \bmod 5) = (\{1, 2, 3, 4\}, \times \bmod 5)$ a group? Check all four axioms directly by building the multiplication table. (You'll need Section 6 for what “mod 5” means if it's new to you — come back to this after reading that section.)

3.1 Subgroups

There is one more group-related idea we need before we can properly define a lattice: a group living *inside* another group.

Definition: Subgroup

Let $(G, *)$ be a group. A subset $H \subseteq G$ is a **subgroup** of G if H is itself a group under the same operation $*$. In practice, this is easiest to check with a simple test: H is a subgroup of G if and only if (1) H is nonempty (usually: it contains the identity e), (2) H is closed under $*$ (for $a, b \in H$, $a * b \in H$), and (3) H is closed under inverses (for $a \in H$, $a^{-1} \in H$). Associativity is automatic, since it already held for all of G .

Example: Subgroups You Can Picture

- For any integer n , the set $n\mathbb{Z} = \{\dots, -2n, -n, 0, n, 2n, \dots\}$ (all multiples of n) is a subgroup of $(\mathbb{Z}, +)$: it contains 0, adding two multiples of n gives another multiple of n , and $-(kn)$ is also a multiple of n . For example, $2\mathbb{Z}$ is “all even integers.”
- The natural numbers $\mathbb{N} = \{0, 1, 2, \dots\}$ are **not** a subgroup of $(\mathbb{Z}, +)$: $\mathbb{N}$ is closed under addition, but it fails closure under inverses — $1 \in \mathbb{N}$, but $-1 \notin \mathbb{N}$.

Looking Ahead: Why This Matters in Two Sections

Keep the example $n\mathbb{Z} \subseteq \mathbb{Z}$ in mind. In Section 8, we will define a lattice as a special kind of subgroup of $\mathbb{R}^n$ — and $n\mathbb{Z}$ is exactly a simple, one-dimensional lattice in disguise: an evenly-spaced, discrete set of points, closed under addition and negation, sitting inside the larger group $\mathbb{Z}$ (or $\mathbb{R}$).

3.2 Cosets and Quotient Groups: Where $\mathbb{Z}_n$ Really Comes From

Section 6 will introduce $\mathbb{Z}_n$ informally, as “remainders with wraparound arithmetic.” That’s the right intuition, but it’s worth seeing the fully rigorous construction now, because it is a direct, satisfying payoff of the subgroup idea — and it is the same construction (“take a group, collapse it by a subgroup”) that shows up again later when we build the ring $R_q = \mathbb{Z}_q[x]/(x^n + 1)$ that Kyber and Dilithium actually use.

Definition: Coset

Let H be a subgroup of an abelian group $(G, +)$. For $a \in G$, the **coset** $a+H$ is the set $\{a+h : h \in H\}$ — “shift H by a .” Two elements $a, b \in G$ turn out to produce the *same* coset, $a+H = b+H$, exactly when $a-b \in H$. This gives a clean way to sort all of G into non-overlapping “buckets,” each one a shifted copy of H .

Example: The Cosets of $3\mathbb{Z}$ in $\mathbb{Z}$

Take $G = \mathbb{Z}$ and $H = 3\mathbb{Z} = \{\dots, -3, 0, 3, 6, \dots\}$ (a subgroup, per the “Subgroups You Can Picture” example in Section 3.1). The cosets are:

- $0 + 3\mathbb{Z} = \{\dots, -3, 0, 3, 6, \dots\}$ (multiples of 3)
- $1 + 3\mathbb{Z} = \{\dots, -2, 1, 4, 7, \dots\}$ (remainder 1 when divided by 3)
- $2 + 3\mathbb{Z} = \{\dots, -1, 2, 5, 8, \dots\}$ (remainder 2 when divided by 3)

Every integer falls into exactly one of these three buckets — and which bucket it falls into is precisely its remainder mod 3. There is no fourth distinct coset: $3 + 3\mathbb{Z} = 0 + 3\mathbb{Z}$ (check: $3 - 0 = 3 \in 3\mathbb{Z}$), so the pattern repeats.

Definition: Quotient Group

The set of all distinct cosets of H in G , with addition defined by $(a+H) + (b+H) := (a+b)+H$, is itself a group — the **quotient group**, written G/H (“ $G \bmod H$ ”). Its identity element is the coset $0 + H = H$ itself.

Remark: $\mathbb{Z}_n$ Is Not Just Like $\mathbb{Z}/n\mathbb{Z}$ — It Is $\mathbb{Z}/n\mathbb{Z}$

Compare the coset example above to Section 6: the three cosets of $3\mathbb{Z}$ in $\mathbb{Z}$ correspond *exactly* to the three elements $\{0, 1, 2\}$ of $\mathbb{Z}_3$, and coset addition matches addition mod 3 exactly. This is not a coincidence or an analogy — it is a theorem: $\mathbb{Z}_n$, as defined informally in Section 6, is precisely the quotient group $\mathbb{Z}/n\mathbb{Z}$. “Wraparound arithmetic” and “sorting integers into cosets of $n\mathbb{Z}$ ” are two descriptions of the exact same object. This is the rigorous justification for a claim we made without proof in Section 6: that $\mathbb{Z}_n$ really is a well-defined group (in fact ring) under addition and multiplication with wraparound — it inherits this structure directly from being a quotient of $\mathbb{Z}$.

Exercise: Practice

Let $H = 4\mathbb{Z} \subseteq \mathbb{Z}$. List all distinct cosets of H in $\mathbb{Z}$, and confirm there are exactly 4 of them, matching $\mathbb{Z}_4 = \{0, 1, 2, 3\}$. Which coset does -5 belong to?

4 Rings

A ring is what you get when you ask a set to support *two* operations — usually called “+” and “×” — that interact the way ordinary addition and multiplication do.

Definition: Ring

A set R with two binary operations $+$ and $\times$ is a **ring** $(R, +, \times)$ if:

(R1) $(R, +)$ is an abelian group (with identity element usually written 0).

(R2) $\times$ is associative: $(a \times b) \times c = a \times (b \times c)$.

(R3) The distributive laws hold: $a \times (b + c) = (a \times b) + (a \times c)$ and $(a + b) \times c = (a \times c) + (b \times c)$.

Note: a ring does *not* need multiplicative inverses, and multiplication need not be commutative in general (though in every ring we use in this thesis, it will be — these are called **commutative rings**). If there is an element $1 \in R$ with $a \times 1 = 1 \times a = a$ for all a , we say R is a **ring with unity**.

Example: Rings You Already Know

- $(\mathbb{Z}, +, \times)$ — the integers form a commutative ring with unity 1. Notice 2 still has no multiplicative inverse, and that’s fine — rings don’t require one.
- **Polynomial rings**, e.g. $\mathbb{Z}[x]$: the set of all polynomials with integer coefficients (like $3x^2 - x + 7$), with ordinary polynomial addition and multiplication. This is a ring for exactly the same reason $\mathbb{Z}$ is.

Looking Ahead: The Ring That Kyber and Dilithium Actually Use

Both Kyber and Dilithium do their arithmetic inside a specific ring of the form

$$R_q = \mathbb{Z}_q[x]/(x^n + 1),$$

read as: “polynomials with coefficients taken modulo q (Section 6), where we also reduce powers of x using the rule $x^n \equiv -1$.” This looks intimidating right now, but it is built from exactly the pieces in this lecture: a polynomial ring (this section), with coefficients from $\mathbb{Z}_q$ (Section 6), further “folded” by a fixed rule. We will unpack this fully in a later lecture — for now, just notice that “ring” is not an abstract curiosity; it’s the literal object the cryptography is built on.

4.1 Zero Divisors: A Structural Reason Some Rings Aren’t Fields

We will soon see (Section 5) that $\mathbb{Z}_n$ is a field exactly when n is prime. Section 6.1 will justify this computationally, via gcd. Here is the deeper, structural reason *why* — worth seeing now, while we’re still inside general ring theory.

Definition: Zero Divisor

In a ring R , a nonzero element a is a **zero divisor** if there exists some *nonzero* $b \in R$ with $a \times b = 0$. A commutative ring with unity that has *no* zero divisors (other than needing a or b to be 0) is called an **integral domain**.

Example: Zero Divisors in $\mathbb{Z}_6$, Absent in $\mathbb{Z}_5$

In $\mathbb{Z}_6$: $2 \times 3 = 6 \equiv 0 \pmod{6}$, but neither 2 nor 3 is 0. So 2 and 3 are both zero divisors — $\mathbb{Z}_6$ is *not* an integral domain. Contrast this with $\mathbb{Z}_5$: checking every pair of nonzero elements (1, 2, 3, 4), no product is ever $\equiv 0 \pmod{5}$ (e.g., $2 \times 3 = 6 \equiv 1$, $4 \times 4 = 16 \equiv 1$, etc.) — $\mathbb{Z}_5$ has no zero divisors.

Remark: Why This Explains the Prime/Composite Split

Fact: every field is automatically an integral domain (if $a \neq 0$ and $ab = 0$, multiply both sides by a^{-1} to get $b = 0$ — so no nonzero b can pair with a nonzero a to give 0). Contrapositive: *if a ring has a zero divisor, it cannot be a field.* Now, $\mathbb{Z}_n$ has a zero divisor exactly when n is composite: write $n = jk$ with $1 < j, k < n$; then $j \times k = n \equiv 0 \pmod{n}$, with neither j nor k equal to 0 in $\mathbb{Z}_n$. So every composite modulus produces zero divisors, which immediately rules out being a field — a purely structural explanation, with no gcd computation needed, of the same fact Section 6.1 reaches computationally.

Exercise: Practice

Find a pair of zero divisors in $\mathbb{Z}_{12}$ (there are several valid answers). Then explain, using the Fact above, why $\mathbb{Z}_{12}$ cannot be a field — without computing any inverses directly.

5 Fields

Definition: Field

A **field** is a commutative ring with unity F in which *every nonzero element has a multiplicative inverse* — that is, $(F \setminus \{0\}, \times)$ is also an abelian group. Informally: a field is a set where you can add, subtract, multiply, and divide (by anything except 0) and always stay inside the set.

Example: Fields and Non-Fields

- $\mathbb{Q}$ (rationals), $\mathbb{R}$ (reals), $\mathbb{C}$ (complex numbers) are all fields — you can divide by any nonzero element and stay inside the set.
- $\mathbb{Z}$ is **not** a field: $2 \in \mathbb{Z}$ has no multiplicative inverse in $\mathbb{Z}$ (again, $1/2 \notin \mathbb{Z}$). $\mathbb{Z}$ is a ring, but not a field.
- $\mathbb{Z}_5 = \{0, 1, 2, 3, 4\}$ under addition and multiplication mod 5 *is* a field — every nonzero element has an inverse (you checked this, or will check this, in the Section 3 subgroup/group exercise). Fields built this way, from $\mathbb{Z}_p$ with p prime, are called **finite fields** or **Galois fields**, written $\mathbb{F}_p$ or $GF(p)$.

Remark: The One-Sentence Summary So Far

Group $\subset$ Ring $\subset$ Field, in the sense that a field is a ring with extra structure (division), and a ring is built from a group with extra structure (a second operation). Each step adds a capability: groups let you undo one operation (inverses); rings add a second, distributive operation; fields guarantee *both* operations are fully invertible (except division by 0).

6 The Modulus: Arithmetic That Wraps Around

Definition: Congruence Modulo n

For a fixed positive integer n (the **modulus**), two integers a and b are **congruent modulo n** , written

$$a \equiv b \pmod{n},$$

if n divides $(a - b)$ evenly — equivalently, a and b leave the same remainder when divided by n . The set of possible remainders, $\{0, 1, 2, \dots, n - 1\}$, together with addition and multiplication performed “with wraparound,” is written $\mathbb{Z}_n$ (or $\mathbb{Z}/n\mathbb{Z}$).




Figure 1: The clock analogy for modular arithmetic: $\mathbb{Z}_{12}$ works exactly like a 12-hour clock face. 9 o'clock plus 5 hours wraps around to 2 o'clock, i.e., $9 + 5 = 14 \equiv 2 \pmod{12}$.

Example: Computing Modulo

$17 \bmod 5$: divide 17 by 5 to get 3 remainder 2, so $17 \equiv 2 \pmod{5}$. Similarly, $23 \equiv 3 \pmod{10}$, and $-1 \equiv n - 1 \pmod{n}$ (e.g., $-1 \equiv 4 \pmod{5}$) — negative numbers wrap around too.

Remark: When Is $\mathbb{Z}_n$ a Ring? When Is It a Field?

$\mathbb{Z}_n$ is *always* a commutative ring with unity, for any $n \geq 2$. But it is a *field* only when $n = p$ is **prime**. If n is composite (e.g., $n = 6$), some nonzero elements fail to have multiplicative inverses — e.g., in $\mathbb{Z}_6$, there is no x with $2x \equiv 1 \pmod{6}$, since 2 shares a common factor with 6. This is exactly why cryptographic schemes that need a field (rather than just a ring) are built with a prime modulus q . (Section 4.1 gave the deeper structural reason: composite n always produces zero divisors, and a ring with zero divisors can never be a field.)

6.1 Finding Inverses Without Guessing: The Extended Euclidean Algorithm

So far, the only way we've found a multiplicative inverse mod n is trial and error (Exercise: “find x with $3x \equiv 1 \pmod{7}$ ” — you could just try $x = 1, 2, 3, \dots$ until one works). That's fine for small numbers, but real cryptographic moduli are enormous (Kyber's modulus is $q = 3329$; RSA moduli have hundreds of digits) — trial and error is completely impractical. We need an actual algorithm.

Definition: Greatest Common Divisor (GCD)

For integers a, b (not both zero), $\gcd(a, b)$ is the largest positive integer that divides both a and b . Two integers are **coprime** (or relatively prime) if $\gcd(a, b) = 1$.

Remark: Fact: When Does an Inverse Exist?

An element $a \in \mathbb{Z}_n$ has a multiplicative inverse if and only if $\gcd(a, n) = 1$. This is why every nonzero element of $\mathbb{Z}_p$ (for prime p) has an inverse: any a with $1 \leq a \leq p - 1$ automatically satisfies $\gcd(a, p) = 1$, since p 's only divisors are 1 and p itself.

The classical **Euclidean Algorithm** computes $\gcd(a, b)$ by repeated division with remainder: $\gcd(a, b) = \gcd(b, a \bmod b)$, stopping when the remainder hits 0. The **Extended Euclidean Algorithm** does this same process but also keeps track of how the running remainder can be written as a combination $a \cdot x + b \cdot y$ (integers x, y) at every step — and when the algorithm finishes with $\gcd(a, b) = 1$, the final x is exactly $a^{-1} \bmod b$.

Example: Finding $3^{-1} \bmod 7$ Properly

We want x with $3x \equiv 1 \pmod{7}$. Run the Euclidean algorithm on $(7, 3)$:

$$7 = 2 \times 3 + 1$$

$$3 = 3 \times 1 + 0 \quad (\text{remainder hits 0, so } \gcd(7, 3) = 1)$$

Now work backward from the first line to express 1 as a combination of 7 and 3:

$$1 = 7 - 2 \times 3.$$

Reading this modulo 7: $1 \equiv -2 \times 3 \pmod{7}$, so $x = -2 \equiv 5 \pmod{7}$ is the inverse. **Check:** $3 \times 5 = 15 = 2 \times 7 + 1 \equiv 1 \pmod{7}$. Correct — and this matches whatever you found earlier by trial and error, but now via a method that scales to numbers with hundreds of digits.

Exercise: Practice

Use the (extended) Euclidean algorithm, not trial and error, to find $5^{-1} \bmod 11$. Show the division steps and the back-substitution step explicitly.

Looking Ahead: Why Cryptography Loves the Modulus

Modular arithmetic gives us two things cryptography needs badly: (1) a **finite** number system (so keys and computations have a fixed, bounded size, no matter how much you compute), and (2) built-in **one-wayness** in places — e.g., given only $a \bmod n$, you cannot tell which of infinitely many integers a actually was. The Learning-With-Errors (LWE) problem from the proposal document hides its secret inside equations that live in exactly this kind of finite, modular number system (there, the modulus is usually called q).

Exercise: Practice

(a) Compute $23 \times 15 \bmod 7$. (b) List all elements of $\mathbb{Z}_8$ that *do not* have a multiplicative inverse, and explain why (hint: compare each element to 8). (c) Confirm $\mathbb{Z}_7$ is a field by checking that 3 has a multiplicative inverse mod 7 (find x such that $3x \equiv 1 \pmod{7}$).

7 Vector Spaces, Basis, and Dimension

Definition: Vector Space

A **vector space** over a field F is a set V (whose elements are called **vectors**) equipped with vector addition and scalar multiplication (multiplying a vector by an element of F), satisfying the “obvious” rules you’d expect: addition makes $(V, +)$ an abelian group, and scalar multiplication distributes sensibly over both vector addition and field addition. The most important example for us: $\mathbb{R}^n$, the set of all n -tuples of real numbers, is a vector space over $\mathbb{R}$.

Definition: Linear Combination, Span, Linear Independence

Given vectors $v_1, \dots, v_k \in V$ and scalars $c_1, \dots, c_k \in F$, the vector $c_1v_1 + c_2v_2 + \dots + c_kv_k$ is called a **linear combination** of $v_1, \dots, v_k$. The set of *all* linear combinations of a set of vectors is called their **span**. The vectors $v_1, \dots, v_k$ are **linearly independent** if the only way to write $0 = c_1v_1 + \dots + c_kv_k$ is with every $c_i = 0$ — informally, no vector in the set is “redundant” (expressible using the others).

Definition: Basis and Dimension

A set of vectors $B = \{b_1, \dots, b_k\}$ is a **basis** of a vector space V if (1) B is linearly independent, and (2) B **spans** V (every vector in V can be written as a linear combination of B). Every basis of a given vector space has the same number of elements k , called the **dimension** of V .

Example: Basis of $\mathbb{R}^2$

The vectors $e_1 = (1, 0)$ and $e_2 = (0, 1)$ form the “standard” basis of $\mathbb{R}^2$: any vector (a, b) is exactly $a \cdot e_1 + b \cdot e_2$. But this is far from the *only* basis — $\{(1, 1), (1, -1)\}$ is also a valid basis of $\mathbb{R}^2$ (check: they’re linearly independent, and any (a, b) can be written as a combination of them). A vector space almost always has infinitely many possible bases; some are more convenient to work with than others. **Keep this fact in mind — it becomes very important in Section 8.**

7.1 Inner Products, Norms, and Orthogonality

To make “good basis” (short, nearly-perpendicular vectors) versus “bad basis” (long, nearly-parallel vectors) precise rather than just intuitive, we need a way to measure length and angle.

Definition: Inner Product and Norm

For $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbb{R}^n$, the (standard) **inner product** is $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$. The **norm** (length) of x is $\|x\| = \sqrt{\langle x, x \rangle}$. Two vectors are **orthogonal** if $\langle x, y \rangle = 0$ (the angle between them is 90°). A basis $\{b_1, \dots, b_k\}$ is an **orthogonal basis** if every pair b_i, b_j ($i \neq j$) is orthogonal.

Example: Computing Norms and Checking Orthogonality

For $x = (3, 4)$: $\|x\| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$. For $x = (1, 1)$ and $y = (1, -1)$: $\langle x, y \rangle = (1)(1) + (1)(-1) = 0$, so these two vectors (the alternate basis of $\mathbb{R}^2$ from the previous example) are orthogonal — a nicer basis, geometrically, than it might have looked at first glance.

Looking Ahead: Quantifying “Good” vs. “Bad” Bases

With norms in hand, we can finally give a real, measurable meaning to the “good basis / bad basis” idea flagged repeatedly in this lecture: a good basis has vectors with small norm $\|b_i\|$ that are close to orthogonal; a bad basis for the *same* lattice can have arbitrarily large norms and near-parallel vectors, despite generating an identical set of points. One common numerical measure of “how good” a basis is is its **orthogonality defect**, $\frac{\prod_i \|b_i\|}{\det(L)}$ (see Section 8.4 below for $\det(L)$) — it equals 1 for a perfectly orthogonal basis and grows the more skewed the basis is.

7.2 Gram–Schmidt Orthogonalization

Given *any* basis, can we systematically produce an orthogonal one (spanning the same space, though generally *not* the same lattice — more on that distinction below)? Yes:

Definition: Gram–Schmidt Process

Given linearly independent $b_1, \dots, b_k$, define $b_1^* = b_1$, and for $i = 2, \dots, k$:

$$b_i^* = b_i - \sum_{j=1}^{i-1} \mu_{i,j} b_j^*, \quad \text{where } \mu_{i,j} = \frac{\langle b_i, b_j^* \rangle}{\langle b_j^*, b_j^* \rangle}.$$

Each b_i^* is b_i with the “shadow” it casts on all previous b_j^* subtracted off, leaving only the genuinely new, orthogonal direction. The result $\{b_1^*, \dots, b_k^*\}$ is an orthogonal basis of the same vector space (though, importantly, its integer combinations generally do *not* produce the same lattice as the original b_i ’s — Gram–Schmidt vectors are a vector-space tool, not automatically a lattice basis).

Example: Gram–Schmidt on a 2×2 Example

Let $b_1 = (2, 0)$, $b_2 = (1, 2)$. Then $b_1^* = b_1 = (2, 0)$. Compute $\mu_{2,1} = \frac{\langle b_2, b_1^* \rangle}{\langle b_1^*, b_1^* \rangle} = \frac{(1)(2) + (2)(0)}{4} = \frac{2}{4} = 0.5$. So $b_2^* = b_2 - 0.5 b_1^* = (1, 2) - (1, 0) = (0, 2)$. Check: $\langle b_1^*, b_2^* \rangle = (2)(0) + (0)(2) = 0$ — orthogonal, as required.

Looking Ahead: Where Gram–Schmidt Leads

This process reappears, in a modified “integer-preserving” form, inside the **LLL algorithm** — the classic algorithm for turning a bad lattice basis into a reasonably good one, and a core tool used both to *attack* weak lattice-based schemes and to reason about the security of well-parametrized ones. We’ll meet it properly in a later lecture; for now, just remember that Gram–Schmidt is the geometric heart of it.

Exercise: Practice

Run Gram–Schmidt on $b_1 = (1, 1)$, $b_2 = (0, 2)$: compute b_1^* , $\mu_{2,1}$, and b_2^* , and verify $\langle b_1^*, b_2^* \rangle = 0$.

8 Lattices, Properly Defined

Everything so far has allowed *any* scalar from the field F (e.g., any real number) when forming linear combinations. Informally, a lattice is what happens when we impose one dramatic restriction: **only integer combinations are allowed**, which turns a solid, continuous vector space into a discrete grid of isolated points. That informal picture (Figure 3 below) is correct and worth keeping in your head. But the definition we will actually use — the one used in serious treatments of the subject, such as Chris Peikert’s graduate lecture notes *Lattices in Cryptography*¹ — starts from the **subgroup** idea from Section 3.1, not from a basis. We will show the two views agree.

8.1 Discrete Sets

Definition: Discrete Subset

A subset $S \subseteq \mathbb{R}^n$ is **discrete** if every point of S is *isolated*: for every $x \in S$, there exists some $\varepsilon > 0$ such that the open ball $B(x, \varepsilon)$ (all points within distance ε of x) contains *no* point of S other than x itself. Informally: you can always draw a small enough circle around any point of S that catches nothing else from S .

¹C. Peikert, *Lattices in Cryptography*, lecture notes, <https://github.com/cpeikert/LatticesInCryptography>.

Remark: A Shortcut for Subgroups

Checking every single point of a set for isolation sounds like a lot of work. But if H is a *subgroup* of $(\mathbb{R}^n, +)$ (Section 3.1), there's a shortcut: H is discrete if and only if the origin $\vec{0}$ alone is isolated, i.e., there is some $\varepsilon > 0$ with $B(\vec{0}, \varepsilon) \cap H = \{\vec{0}\}$. *Why this works:* suppose $\vec{0}$ is isolated with radius ε , and suppose some other point $y \in H$ were within ε of some $x \in H$. Since H is a subgroup, $y - x \in H$ as well, and $y - x$ is within ε of $\vec{0}$ — so by our assumption, $y - x = \vec{0}$, i.e., $y = x$. So checking one point (the origin) is enough for the whole subgroup, thanks to closure under subtraction.

8.2 The Definition

Definition: Lattice

A **lattice** is a discrete additive subgroup L of $\mathbb{R}^n$. That is, $L \subseteq \mathbb{R}^n$ is a lattice if:

- (L1) L is a subgroup of $(\mathbb{R}^n, +)$ (Section 3.1): it contains $\vec{0}$, and is closed under addition and negation.
- (L2) L is discrete: by the shortcut above, there exists $\varepsilon > 0$ such that $B(\vec{0}, \varepsilon) \cap L = \{\vec{0}\}$.

Example: $\mathbb{Q}$ Is a Subgroup, But *Not* a Lattice

Consider $\mathbb{Q} \subseteq \mathbb{R}$ (the rational numbers) under addition. This *is* a subgroup: it contains 0, the sum of two rationals is rational, and $-a$ is rational whenever a is. So condition (L1) holds. But condition (L2) fails badly: the rationals are **dense** in $\mathbb{R}$, meaning every open ball around 0, no matter how tiny, contains infinitely many other rational numbers. There is no $\varepsilon > 0$ that isolates 0. So $\mathbb{Q}$ is a perfectly good subgroup that is **not** a lattice — this is exactly why condition (L2) has to be part of the definition; subgroup alone is not a strong enough requirement.




Figure 2: Discreteness, visualized. Left: in $\mathbb{Z}$, every point has a gap of size 1 around it with nothing else nearby — discrete. Right: in $\mathbb{Q}$, no matter how small a circle you draw around a point, there are always infinitely many more rational points crammed inside it — dense, and therefore *not* discrete.

Example: $\mathbb{Z}$ and $\mathbb{Z}^n$ Are Lattices

$\mathbb{Z} \subseteq \mathbb{R}$ is a subgroup (trivially — and recall from Section 3.1 that $n\mathbb{Z}$ is a subgroup of $\mathbb{Z}$ for any n ; here we're just looking at $\mathbb{Z}$ itself sitting inside $\mathbb{R}$), and it is discrete: taking $\varepsilon = 0.5$, $B(0, 0.5) \cap \mathbb{Z} = \{0\}$, since the nearest other integers (± 1) are a full unit away. So $\mathbb{Z}$ is a lattice in $\mathbb{R}$ — the simplest possible example. The same argument, coordinate by coordinate, shows $\mathbb{Z}^n \subseteq \mathbb{R}^n$ (all integer-coordinate points) is a lattice in $\mathbb{R}^n$; it is often called **the integer lattice**.

8.3 Recovering the Basis Picture

The subgroup definition is more general and more standard, but the “grid generated by a basis” picture you may already have in your head is not wrong — it is a theorem, not a definition, and it's worth seeing why.

Basis Representation of Lattices Let $L \subseteq \mathbb{R}^n$. Then L is a lattice (in the sense of the subgroup definition above, i.e., a discrete additive subgroup) **if and only if** there exist linearly independent vectors

$b_1, \dots, b_k \in \mathbb{R}^n$ (with $k \leq n$) such that

$$L = \mathcal{L}(b_1, \dots, b_k) = \left\{ \sum_{i=1}^k a_i b_i \mid a_i \in \mathbb{Z} \right\}.$$

The vectors $b_1, \dots, b_k$ are called a **basis** of L ; the number k is the **rank** of L . If $k = n$ (the basis spans all of $\mathbb{R}^n$), L is called **full-rank** — this is the common case in cryptography.

Remark: Proof Idea (Sketch Only)

Easy direction ($\Leftarrow$): given linearly independent $b_1, \dots, b_k$, the set of integer combinations is clearly a subgroup (adding two integer combinations, or negating one, gives another integer combination). Discreteness holds because the b_i point in genuinely independent directions: you cannot make $\sum a_i b_i$ arbitrarily close to $\vec{0}$ without every a_i being exactly 0 — linear independence rules out any “near-cancellation.” (Making this fully rigorous uses the fact that the linear map $(a_1, \dots, a_k) \mapsto \sum a_i b_i$ has a bounded inverse on its image.)

Harder direction ($\Rightarrow$): given only that L is a discrete subgroup, we must *construct* a basis. The idea is greedy and inductive: pick b_1 to be a shortest nonzero vector in L (this exists precisely *because* L is discrete — in a non-discrete set like $\mathbb{Q}$, there is no shortest nonzero element, since you can always find a smaller one). Then look at the part of L not lying on the line through b_1 , and pick b_2 to be a shortest vector there, and so on. A careful argument (using discreteness at each step, and projecting onto the space orthogonal to the vectors already chosen) shows this process terminates with a valid basis after $k \leq n$ steps. The full details are carried out rigorously in Peikert’s notes; we take the result as given here.

Looking Ahead: A Name for the “Shortest Vector”

The shortest nonzero vector used in the proof sketch above has a standard name: $\lambda_1(L)$, the **minimum distance** of the lattice. Computing $\lambda_1(L)$ — or even just estimating it — for a lattice given only a (possibly bad) basis is called the **Shortest Vector Problem (SVP)**, and it is believed to be extremely hard in high dimensions. This hardness is one of the two or three core hard problems that lattice-based cryptography’s security ultimately rests on. We’ll return to this once we reach the LWE problem.




Figure 3: A 2-dimensional, full-rank lattice generated by a basis $\{b_1, b_2\}$: every grid dot is reachable by some *integer* combination $a_1 b_1 + a_2 b_2$ (with $a_1, a_2 \in \mathbb{Z}$), starting from any chosen origin dot. This is the same object as the subgroup definition above — just viewed through its basis instead of as an abstract subgroup.

Remark: Same Lattice, Different Basis

A single lattice has *many* different valid bases (just as a vector space does — recall the “Basis of $\mathbb{R}^2$ ” example in Section 7). Two bases generate the exact same set of points if and only if one can be converted to the other using an integer matrix with determinant ± 1 (a **unimodular** transformation) — we won’t need the details of this fact yet, but the consequence is critical: some bases consist of short, nearly-perpendicular vectors (a “good” basis, easy to compute with), while other, equally valid bases for the *same* lattice consist of long, nearly-parallel vectors (a “bad” basis, hard to compute with). **This gap between good and bad bases for the same lattice is the single most important idea in lattice-based cryptography:** the secret key is typically a good (short) basis, while the public key describes the same lattice using a bad (long, scrambled-looking) basis. Recovering the good basis from the bad one is believed to be extremely hard once the dimension n is large enough.

Example: A Tiny Lattice by Hand

Let $b_1 = (2, 0)$ and $b_2 = (0, 3)$ in $\mathbb{R}^2$. The lattice $\mathcal{L}(b_1, b_2)$ consists of every point $(2a_1, 3a_2)$ for integers a_1, a_2 — so it includes $(0, 0)$, $(2, 0)$, $(-2, 0)$, $(0, 3)$, $(2, 3)$, $(4, -3)$, and so on, but *not* $(1, 1)$ or $(1, 0)$, since those cannot be written as $(2a_1, 3a_2)$ for integer a_1, a_2 . As a sanity check against the subgroup definition directly: is this set a subgroup? Yes — closed under addition and negation, contains $(0, 0)$. Is it discrete? Yes — the nearest point to the origin is at distance 2, so any $\varepsilon < 2$ isolates $\vec{0}$.

8.4 The Determinant (Volume) of a Lattice

One more piece of structure will matter a great deal in later lectures: every lattice has a well-defined “volume,” independent of which basis you use to describe it.

Definition: Fundamental Domain and Determinant

Given a basis $b_1, \dots, b_n$ of a full-rank lattice $L \subseteq \mathbb{R}^n$, the **fundamental parallelepiped** is

$$\mathcal{P}(B) = \left\{ \sum_{i=1}^n t_i b_i \mid 0 \leq t_i < 1 \right\},$$

i.e., the region “tiled out” by taking all *fractional* (not just integer) combinations between 0 and 1. Copies of $\mathcal{P}(B)$, shifted by every lattice point, tile all of $\mathbb{R}^n$ with no gaps or overlaps. The **determinant** of the lattice, $\det(L)$, is the volume of $\mathcal{P}(B)$, computed as $\det(L) = |\det(B)|$, where B is the matrix whose rows (or columns) are $b_1, \dots, b_n$.

Remark: Why $\det(L)$ Doesn’t Depend on Which Basis You Pick

Recall from the “Same Lattice, Different Basis” remark above: any two bases of the same lattice are related by a unimodular matrix U (integer entries, $\det(U) = \pm 1$), i.e., $B' = UB$. Then $\det(B') = \det(U) \det(B) = \pm \det(B)$, so $|\det(B')| = |\det(B)|$. This confirms $\det(L)$ is a genuine property of the lattice itself, not an artifact of which basis happened to be used to compute it — exactly the kind of basis-independence you should expect from something called “the volume of the lattice.”

Example: Computing $\det(L)$ for the Tiny Lattice

For $b_1 = (2, 0)$, $b_2 = (0, 3)$ from the example above, $B = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$, so $\det(L) = |\det(B)| = |2 \times 3 - 0 \times 0| = 6$. Geometrically: the fundamental parallelepiped is just the 2×3 rectangle $[0, 2) \times [0, 3)$, and copies of this rectangle exactly tile the plane, anchored at every lattice point — matching the picture in Figure 3’s style, just for this specific (axis-aligned) lattice.

Looking Ahead: Minkowski’s Theorem: Connecting Volume Back to $\lambda_1(L)$

Here is a genuinely deep fact we will only preview, not prove: **Minkowski’s First Theorem** says that any symmetric convex region of volume greater than $2^n \det(L)$ must contain a nonzero lattice point. Applying this to a ball gives a bound relating the two quantities we’ve now defined: $\lambda_1(L) \leq \sqrt{n} \cdot \det(L)^{1/n}$ (up to constants). In plain terms: a lattice cannot “hide” an arbitrarily long shortest vector if its volume is small — volume and minimum distance are locked together by the geometry, no matter which basis you’re handed. This is part of *why* SVP is believed hard only in the “regime” cryptographers carefully choose (high dimension n , appropriately scaled $\det(L)$) rather than for arbitrary lattices — a preview of parameter selection, which we’ll return to once we discuss how Kyber and Dilithium choose their concrete dimensions and moduli.

Exercise: Practice

For the lattice generated by $b_1 = (1, 2)$, $b_2 = (3, 1)$ (from the earlier homework exercise): compute $\det(L)$ directly from the basis matrix. Then pick any *other* valid basis for the same lattice (e.g., $b'_1 = b_1$, $b'_2 = b_2 - b_1$) and confirm you get the same $\det(L)$.

Looking Ahead: Where This Leads Next Lecture

With Groups $\rightarrow$ Subgroups $\rightarrow$ Rings $\rightarrow$ Fields $\rightarrow$ Modular arithmetic $\rightarrow$ Vector spaces $\rightarrow$ Basis $\rightarrow$ Lattice now in place, the next lecture can finally state the **Learning With Errors (LWE)** problem precisely: a secret vector, hidden inside noisy linear equations over $\mathbb{Z}_q$ (a ring, Section 6), which is exactly the kind of “bad basis” hardness described above, dressed up in a specific, standardized cryptographic form. We will also unpack the ring $R_q = \mathbb{Z}_q[x]/(x^n + 1)$ flagged in Section 4, since **module lattices** (lattices whose coordinates come from a ring like R_q instead of plain integers) are exactly what Kyber and Dilithium use in practice.

9 Summary Table

Structure	Requires	Example
Group	1 operation, inverses exist	$(\mathbb{Z}, +)$
Subgroup	A subset that is itself a group	$2\mathbb{Z} \subseteq \mathbb{Z}$
Coset / Quotient group	G sorted into shifted copies of H	$\mathbb{Z}/3\mathbb{Z} = \{0, 1, 2\}$
Ring	2 operations, no division needed	$(\mathbb{Z}, +, \times), \mathbb{Z}[x]$
Integral domain	Ring, no zero divisors	$\mathbb{Z}, \mathbb{Z}_5$ (not $\mathbb{Z}_6$)
Field	Ring + division by nonzero elements	$\mathbb{Q}, \mathbb{R}, \mathbb{Z}_p$ (p prime)
Modulus	Wraparound arithmetic	$\mathbb{Z}_n = \{0, \dots, n-1\}$
Vector space	Field + vectors + scalar mult.	$\mathbb{R}^n$
Basis	Independent, spanning vector set	$\{(1, 0), (0, 1)\}$ in $\mathbb{R}^2$
Orthogonal basis	Basis, all pairs perpendicular	Gram–Schmidt output
Lattice	Discrete additive subgroup of $\mathbb{R}^n$	$\mathbb{Z}^n$; grid of points
$\det(L)$	Volume of the fundamental domain	$ \det(B) $, any basis B

10 Full List of Exercises (Assign as Homework Before Lecture 2)

Exercise: Homework Set

1. Verify $(\{1, 2, 3, 4\}, \times \bmod 5)$ is a group by writing out its full multiplication table (Section 3 exercise, above).
2. Is $3\mathbb{Z} = \{\dots, -6, -3, 0, 3, 6, \dots\}$ a subgroup of $(\mathbb{Z}, +)$? Justify using the subgroup test (Section 3.1). Is $\{0, 3, 6, 9, \dots\}$ (only the non-negative multiples of 3) a subgroup of $(\mathbb{Z}, +)$? Why or why not?
3. List all distinct cosets of $4\mathbb{Z}$ in $\mathbb{Z}$, confirm there are exactly 4, and identify which coset -5 belongs to (Section 3.2 exercise, above).
4. Find a pair of zero divisors in $\mathbb{Z}_{12}$, and use the zero-divisor fact (Section 4.1) to explain why $\mathbb{Z}_{12}$ cannot be a field, without computing any inverses.
5. Compute $23 \times 15 \bmod 7$; list the non-invertible elements of $\mathbb{Z}_8$; use the Extended Euclidean Algorithm (Section 6.1) to find the inverse of $3 \bmod 7$ — confirm it matches the worked example.
6. Use the Extended Euclidean Algorithm to find $5^{-1} \bmod 11$ (Section 6.1 exercise, above), showing all steps.
7. Is $\mathbb{Z}_9$ a field? Why or why not? Answer it two ways: (a) using gcd (Section 6), and (b) using zero divisors (Section 4.1).
8. Show that $\{(1, 1), (1, -1)\}$ is a valid basis of $\mathbb{R}^2$ by (a) checking linear independence and (b) writing the vector $(4, 2)$ as a linear combination of them.
9. Run Gram–Schmidt on $b_1 = (1, 1)$, $b_2 = (0, 2)$ (Section 7.2 exercise, above), and verify the result is orthogonal.
10. For the lattice generated by $b_1 = (1, 2)$ and $b_2 = (3, 1)$: (a) is the point $(7, 5)$ in the lattice? (b) compute $\det(L)$ from this basis; (c) pick a different valid basis for the same lattice and confirm $\det(L)$ is unchanged (Section 8.4 exercise, above).
11. Let $L = \{(x, y) \in \mathbb{Z}^2 : x + y \text{ is even}\}$. (a) Show L is a subgroup of $(\mathbb{R}^2, +)$. (b) Show L is discrete (find an ε that isolates $\vec{0}$). (c) Since L is therefore a lattice, find a basis $\{b_1, b_2\}$ for it, and compute $\det(L)$.
12. In your own words (3–4 sentences): why does the definition of a lattice need *both* “subgroup” and “discrete”? Use the $\mathbb{Q}$ example from Section 8.2 to explain what goes wrong if you drop the discreteness requirement.
13. In your own words (3–4 sentences), explain why a “good basis” and a “bad basis” for the same lattice can make the same mathematical object feel completely different to work with computationally, now that you can measure this with norms (Section 7.1).