1 Problem

Compute the derivatives of the least squares loss $L[\phi]$ with respect to ϕ_0 and ϕ_1 for the Gabor model equation 6.8.

2 Answer

$$f[x,\phi] = sin(\phi_0 + 0.06 \cdot \phi_1 x) \cdot exp\left(\frac{-(\phi_0 + 0.06 \cdot \phi_1 x)^2}{32.0}\right) \tag{1}$$

$$L[\phi] = \sum_{i=1}^{I}(f[x_i,\phi] - y_i)^2 \tag{2}$$

$$\tag{3}$$

$$\begin{aligned}
\text{Let } u_i &= f[x_i,\phi] - y_i \\
\text{Then } L[\phi] &= \sum_{i=1}^{I}(u_i)^2 \\
\frac{\partial L[\phi]}{\partial \phi_0} &= \sum_{i=1}^{I}\frac{\partial L[\phi]}{\partial u_i} \cdot \frac{\partial u_i}{\partial \phi_0} \\
\text{Let us break it down } \frac{\partial L[\phi]}{\partial u_i} &= 2 \cdot u_i = 2 \cdot \left(sin(\phi_0 + 0.06 \cdot \phi_1 x) \cdot exp\left(\frac{-(\phi_0 + 0.06 \cdot \phi_1 x_i)^2}{32.0}\right) - y_i\right) \\
\frac{\partial u_i}{\partial \phi_0} &= \frac{\partial sin(\phi_0 + 0.06 \cdot \phi_1 x) \cdot exp\left(\frac{-(\phi_0+0.06\cdot\phi_1 x_i)^2}{32.0}\right)}{\partial \phi_0} \\
\text{Let } p &= sin(\phi_0 + 0.06 \cdot \phi_1 x_i) \\
\text{Let } q &= exp\left(\frac{-(\phi_0 + 0.06 \cdot \phi_1 x_i)^2}{32.0}\right) \\
\text{So that } \frac{\partial u_i}{\partial \phi_0} &= p\frac{\partial q}{\partial \phi_0} + q\frac{\partial p}{\partial \phi_0} \\
\text{Let } r &= \phi_0 + 0.06 \cdot \phi_1 x_i \\
\text{So that } \frac{\partial p}{\partial \phi_0} &= \frac{\partial sin(r)}{\partial r} \cdot \frac{\partial r}{\partial \phi_0} = cos(r) \cdot (1) = cos(\phi_0 + 0.06 \cdot \phi_1 x_i) \\
\text{Let } s &= \left(\frac{-(\phi_0 + 0.06 \cdot \phi_1 x_i)^2}{32.0}\right)
\end{aligned}$$