

DIRECTORY OF SERVICES



NABL
CERTIFICATE
No.MC-3108

Laboratory Accredited to
National Accreditation Board for
Testing & Calibration Laboratories

We believe ...

Quality is never an accident; it is always the result of high intention, sincere effort, intelligent direction and skillful execution; represents the wise choice of many alternatives.



Chinthavalappu Junction, Jail Road, Calicut-673004, Kerala
Web: www.aswinicalicut.net, E-mail: mail@aswinicalicut.net
Phone: 0495 2338888, 7990200300

profile

Aswini Diagnostic Services, made it's humble beginning in 1994. Since then we have been treading the path of steady growth and development, we are now amongst the topmost diagnostic centres of the region and continue to provide most modern diagnostic facilities of superior quality to the patients and doctors of Calicut and neighbouring areas of North Kerala.

Aswini added a feather to its cap in 2010, by being the first lab based at North Malabar, to be accredited to NABL, the board constituted by Government of India to assure quality and reliability in the field of laboratory science.

Thirty years of glorious service, a host of satisfied customers, outstanding excellence in all high performance parameters. All have made Aswini the most preferred diagnostic centre in North Kerala

Why Aswini ?

Behind Aswini's accomplishment, are a lot of remarkable factors that paved the way for its success.

Focus on quality

Total automation of lab tests

Wider selection of clinically reliable reports

Accurate result in the shortest possible time

Extraordinary levels of technical support and customer care

Economical rates

We Insist

On quality results

On proper quality checking

On minimum cost to the patients

At the shortest possible time

We Have

NABL Accreditation

ISO 15189 accredited laboratory

Highly trained professionals managing testing

Equipments of international quality

Quality control checking by All India Institute of Medical Science, CMC-Vellore, Bio Rad, IfQ-Lübeck, Randox and RML-QAP leaders in quality testing.

24*7 service and home collection facility

Copyright © 2024 by Aswini Diagnostic Services.

No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the publisher.

The test menu is subject to revision. Please contact our customer service for updated details.



Dear Esteemed Patients and Team Members,

As we embark on 32 years of service and commitment to healthcare excellence, I find great satisfaction in reflecting on our journey and looking ahead to the promising future that lies before us. For an institution 30 years is not a big period, but as an individual who had been part of it from its inception, it gives me intense pleasure and satisfaction for what ever I could do in its growth from the very small beginning since 1994 to the present.

There had been moments of joy and periods of trials and tribulations and I am thankful to the almighty for allowing me to take everything with equanimity.

At Aswini we are dedicated to upholding the highest standards of quality, precision, and ethics in diagnostic services. Our mission is to consistently deliver accurate and reliable results, and compassionate diagnostic services that contribute to the well-being of our customers. We understand the pivotal role diagnostic testing plays in healthcare decision- making, and our dedicated team of professionals is committed to upholding the highest standards of quality and accuracy.

As we move forward, we remain focused on enhancing the patient experience. Your health and satisfaction are at the heart of everything we do. Our team is dedicated to providing a seamless and comfortable journey through the diagnostic process, ensuring that you receive the information and support you need to make informed decisions about your health.

As we face new challenges and opportunities, I am confident that the spirit of teamwork and commitment that defines our lab will propel us to even greater heights.

In closing, I invite you to join us in this exciting journey of progress and service. Whether you are a customer entrusting us with your health, or a team member contributing to our success, your role is invaluable. Thank you for being an integral part of the Aswini Diagnostic Services family. Together, let us continue to make a positive impact on healthcare and contribute to a healthier, happier community.

Wishing you good health and prosperity in the coming years.

Warm regards.



Dr. P.C.Rajalakshmi, MD (PATH)
Medical Director, Senior Consultant Pathologist,
Aswini Diagnostic Services A unit of ADS research centre LLP
Former Asso.Prof.Govt.MCH

Certifications



National Accreditation Board for
Testing and Calibration Laboratories

CERTIFICATE OF ACCREDITATION

**ASWINI DIAGNOSTIC SERVICES (A UNIT OF ADS
RESEARCH CENTRE LLP)**

has been assessed and accredited in accordance with the standard

ISO 15189:2022

**"Medical laboratories - Requirements for quality and
competence"**

for its facilities at

CHINTHAVALAPPU JUNCTION, JAIL ROAD, CALICUT, KOZHIKODE, KERALA, INDIA

In the field of

Medical Testing

Certificate Number: MC-3108

Issue Date: 12/03/2025

Valid Until: 17/01/2026

This certificate supersedes the Certificate No. MC-3108 with issue date 17/02/2024 in view of change in legal entity of the laboratory.

This certificate supersedes the Certificate No. MC-3108 with issue date 18/01/2024 in view of the transition of Accreditation Standard from ISO 15189: 2012 to ISO 15189: 2022.

This certificate remains valid for the Scope of Accreditation as specified in the annexure subject to continued satisfactory compliance to the above standard & the relevant requirements of NABL.
(To see the scope of accreditation of this laboratory, you may also visit NABL website www.nabl-india.org)

Name of Legal Entity: ADS RESEARCH CENTRE LLP

Signed for and on behalf of NABL




Pankaj Johri
Director


N. Venkateswaran
Chief Executive Officer



PROFICIENCY TESTING REPORT
ISHTM-AIIMS EXTERNAL QUALITY ASSURANCE PROGRAMME
NABL accredited program as per ISO/IEC 17043:2023 standard
Organized By Department of Hematology, AIIMS, New Delhi-110029



PARTICIPATION CERTIFICATE

[Certificate No. EQAP/37/2024/38]

Date 30.12.2024

This is to certify that **"ASWINI DIAGNOSTIC SERVICES(A UNIT OF ADS RESERCH CENTRE LLP), KOZHIKODE, KERALA, 673004"** has participated in the "ISHTM-AIIMS External Quality Assurance Program" for the period "January 2024 to December 2024".

Report authorized by,

Dr. Manoranjan Mahapatra (Prof. & Head)
PT Co-ordinator: ISHTM-AIIMS-EQAP
Department of Hematology, AIIMS, New Delhi



CMC External Quality Assessment Scheme
Christian Medical College Vellore

TRANSFUSION MEDICINE MODULE

Certificate of Participation

This is to certify that the following laboratory has participated in the External Quality Assessment Scheme conducted by the Christian Medical College Vellore

RIN Number : T2061

Name of Laboratory / Blood Bank : Aswini Diagnostic Services(A UNIT OF ADS RESERCH CENTRE LLP)

Year : 2024

Parameters	Results returned
Immunohaematology	
Blood Grouping and Typing	3/3 Success
Direct Coombs and Indirect Coombs test	3/3 Success

Issued under authorization of

Professor Anil Srinivasan, MD, FRACP, FRCPA, FRCP

Chairperson

Department of Transfusion Medicine & Immunohaematology
Christian Medical College Vellore 632004, Tamil Nadu, India

Please keep this certificate in a safe place as evidence of your participation

Certificate No: Transfusion / 2024 / T2061





INDIAN ACADEMY of CYTOLOGISTS

National External Quality Assurance Programme

Certificate of Participation

This certificate is to certify that

Aswini Diagnostic Services (A Unit of ADS research centre LLP), Calicut

with EQA Registration No. **142/25**

has successfully participated in the

National External Quality Assurance Programme
of the India Academy of Cytologists for the year 2025.

Dr Meherbano Kamal
Chairperson
Accreditation and Examinations
Committee

Dr. Venkateswaran K. Iyer
President, IAC

Dr. Bharat Rekhi
Secretary, IAC

Dr. Nalini Gupta
Chairperson
EQA Committee





Awards and recognitions








TABLE OF CONTENTS

DISCIPLINES	PAGE NO
INDEX PAGE	I
PROFILE	II
MEDICAL DIRECTORS MESSAGE	IV
CERTIFICATIONS	V
AWARDS AND RECOGNITIONS	XIII
TABLE OF CONTENTS	XVI
DIRECTORY OF SERVICES - ALL DISCIPLINES	
CLINICAL BIOCHEMISTRY	1-31
CLINICAL PATHOLOGY	32-39
HEMATOLOGY	40-46
HISTOPATHOLOGY AND CYTOPATHOLOGY	47-66
MICROBIOLOGY	67-79
MOLECULAR BIOLOGY	80-82
SEROLOGY	83-90



CLINICAL BIOCHEMISTRY



DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
1	820	ACE ANGIOTENSIN CONVERTING ENZYME	FAPGG	SERUM	SERUM CONTAINER	3 Hours	ACE IS AN ENZYME THAT PLAYS A CRITICAL ROLE IN THE RENIN-ANGIOTENSIN-ALDOSTERONE SYSTEM (RAAS), WHICH REGULATES BLOOD PRESSURE, FLUID BALANCE, AND ELECTROLYTE BALANCE IN THE BODY. THE TEST IS PRIMARILY USED TO DIAGNOSE AND MONITOR CERTAIN MEDICAL CONDITIONS LIKE SARCOIDOSIS, CHRONIC OBSTRUCTIVE PULMONARY DISORDER(COPD), HYPOTHYROIDISM, CYSTIC FIBROSIS, EMPHYSEMA, ETC.
2	10	ACID PHOSPHATASE	ALPHA NAPHTHYL P04	SERUM	SERUM CONTAINER	3 Hours	TOTAL ACID PHOSPHATASE ACTIVITY IS USEFUL IN THE EVALUATION OF PROSTATE CANCER, PAGET'S DISEASE, HYPERPARATHYROIDISM WITH SKELETAL INVOLVEMENT, AND GAUCHER'S DISEASE.
3	1638	ACTH (ADRENO-CORTICO TROPIC HORMONE)	ECLIA	EDTA PLASMA	EDTA CONTAINER	3 Hours	THE ACTH TEST IS OFTEN USED TO EVALUATE THE FUNCTIONING OF ADRENAL GLANDS. ACTH IS A HORMONE PRODUCED BY THE PITUITARY GLAND IN BRAIN, AND IT PLAYS A CRUCIAL ROLE IN REGULATING THE PRODUCTION AND RELEASE OF CORTISOL, A STEROID HORMONE PRODUCED BY THE ADRENAL GLANDS.
4	70	ADENOSINE DEAMINASE (CSF)	KINETIC METHOD	CSF	AIR TIGHT CONTAINER	3 Hours	ADENOSINE DEAMINASE (ADA) IN CEREBROSPINAL FLUID (CSF) IS CONSIDERED TO BE A USEFUL BIOMARKER IN DIFFERENTIATING TUBERCULOUS MENINGITIS (TBM) FROM OTHER MENINGITIS IN NON-HIV PATIENTS.
5	739	ADENOSINE DEAMINASE (FLUID)	KINETIC METHOD	FLUID	AIR TIGHT CONTAINER	3 Hours	ADENOSINE DEAMINASE IS AN ENZYME INVOLVED IN PURINE METABOLISM, AND ITS LEVEL CAN BE MEASURED IN VARIOUS BODY FLUIDS SUCH AS CSF, PLEURAL FLUID, AND PERITONEAL FLUID. THE TEST IS PARTICULARLY RELEVANT IN THE DIAGNOSIS AND MONITORING OF MEDICAL CONDITIONS LIKE PLEURAL TB, THAT AFFECTS THE LINING OF THE LUNGS, IN CASES OF SUSPECTED CENTRAL NERVOUS SYSTEM INFECTIONS, SUCH AS TUBERCULOUS MENINGITIS, ETC.
6	59	ADENOSINE DEAMINASE	KINETIC METHOD	SERUM	SERUM CONTAINER	3 Hours	THE ADA BLOOD TEST IS USED PRIMARILY FOR ASSESSING IMMUNE SYSTEM FUNCTION, PARTICULARLY IN THE CONTEXT OF CERTAIN MEDICAL CONDITIONS, SUCH AS: SEVERE COMBINED IMMUNODEFICIENCY (SCID), TUBERCULOSIS (TB) ETC.
7	11	ALBUMIN	BROMOCRESOL GREEN	SERUM	SERUM CONTAINER	2 Hours	THIS TEST MEASURES THE AMOUNT OF ALBUMIN IN BLOOD. LOW ALBUMIN LEVELS CAN BE A SIGN OF LIVER OR KIDNEY DISEASE OR ANOTHER MEDICAL CONDITION. HIGH LEVELS MAY BE A SIGN OF DEHYDRATION.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
8	1	ALBUMIN (ASCITIC FLUID)	BROMOCRESOL GREEN	FLUID	AIR TIGHT CONTAINER	2 Hours	THE ASCITIC ALBUMIN TEST IS THE MOST ACCURATE WAY TO DIFFERENTIATE DIFFERENT CAUSES OF ASCITES. THE ALBUMIN GRADIENT, WHICH IS THE DIFFERENCE BETWEEN THE SERUM ALBUMIN AND THE ASCITIC FLUID ALBUMIN, IS THE BEST WAY TO DIFFERENTIATE BETWEEN LIVER CAUSES OF ASCITES AND PERITONEAL CAUSES OF ASCITES.
9	13	ALKALINE PHOSPHATASE	PARA- NITROPHENYL PHOSPHATE	SERUM	SERUM CONTAINER	2 Hours	AALKALINE PHOSPHATASE (ALP) IS AN ENZYME THAT HELPS BREAK DOWN PROTEINS AND REMOVE PHOSPHATE GROUPS FROM MOLECULES IN THE BODY, WITH SIGNIFICANT AMOUNTS PRODUCED IN THE LIVER AND BONES, AND ALSO FOUND IN THE INTESTINES, KIDNEYS, AND PLACENTA DURING PREGNANCY. A BLOOD TEST MEASURES ALP LEVELS TO SCREEN FOR LIVER DISEASE, BONE DISORDERS, AND OTHER CONDITIONS, AS ELEVATED LEVELS CAN INDICATE PROBLEMS WITH THESE ORGANS OR SPECIFIC CANCERS.
10	1617	ALKALINE PHOSPHATASE (CSF)	PARA- NITROPHENYL PHOSPHATE	CSF	AIR TIGHT CONTAINER	2 Hours	ALP IS AN ENZYME THAT IS PRESENT IN VARIOUS TISSUES THROUGHOUT THE BODY, INCLUDING THE CENTRAL NERVOUS SYSTEM. ELEVATED LEVELS OF ALP IN THE CSF CAN INDICATE CERTAIN MEDICAL CONDITIONS LIKE BRAIN TUMOR, OTHER SPACE-OCCUPYING LESIONS, BACTERIAL OR VIRAL INFECTIONS.
11	1174	ALLERGY PANEL, FOOD SCREENING (64) Allergens	IMMUNOBLOT	SERUM	SERUM CONTAINER	24 Hours	A DIAGNOSTIC TEST USED TO IDENTIFY SPECIFIC FOOD ALLERGENS THAT MAY TRIGGER ALLERGIC REACTIONS IN INDIVIDUALS. ALLERGY PANELS HELP TO IDENTIFY SPECIFIC FOODS RESPONSIBLE FOR CAUSING ALLERGIC REACTIONS SO THAT APPROPRIATE AVOIDANCE STRATEGIES CAN BE IMPLEMENTED.
12	1746	ALLERGY PANEL, INHALATION SCREENING	IMMUNOBLOT	SERUM	SERUM CONTAINER	24 Hours	A DIAGNOSTIC TEST USED TO IDENTIFY SPECIFIC AIR BORNE ALLERGENS THAT MAY TRIGGER ALLERGIC REACTIONS IN INDIVIDUALS. THIS PANEL HELP TO IDENTIFY SPECIFIC INHALANTS RESPONSIBLE FOR CAUSING ALLERGIC REACTIONS, SO THAT APPROPRIATE AVOIDANCE STRATEGIES CAN BE IMPLEMENTED.
13	1747	ALLERGY, PAEDIATRIC FOOD AND INHALATION SCREENING PANEL	IMMUNOBLOT	SERUM	SERUM CONTAINER	24 Hours	A DIAGNOSTIC TEST USED TO IDENTIFY SPECIFIC FOOD AND AIR BORNE ALLERGENS THAT MAY TRIGGER ALLERGIC REACTIONS IN CHILDREN. THIS PANEL HELP TO IDENTIFY SPECIFIC FOOD AND INHALANTS RESPONSIBLE FOR CAUSING ALLERGIC REACTIONS, SO THAT APPROPRIATE AVOIDANCE STRATEGIES CAN BE IMPLEMENTED.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
14	108	ALPHA FETO PROTEIN (AFP)	TWO STEP IMMUNO ASSAY CMIA	SERUM	SERUM CONTAINER	2 Hours	ALPHA-FETOPROTEIN IS A PROTEIN THAT IS NORMALLY PRODUCED BY THE LIVER AND YOLK SAC OF A DEVELOPING FETUS. AFP LEVELS CAN BE MEASURED DURING PREGNANCY TO SCREEN FOR CERTAIN BIRTH DEFECTS AND CHROMOSOMAL ABNORMALITIES, SUCH AS NEURAL TUBE DEFECTS AND DOWN SYNDROME. AFP LEVELS CAN BE MONITORED IN PATIENTS WHO HAVE A HISTORY OF CERTAIN CONDITIONS, SUCH AS LIVER CANCER OR TESTICULAR CANCER, TO TRACK THE EFFECTIVENESS OF TREATMENT AND DETECT ANY POTENTIAL RECURRENCE.
15	1174	AFP CSF	TWO STEP IMMUNO ASSAY CMIA	CSF	AIR TIGHT CONTAINER	2 Hours	AFP TEST CAN BE USED TO DETECT NEURAL TUBE DEFECTS IN THE DEVELOPING FETUS. ELEVATED LEVELS OF AFP IN THE CSF MIGHT INDICATE CONDITIONS LIKE SPINA BIFIDA OR ANENCEPHALY. ELEVATED CSF AFP LEVELS ARE ASSOCIATED WITH CERTAIN GERM CELL TUMORS, PARTICULARLY THOSE AFFECTING THE CENTRAL NERVOUS SYSTEM, TUMORS OF THE PINEAL GLAND ETC.
16	1413	AMMONIA	ENZYMATIC GLDH METHOD	EDTA PLASMA	EDTA CONTAINER	3 Hours	AMMONIA A NORMAL WASTE PRODUCT OF THE BODY GETS CONVERTED TO UREA BY THE LIVER.ELEVATED LEVELS OF AMMONIA ARE SEEN IN VARIOUS DISEASES LIKE LIVER CIRRHOSIS, HEPATIC ENCEPHALOPATHY, SEVERE GASTROINTESTINAL BLEEDING AND METABOLIC DISEASES LIKE UREA CYCLE DISORDERS.
17	16	AMYLASE	CNPG3	SERUM	SERUM CONTAINER	2 Hours	ELEVATED AMYLASE LEVELS CAN INDICATE PANCREATITIS, WHICH IS THE INFLAMMATION OF THE PANCREAS. TRAUMA OR INJURY TO THE PANCREAS, GASTRO INTESTINAL PERFORATIONS, ETC, CAN ALSO CAUSE ELEVATED FLUID AMYLASE LEVELS IN THE ABDOMINAL OR PLEURAL FLUID.
18	108	AMYLASE (FLUID)	CNPG3	FLUID	AIR TIGHT CONTAINER	2 Hours	ELEVATED FLUID AMYLASE LEVELS CAN INDICATE PANCREATITIS, WHICH IS THE INFLAMMATION OF THE PANCREAS. TRAUMA OR INJURY TO THE PANCREAS, GASTRO INTESTINAL PERFORATIONS, ETC, CAN ALSO CAUSE ELEVATED FLUID AMYLASE LEVELS IN THE ABDOMINAL OR PLEURAL FLUID.
19	2	AMYLASE (ASCITIC FLUID)	CNPG3	FLUID	AIR TIGHT CONTAINER	2 Hours	AMYLASE MEASUREMENTS IN ASCITIC FLUID ARE USED TO DETERMINE WHETHER AN EFFUSION ORIGINATES FROM THE PANCREAS OR OTHER ORGANS

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
20	109	ANA -IF (ANTI-NUCLEAR AB- IFA, HEP2, SERUM)	IMMUNO FLUORESCENCE	SERUM	SERUM CONTAINER	24 Hours	AN ANTINUCLEAR ANTIBODY (ANA) TEST IS A BLOOD TEST THAT CHECKS FOR THE PRESENCE OF ANTIBODIES AGAINST THE CELL NUCLEI. THE ANA TEST IS PRIMARILY USED TO HELP DIAGNOSE AUTOIMMUNE DISEASES.
21	65	ANA PROFILE	IMMUNOBLOT	SERUM	SERUM CONTAINER	24 Hours	AN ANA (ANTINUCLEAR ANTIBODY) PROFILE TEST IS A BLOOD TEST THAT EXAMINES THE PRESENCE OF SPECIFIC ANTIBODIES THAT TARGET COMPONENTS OF THE CELL NUCLEUS. THE ANA PROFILE TEST HELPS IN DIAGNOSING AUTOIMMUNE DISORDERS, SUCH AS SYSTEMIC LUPUS ERYTHEMATOSUS (SLE), RHEUMATOID ARTHRITIS (RA), SJÖGREN'S SYNDROME, AND OTHER CONNECTIVE TISSUE DISEASES.
22	170	ANTI CCP	CMIA	SERUM	SERUM CONTAINER	2 Hours	THE PRESENCE OF ANTI-CCP ANTIBODIES IN THE BLOOD IS A SPECIFIC MARKER ASSOCIATED WITH RHEUMATOID ARTHRITIS. THE LEVELS OF ANTI- CCP ANTIBODIES CAN PROVIDE INFORMATION ABOUT THE SEVERITY AND PROGRESSION OF RHEUMATOID ARTHRITIS.
23	478	ANTI MULLARIAN HORMONE (AMH)	CLIA	SERUM	SERUM CONTAINER	3 Hours	THE AMH TEST IS PRIMARILY USED FOR ASSESSING A WOMAN'S OVARIAN RESERVE, TO ESTIMATE THE TIMING OF MENOPAUSE, ASSESSMENT OF PCOS, ETC.
24	1082	ANTI SPERM ANTIBODY	ELISA	SERUM	SERUM CONTAINER	24 Hours	AN ANTI-SPERM ANTIBODY (ASA) TEST IS A BLOOD TEST THAT CHECKS FOR THE PRESENCE OF ANTIBODIES THAT TARGET SPERM. THESE ANTIBODIES CAN POTENTIALLY AFFECT THE ABILITY OF SPERMS TO MOVE PROPERLY AND INTERACT WITH EGG CELLS, WHICH CAN LEAD TO FERTILITY ISSUES. THE TEST IS USUALLY PERFORMED AS PART OF FERTILITY EVALUATION.
25	736	ANTI THYRO GLOBULIN ANTIBODY	TWO STEP IMMUNO ASSAY- CMIA	SERUM	SERUM CONTAINER	2 Hours	ANTI-TG (THYROGLOBULIN ANTIBODY) TESTING IS A BLOOD TEST THAT MEASURES THE LEVELS OF THYROGLOBULIN ANTIBODIES IN THE BLOOD. ANTI- TG TESTING IS PRIMARILY USED FOR: THYROID CANCER MONITORING AND ASSESSING THYROID AUTOIMMUNE DISORDERS.
26	112	ANTI THYRO PEROXIDASE	CMIA	SERUM	SERUM CONTAINER	2 Hours	AN ANTI-TPO TEST IS A BLOOD TEST THAT MEASURES THE LEVELS OF ANTIBODIES AGAINST THYROID PEROXIDASE IN THE BODY. THE PRESENCE OF ANTI-TPO ANTIBODIES IS INDICATIVE OF AUTOIMMUNE THYROID DISORDERS, PARTICULARLY HASHIMOTO'S THYROIDITIS AND SOMETIMES GRAVES' DISEASE.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
27	65	APTT	COAGULATION	PLASMA	3.2% SODIUM CITRATE CONTAINING CONTAINER	2 Hours	PARTIAL THROMBOPLASTIN TIME (PTT) TEST, IS A MEDICAL LABORATORY TEST USED TO ASSESS THE BLOOD'S ABILITY TO FORM CLOTS. THE TEST IS COMMONLY ORDERED TO EVALUATE BLEEDING DISORDERS, MONITOR ANTICOAGULANT THERAPY, AND ASSESS OVERALL COAGULATION FUNCTION.
28	1748	AUTO IMMUNE LIVER PROFILE- LIVER LIA	LINE IMMUNO ASSAY	SERUM	SERUM CONTAINER	24 Hours	AN AUTO IMMUNE LIVER DISEASE PANEL IS A GROUP OF TESTS THAT IS DONE TO CHECK FOR AUTO IMMUNE LIVER DISEASE. AN AUTO IMMUNE LIVER. DISEASE MEANS THAT THE BODY'S IMMUNE SYSTEM IS ATTACKING THE LIVER CELLS. THE PANEL INCLUDES THE FOLLOWING PARAMETERS: AMA M2, LKM1, LC1, SLA, GP210 AND SP 100.
29	POO 160	AUTO IMMUNE LIVER DISEASE PANEL-1	VARIOUS METHODS	SERUM	SERUM CONTAINER	24 Hours	AN AUTO IMMUNE LIVER DISEASE PANEL IS A GROUP OF TESTS THAT IS DONE TO CHECK FOR AUTO IMMUNE LIVER DISEASE. AN AUTO IMMUNE LIVER DISEASE MEANS THAT THE BODY'S IMMUNE SYSTEM IS ATTACKING THE LIVER CELLS. THE PANEL INCLUDES THE FOLLOWING PARAMETERS: ASMA, ANA-IF, AMA M2, LKM1, LC1, SLA, GP210 AND SP100.
30	POO 161	AUTO IMMUNE LIVER DISEASE PANEL -2	VARIOUS METHODS	SERUM	SERUM CONTAINER	24 Hours	AN AUTO IMMUNE LIVER DISEASE PANEL IS A GROUP OF TESTS THAT IS DONE TO CHECK FOR AUTO IMMUNE LIVER DISEASE. AN AUTO IMMUNE LIVER DISEASE MEANS THAT THE BODY'S IMMUNE SYSTEM IS ATTACKING THE LIVER CELLS. THE PANEL INCLUDES THE FOLLOWING PARAMETERS: IGG TOTAL, ASMA, ANA-IF, AMA, AMA M2, LKM1, LC1, SLA, GP210 AND SP100.
31	1673	BETA 2 MICRO GLOBULIN	ELFA(Enzyme Linked Fluorescent Assay)	SERUM	SERUM CONTAINER	3 Hours	BETA 2-MICROGLOBULIN CAN BE USED IN ASSESSING RENAL FUNCTION, PARTICULARLY IN KIDNEY-TRANSPLANT RECIPIENTS AND IN PATIENTS SUSPECTED OF HAVING RENAL TUBULOINTERSTITIAL DISEASE. IT ALSO CAN SERVE AS A NONSPECIFIC BUT RELATIVELY SENSITIVE MARKER OF VARIOUS NEOPLASTIC, INFLAMMATORY, AND INFECTIOUS CONDITIONS.
32	882	BETA 2 GLYCOPROTEIN (IgG)	ELISA	SERUM	SERUM CONTAINER	24 Hours	BETA-2 GLYCOPROTEIN IGG AND IGM ARE ASSOCIATED WITH A CONDITION KNOWN AS ANTIPHOSPHOLIPID SYNDROME (APS).IN APS, THE IMMUNE SYSTEM MISTAKENLY PRODUCES ANTIBODIES (IGG AND IGM) AGAINST THIS PROTEIN. THESE ANTIBODIES, KNOWN AS ANTI-BETA-2 GLYCOPROTEIN I ANTIBODIES, CAN LEAD TO CLOTTING EVENTS BY INTERFERING WITH THE BODY'S NATURAL CLOTTING REGULATION.
33	109	BETA 2 GLYCOPROTEIN (IgM)	ELISA	SERUM	SERUM CONTAINER	24 Hours	BETA-2 GLYCOPROTEIN IGG AND IGM ARE ASSOCIATED WITH A CONDITION KNOWN AS ANTIPHOSPHOLIPID SYNDROME (APS).IN APS, THE IMMUNE SYSTEM MISTAKENLY PRODUCES ANTIBODIES (IGG AND IGM) AGAINST THIS PROTEIN. THESE ANTIBODIES, KNOWN AS ANTI-BETA-2 GLYCOPROTEIN I ANTIBODIES, CAN LEAD TO CLOTTING EVENTS BY INTERFERING WITH THE BODY'S NATURAL CLOTTING REGULATION.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
34	92	BETA HCG	TWO STEP IMMUNO ASSAY CMIA	SERUM	SERUM CONTAINER	2 Hours	THE BETA HCG TEST IS PRIMARILY USED FOR CONFIRMING PREGNANCY. IT HELPS DETERMINE THE APPROXIMATE AGE OF THE FETUS. IT CAN DIAGNOSE A POTENTIAL MISCARRIAGE AND SCREEN FOR DOWN'S SYNDROME. SECONDARY BETA HCG TEST USES ARE EVALUATING CANCERS LIKE LUNG, BREAST, OVARIAN, OR UTERUS AND NON-CANCEROUS CONDITIONS LIKE ULCERS, INFLAMMATORY BOWEL DISEASE, AND CIRRHOSIS.
35	147	BETA-HCG (CSF)	TWO STEP IMMUNO ASSAY CMIA	CSF	AIR TIGHT CONTAINER	2 Hours	THE CSF BETA HCG TEST IS USUALLY PERFORMED WHEN THERE IS A SUSPICION OF A SPECIFIC MEDICAL CONDITION, OFTEN RELATED TO CERTAIN TYPES OF CANCERS OR NEUROLOGICAL DISORDERS. THE PRESENCE OF HCG IN CSF IS SUGGESTIVE OF TUMOR PRESENCE. PURE GERMINOMAS ARE ASSOCIATED WITH LOW HCG CONCENTRATIONS IN BOTH SERUM AND CSF.
36	1641	BICARBONATE, SERUM	PEP-PEPC	SERUM	SERUM CONTAINER	3 Hours	THE BICARBONATE TEST OR A TOTAL CO ₂ TEST, MEASURES THE LEVEL OF BICARBONATE IONS (HCO ₃ ⁻) IN THE BLOOD. BICARBONATE IS AN IMPORTANT ELECTROLYTE AND A COMPONENT OF THE BODY'S ACID-BASE BALANCE SYSTEM, WHICH HELPS MAINTAIN THE BODY'S PH WITHIN A NARROW AND HEALTHY RANGE.
37	18	BILIRUBIN DIRECT	DIAZO REACTION	SERUM	SERUM CONTAINER	2 Hours	IT IS THE FORM OF BILIRUBIN WHICH HAS BEEN CONJUGATED WITH GLUCORONIC ACID AND IS EXCRETED IN THE BILE. PURPOSE OF THIS TEST IS MAINLY TO INVESTIGATE JAUNDICE. THIS TEST IS COMMONLY USED TO MEASURE BILIRUBIN LEVELS IN NEWBORNS WITH INFANT JAUNDICE, DETERMINE WHETHER THERE MIGHT BE BLOCKAGE IN THE BILE DUCTS, ETC.
38	17	BILIRUBIN TOTAL	DIAZONIUM SALT	SERUM	SERUM CONTAINER	2 Hours	THIS TEST IS USED TO FIND OUT HOW WELL THE LIVER IS WORKING. IT IS OFTEN A PART OF THE PANEL OF TESTS THAT MEASURE LIVER FUNCTION. A SMALL AMOUNT OF BILIRUBIN IN YOUR BLOOD IS NORMAL, BUT A HIGH LEVEL MAY BE A SIGN OF LIVER DISEASE.
39	1674	BLADDER TUMOR ANTIGEN (BTA)	IMMUNO- CHROMATOGRAPHY	URINE	URINE CONTAINER	4 Hours	QUALITATIVE ASSAY, WHICH DETECTS BLADDER TUMOR-ASSOCIATED ANTIGEN IN URINE OF PATIENTS DIAGNOSED WITH BLADDER CANCER.
40	19	BUN (BLOOD UREA NITROGEN)	CALCULATION	SERUM	SERUM CONTAINER	2 Hours	THE BLOOD UREA NITROGEN TEST, WHICH IS ALSO CALLED A BUN OR SERUM BUN TEST, MEASURES HOW MUCH WASTE PRODUCT YOU HAVE IN YOUR BLOOD. IF YOUR LEVELS ARE OFF THE NORMAL RANGE, THIS COULD MEAN THAT EITHER YOUR KIDNEYS OR YOUR LIVER MAY NOT BE WORKING PROPERLY.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
41	737	C.PEPTIDE	CMIA	SERUM	SERUM CONTAINER	2 Hours	THE C-PEPTIDE TEST, ALONGSIDE BLOOD GLUCOSE AND INSULIN TESTS, AIDS IN ASSESSING DIABETES. ELEVATED C-PEPTIDE LEVELS MAY INDICATE INSULINOMA OR EARLY TYPE 2 DIABETES, WHILE LOW LEVELS ARE LINKED TO TYPE 1 DIABETES, ADVANCED TYPE 2 DIABETES, OR IMPAIRED INSULIN PRODUCTION.
42	151	C3 (COMPLEMENT C3)	NEPHELOMETRY	SERUM	SERUM CONTAINER	3 Hours	A C3 COMPLEMENT BLOOD TEST MEASURES THE LEVELS OF A SPECIFIC TYPE OF PROTEIN IN THE BLOOD (C3 PROTEINS). AS PART OF THE IMMUNE SYSTEM, C3 PROTEINS DESTROY MICROBES (GERMS) THAT CAN MAKE A PERSON SICK. SOMETIMES, C3 PROTEINS LAUNCH AN ATTACK AGAINST HEALTHY CELLS BY MISTAKE. A C3 COMPLEMENT BLOOD TEST GIVES THE HEALTHCARE PROVIDER INFORMATION ABOUT THE PERSON'S IMMUNE SYSTEM. IT SHOWS HOW PARTS OF THE IMMUNE SYSTEM ARE RESPONDING TO HARMFUL SUBSTANCES. THIS TEST CAN HELP THE HEALTHCARE PROVIDER DIAGNOSE. AUTOIMMUNE DISORDERS (LIKE LUPUS), AS WELL AS OTHER CONDITIONS. IT ALSO HELP TO MONITOR TREATMENT FOR AUTOIMMUNE DISEASE.
43	153	C4 (COMPLEMENT C4)	NEPHELOMETRY	SERUM	SERUM CONTAINER	3 Hours	A COMPLEMENT C4 (ALSO KNOWN AS A COMPLEMENT COMPONENT 4, OR SIMPLY C4) IS A TEST TO MEASURE THE LEVEL OF C4 PROTEINS IN THE BLOOD. THESE PROTEINS PLAY A ROLE IN HOW THE IMMUNE SYSTEM FUNCTIONS AND DEFENDS THE BODY FROM HARMFUL BACTERIA AND VIRUSES. MONITORING C4 PROTEIN LEVELS CAN HELP DOCTORS TO DIAGNOSE AND MONITOR CERTAIN CHRONIC MEDICAL CONDITIONS. C4 TESTS ARE USEFUL BECAUSE AFTER AN INJURY, ILLNESS, OR INFECTION WHEN THE BODY IS ACTIVELY FIGHTING INFLAMMATION, THE AMOUNT OF C4 PROTEIN IN YOUR BLOOD CHANGES. DOCTORS MIGHT ALSO ADVISE TO TAKE A C4 TEST IF A PERSON SHOW SIGNS OF CHRONIC INFECTIONS, KIDNEY PROBLEMS, HEPATITIS, OR AUTOIMMUNE DISEASES LIKE RHEUMATOID ARTHRITIS AND LUPUS.
44	100	CA 125	TWO STEP IMMUNO ASSAY- CMIA	SERUM	SERUM CONTAINER	2 Hours	THE CA-125 TEST IS USED FOR SEVERAL PURPOSES, INCLUDING MONITORING THE PROGRESS OF TREATMENT IN OVARIAN CANCER, ENDOMETRIOSIS, UTERINE FIBROIDS, PELVIC INFLAMMATORY DISEASE, AND SOME OTHER TYPES OF CANCER.
45	349	CA 19-9	CLIA	SERUM	SERUM CONTAINER	3 Hours	CA 19-9 LEVELS ARE OFTEN ELEVATED IN PANCREATIC CANCER. IT'S USED AS A TUMOR MARKER TO MONITOR DISEASE PROGRESSION, RESPONSE TO TREATMENT, AND RECURRENCE IN INDIVIDUALS WITH PANCREATIC CANCER. CA 19-9 LEVELS CAN ALSO BE ELEVATED IN OTHER GASTROINTESTINAL CANCERS, INCLUDING THOSE OF THE BILE DUCTS, LIVER, AND STOMACH.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
46	21	CALCIUM	ARSENazo III	SERUM	SERUM CONTAINER	2 Hours	CALCIUM IS AN ESSENTIAL MINERAL THAT IS CRUCIAL FOR VARIOUS PHYSIOLOGICAL FUNCTIONS. THIS TEST IS USED TO HELP DIAGNOSE OR MONITOR MANY TYPES OF MEDICAL CONDITIONS, INCLUDING CONDITIONS THAT AFFECT BONES, KIDNEYS, DIGESTIVE SYSTEM, NERVOUS SYSTEM, THYROID, AND PARATHYROID GLANDS.
47	89	CEA	CMIA	SERUM	SERUM CONTAINER	2 Hours	HIGH LEVELS OF CEA ARE NORMAL IN HEALTHY, UNBORN BABIES. AFTER BIRTH, CEA LEVELS BECOME VERY LOW OR DISAPPEAR COMPLETELY. A CEA TEST IS MOSTLY USED IN PEOPLE WHO HAVE BEEN DIAGNOSED WITH CANCERS THAT OFTEN INCREASE CEA LEVELS. IT'S MOST COMMONLY USED IN COLORECTAL CANCER.
48	23	CERULO-PLASMIN	IMMUNO TURBIDIMETRY	SERUM	SERUM CONTAINER	2 Hours	CERULOPLASMIN IS A COPPER-BINDING PROTEIN PRODUCED BY THE LIVER THAT PLAYS A ROLE IN TRANSPORTING AND REGULATING COPPER IN THE BODY. THE TEST IS USED TO EVALUATE COPPER METABOLISM AND DIAGNOSE CERTAIN MEDICAL CONDITIONS RELATED TO COPPER IMBALANCES.
49	24	CHLORIDE (CI)	ION-SELECTIVE ELECTRODE (INDIRECT)	SERUM	SERUM CONTAINER	2 Hours	CHLORIDE IS AN ESSENTIAL ELECTROLYTE, ALONG WITH SODIUM AND POTASSIUM, IT HELPS MAINTAIN PROPER FLUID BALANCE, ELECTROLYTE BALANCE, AND ACID-BASE BALANCE IN THE BODY. THE TEST IS USED TO ASSESS CHLORIDE LEVELS AND ELECTROLYTE BALANCE, AS WELL AS DIAGNOSE AND MONITOR VARIOUS MEDICAL CONDITIONS.
50	137	CHLORIDE (CSF)	ION-SELECTIVE ELECTRODE (INDIRECT)	CSF	AIR TIGHT CONTAINER	2 Hours	IT IS AN IMPORTANT ELECTROLYTE IN THE CSF AND PLAYS A ROLE IN MAINTAINING THE PROPER BALANCE OF FLUIDS AND ELECTROLYTES IN THE CENTRAL NERVOUS SYSTEM. ABNORMAL LEVELS OF CSF CHLORIDE MAY INDICATE UNDERLYING NEUROLOGICAL CONDITIONS, SUCH AS MENINGITIS, ENCEPHALITIS, OR HYDROCEPHALUS.
51	1593	CHLORIDE (FLUID)	ION-SELECTIVE ELECTRODE (INDIRECT)	FLUID	AIR TIGHT CONTAINER	2 Hours	USED TO DETECT THE CHLORIDE LEVEL IN FLUID.
52	1412	CHLORIDE, RANDOM URINE	ION-SELECTIVE ELECTRODE (INDIRECT)	URINE	URINE CONTAINER	3 Hours	MONITORING URINE CHLORIDE LEVELS CAN AID IN DIAGNOSING AND MANAGING VARIOUS MEDICAL CONDITIONS, SUCH AS KIDNEY DISORDERS, METABOLIC ACIDOSIS, AND CERTAIN TYPES OF KIDNEY STONES.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
53	1390	CHOLESTEROL FLUID	ENZY MATIC(CHOLEST EROL ESTERASE AND CHOLESTROL OXIDASE)	FLUID	AIR TIGHT CONTAINER	2 Hours	THIS TEST PROVIDES A MEASUREMENT OF CHOLESTEROL IN BODY FLUIDS, WHICH IS USEFUL FOR SCREENING OF A CHOLESTEROL EFFUSION OR CHOLESTEROL-RICH PSEUDOCYHOUS EFFUSION AND TO DISTINGUISH BETWEEN CHYLOUS AND NONCHYLOUS EFFUSIONS.
54	27	CK-MB	KINETIC/IFCC	SERUM	SERUM CONTAINER	3 Hours	CK-MB TEST IS A SPECIFIC FORM OF THE CREATINE KINASE TEST THAT MEASURES THE LEVELS OF THE CK-MB ENZYME IN THE BLOOD. CK-MB IS AN ISOFORM OF CREATINE KINASE THAT IS PRIMARILY FOUND IN HEART MUSCLE (MYOCARDIUM). THE TEST IS USED TO ASSESS DAMAGE TO THE HEART MUSCLE, SPECIFICALLY IN THE CONTEXT OF SUSPECTED HEART ATTACKS.
55	26	CK (CPK TOTAL)	N-ACETYLCYSTEIN (NAC) ACTIVATED	SERUM	SERUM CONTAINER	2 Hours	CK IS AN ENZYME FOUND IN VARIOUS TISSUES, INCLUDING SKELETAL MUSCLES, THE HEART, AND THE BRAIN. THE TEST IS USED TO ASSESS MUSCLE DAMAGE AND DIAGNOSE CONDITIONS THAT AFFECT MUSCLES AND OTHER TISSUES.
56	29	COPPER	DI-BR-PAESA	SERUM	SERUM CONTAINER	3 Hours	COPPER IS AN ESSENTIAL TRACE ELEMENT THAT PLAYS A CRUCIAL ROLE IN VARIOUS PHYSIOLOGICAL PROCESSES, INCLUDING THE FORMATION OF ENZYMES, IRON METABOLISM, CONNECTIVE TISSUE SYNTHESIS, AND THE FUNCTION OF THE NERVOUS AND IMMUNE SYSTEMS. THE TEST IS USED TO ASSESS COPPER LEVELS AND DIAGNOSE CONDITIONS RELATED TO COPPER METABOLISM.
57	106	CORTISOL	ONE STEP IMMUNO ASSAY- CMIA	SERUM	SERUM CONTAINER	2 Hours	CORTISOL TEST CAN HELP DIAGNOSE CONDITIONS THAT AFFECT ADRENAL GLAND FUNCTION, SUCH AS ADDISON'S DISEASE OR CUSHING'S SYNDROME. CORTISOL LEVELS CAN BE USED TO ASSESS THE BODY'S STRESS RESPONSE ALSO.
58	30	CREATININE	KINETIC ALKALINE PICRATE	SERUM	SERUM CONTAINER	2 Hours	CREATININE IS A WASTE PRODUCT THAT IS PRODUCED BY MUSCLES DURING NORMAL METABOLIC PROCESSES AND IS FILTERED OUT OF THE BLOOD BY THE KIDNEYS. THE CREATININE TEST IS COMMONLY USED TO ASSESS KIDNEY FUNCTION AND TO MONITOR KIDNEY HEALTH.
59	7	CREATININE CLEARANCE*	KINETIC ALKALINE PICRATE	SERUM & 24 HOUR URINE	SERUM/URINE CONTAINER	3 Hours	THE PURPOSE OF A CREATININE CLEARANCE TEST IS TO ASSESS KIDNEY FUNCTION. IN PARTICULAR, CREATININE CLEARANCE IS USED TO DETERMINE THE ESTIMATED GLOMERULAR FILTRATION RATE (EGFR), WHICH DESCRIBES HOW WELL THE KIDNEYS ARE FILTERING THE BLOOD.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
60	1200	D-DIMER	IMMUNOTURBIDIMETRY	PLASMA	3.2% SODIUM CITRATE CONTAINING CONTAINER	3 Hours	D-DIMER IS A SMALL PROTEIN FRAGMENT THAT IS PRODUCED WHEN A BLOOD CLOT DISSOLVES IN THE BODY. THE TEST IS USED TO HELP DIAGNOSE AND RULE OUT CONDITIONS INVOLVING BLOOD CLOT FORMATION AND BREAKDOWN.
61	146	DHEA SULPHATE	CLIA	SERUM	SERUM CONTAINER	2 Hours	DHEA-S IS A PRECURSOR HORMONE THAT CAN BE CONVERTED INTO OTHER HORMONES, INCLUDING TESTOSTERONE AND ESTROGEN. DHEA-S TEST MAINLY ASSESSES HEALTH AND FUNCTION OF THE ADRENAL GLANDS, FERTILITY AND REPRODUCTIVE HEALTH IN BOTH MEN AND WOMEN, PCOS, ETC.
62	1544	ERYTHROPOIETIN	CLIA	SERUM	SERUM CONTAINER	3 Hours	EPO IS A HORMONE PRODUCED BY THE KIDNEYS IN RESPONSE TO LOW OXYGEN LEVELS IN THE BODY. THE EPO TEST IS USUALLY DONE FOR DIAGNOSING ANEMIA, MONITORING KIDNEY FUNCTION, ASSESSING CHRONIC DISEASE ETC.
63	1107	ESTIMATED GLOMERULAR FILTRATION RATE (eGFR)	CALCULATION	SERUM	SERUM CONTAINER	3 Hours	THE EGFR TEST, OR ESTIMATED GLOMERULAR FILTRATION RATE TEST, IS A BLOOD TEST USED TO ASSESS HOW WELL THE KIDNEYS ARE FUNCTIONING. THE EGFR TEST PROVIDES AN ESTIMATE OF THE RATE AT WHICH THE KIDNEYS ARE ABLE TO FILTER BLOOD TO PRODUCE URINE.
64	730	ESTRADIOL	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	THE ESTRADIOL TEST IS USED FOR ASSESSING REPRODUCTIVE HEALTH, TO ENSURE PROPER HORMONE BALANCE AND THERAPEUTIC EFFECTIVENESS IN HORMONE REPLACEMENT THERAPY, TO EVALUATE FERTILITY ISSUES AND ASSIST IN DETERMINING THE CAUSE OF INFERTILITY.
65	1667	EXTENDED DOUBLE MARKER WITH PREECLAMPSIA SCREENING	ECLIA	SERUM	SERUM CONTAINER	24 Hours	THIS TEST HELPS TO ASSESS THE RISK OF CHROMOSOMAL ABNORMALITIES SUCH AS DOWN SYNDROME (TRISOMY 21) AND TRISOMY 18/13 TOGETHER WITH RISK ANALYSIS FOR PREECLAMPSIA. MARKERS TESTED ARE PAPP-A, FREE BETA-HCG, AND PIGF.
66	1768	FECAL CALPROTECTIN	CLIA	STOOL	AIR TIGHT CONTAINER	2 Days	FECAL TEST QUANTITATIVELY MEASURES CALPROTECTIN, A PROTEIN FOUND IN NEUTROPHILS, FROM A SIMPLE STOOL SAMPLE. CALPROTECTIN LEVELS REFLECT THE NUMBER OF NEUTROPHILS PRESENT, INDICATING THE SEVERITY OF INTESTINAL INFLAMMATION.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
67	94	FERRITIN	TWO STEP IMMUNO ASSAY. CMIA	SERUM	SERUM CONTAINER	2 Hours	THE SERUM FERRITIN LEVEL IS DIRECTLY RELATED TO THE AMOUNT OF IRON STORED IN THE BODY.FERRITIN LEVELS LOWER THAN THE TYPICAL RANGE SHOW IRON DEFICIENCY. FERRITIN BEING AN ACUTE-PHASE REACTANT PROTEIN, TENDS TO RISE DURING INFLAMMATION. HIGHER VALUES ARE SEEN IN CONDITIONS LIKE HEMOCHROMATOSIS, LIVER DISEASE, HYPERTHYROIDISM AND LEUKEMIAS.
68	1197	FIBRINOGEN	CLOTTING METHOD	PLASMA	3.2% SODIUMCITRATE CONTAINING CONTAINER	3 Hours	FIBRINOGEN IS A PROTEIN PRODUCED BY THE LIVER AND IS ESSENTIAL FOR THE BLOOD CLOTTING PROCESS. THIS TEST IS USED TO ASSESS THE BLOOD'S ABILITY TO FORM CLOTS AND IS OFTEN PART OF ASSESSMENT OF BLOOD CLOTTING DISORDERS AND RELATED CONDITIONS.
69	1637	FIRST TRIMESTER DOUBLE MARKER	ECLIA	SERUM	SERUM CONTAINER	24 Hours	PRENATAL SCREENING TEST ASSESSES THE RISK OF CHROMOSOMAL ABNORMALITIES IN THE FETUS. THIS TEST IS TYPICALLY PERFORMED DURING THE FIRST TRIMESTER OF PREGNANCY, USUALLY 11 TO 13 WEEKS. THE DOUBLE MARKER TEST COMBINES TWO SPECIFIC MEASUREMENTS OF FREE BETA-HCG AND PAPP-A, ALONG WITH MATERNAL AGE TO PROVIDE AN ESTIMATE OF THE RISK FOR ABNORMAL CHROMOSOMAL CONDITIONS, MOSTLY DOWN SYNDROME (TRISOMY 21) AND TRISOMY 18 AND TRISOMY 13.
70	1329	FLUID CREATININE	KINETIC ALKALINE PICRATE	FLUID	AIR TIGHT CONTAINER	3 Hours	TO CHECK FLUID CREATININE LEVELS.
71	78	FLUID- PROTEIN	BIURET	FLUID	AIR TIGHT CONTAINER	3 Hours	A FLUID PROTEIN TEST MEASURES THE AMOUNT OF PROTEIN IN A BODILY FLUID, LIKE ASCITIC, SYNOVIAL OR PLEURAL FLUID. IT HELPS DIAGNOSE VARIOUS CONDITIONS RELATED TO INFLAMMATION, INFECTION, AND CERTAIN DISEASES.
72	1081	FOLATE (FOLIC ACID)	CMIA	SERUM	SERUM CONTAINER	2 Hours	A FOLIC ACID TEST, ALSO KNOWN AS A FOLATE TEST, MEASURES THE LEVELS OF FOLIC ACID (VITAMIN B9) IN THE BLOOD. DURING PREGNANCY, FOLIC ACID INTAKE IS ESSENTIAL TO PREVENT NEURAL TUBE DEFECTS IN THE DEVELOPING FETUS. A FOLIC ACID DEFICIENCY DURING PREGNANCY CAN INCREASE THE RISK OF BIRTH DEFECTS
73	1139	FOOD INTOLERANCE (46 SPECIFIC IGG ANTIBODIES)	MICROARRAY	SERUM/PLASMA	SERUM OR EDTA CONTAINER	5 Hours	THIS TEST UTILISES AN INNOVATIVE, COLORIMETRIC MICROARRAY-BASED ELISA TECHNOLOGY FOR THE MEASUREMENT OF FOOD-SPECIFIC IGG ANTIBODIES IN HUMAN SERUM OR PLASMA FOR OVER 200 DIFFERENT FOODS. THIS TEST USE SPECIFIC FOOD EXTRACTS TO IDENTIFY THE CORRESPONDING LEVEL OF CIRCULATING IGG ANTIBODIES TO THESE POTENTIAL ANTIGENS AND CAN THEREFORE DETECT FOODS TO WHICH THE IMMUNE SYSTEM IS REACTING.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
74	1084	FREE KAPPA LAMBDA LIGHT CHAIN ASSAY	NEPHELOMETRY	SERUM	SERUM CONTAINER	4 Hours	THESE LIGHT CHAINS ARE COMPONENTS OF ANTIBODIES, WHICH ARE PROTEINS PRODUCED BY PLASMA CELLS IN THE IMMUNE SYSTEM. THE TEST IS OFTEN USED IN THE DIAGNOSIS AND MONITORING OF CONDITIONS RELATED TO ABNORMAL ANTIBODY PRODUCTION, SUCH AS MULTIPLE MYELOMA, A TYPE OF CANCER THAT AFFECTS PLASMA CELLS.
75	818	FREE PSA	CLIA	SERUM	SERUM CONTAINER	3 Hours	THE FREE PROSTATE-SPECIFIC ANTIGEN (PSA) TEST MEASURES THE LEVEL OF UNBOUND OR FREE PSA IN THE BLOOD. THE FREE PSA TEST, ALONG WITH THE TOTAL PSA TEST, IS USED TO ASSESS THE RISK OF PROSTATE CANCER, AND TO AID IN MAKING DECISIONS ABOUT FURTHER DIAGNOSTIC TESTS.
76	95	FREE T3	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	FREE T3 REPRESENTS THE BIOLOGICALLY ACTIVE FORM OF T3 THAT IS AVAILABLE FOR USE BY CELLS. FOR INDIVIDUALS UNDERGOING TREATMENT FOR THYROID DISORDERS, SUCH AS MEDICATION FOR HYPOTHYROIDISM, THE FREE T3 TEST IS USED TO MONITOR TREATMENT.
77	102	FREE T4	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	FREE T4 REPRESENTS THE BIOLOGICALLY ACTIVE FORM OF T4 THAT IS AVAILABLE FOR USE BY CELLS. THE TEST IS VALUABLE FOR DIAGNOSING THYROID DISORDERS, ESPECIALLY THOSE RELATED TO T4 LEVELS.
78	105	FSH	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	FSH PLAYS AN IMPORTANT ROLE IN SEXUAL DEVELOPMENT AND FUNCTIONING. IT HELPS TO FIND THE CAUSE OF INFERTILITY, PROBLEMS WITH OVARIAN FUNCTION, REASON FOR IRREGULAR OR STOPPED MENSTRUAL PERIODS, CONFIRMATION OF MENOPAUSE, OR PERIMENOPAUSE.
79	121	FSH (MALE < 19 YEARS)	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	TESTING FSH LEVELS IN PREPUBERTAL MALES MIGHT BE CONDUCTED FOR A VARIETY OF MEDICAL REASONS, SUCH AS TO EVALUATE THE FUNCTIONING OF THE REPRODUCTIVE SYSTEM, DIAGNOSE CERTAIN CONDITIONS, OR ASSESS THE DEVELOPMENT OF PUBERTY, INCASE A MALE CHILD IS NOT SHOWING SIGNS OF PUBERTY BY THE AGE OF 14, OR IF THERE ARE CONCERNS ABOUT THE LACK OF SECONDARY SEXUAL FEATURES ETC.
80	123	FSH(FEMALE <19 YEARS)	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	FSH TEST IS USED TO HELP DIAGNOSE OR EVALUATE:WOMEN WHO HAVE PCOS, OVARIAN CYSTS, ABNORMAL VAGINAL OR MENSTRUAL BLEEDING, PROBLEMS BECOMING PREGNANT, OR INFERTILITY. FSH TESTING IN PREPUBERTAL FEMALES MIGHT BE CONSIDERED IN CASES LIKE EARLY OR PRECOCIOUS PUBERTY, DELAYED PUBERTY, ETC.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
81	31	G6 PD	KINETIC	BLOOD	EDTA CONTAINER	3 Hours	THE TEST IS USED TO MONITOR G6PD ENZYME DEFICIENCY, WHICH IS AN ENZYME THAT PLAYS A CRUCIAL ROLE IN PROTECTING RED BLOOD CELLS FROM DAMAGE CAUSED BY CERTAIN SUBSTANCES. A DEFICIENCY IN G6PD CAN LEAD TO A CONDITION CALLED G6PD DEFICIENCY OR FAVISM.
82	32	GAMMA GT (GGT)	SZASZ & PERSIEN	SERUM	SERUM CONTAINER	3 Hours	GGT IS FOUND PRIMARILY IN THE LIVER AND BILE DUCTS, AND IT CAN PROVIDE INFORMATION ABOUT THE HEALTH OF THE LIVER AND THE BILE DUCTS. IT IS OFTEN USED AS PART OF LIVER FUNCTION TESTS, TO EVALUATE THE FUNCTIONING OF THE LIVER AND TO DETECT POTENTIAL LIVER DISEASES OR CONDITIONS.
83	668	GLOBULIN	CALCULATION	SERUM	SERUM CONTAINER	2 Hours	THE TEST REFERS TO TOTAL SERUM GLOBULIN LEVELS WHICH IS TESTED TO MONITOR CONDITIONS SUCH AS CHRONIC INFLAMMATION, AUTOIMMUNE DISORDERS, INFECTIONS, CERTAIN CANCERS, LIVER DISEASES, KIDNEY DISEASES, MALNUTRITION, AND CERTAIN GENETIC DISORDERS.
84	40	GLUCOSE (PP 1 1/2 HOURS)	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	2 Hours	THE POSTPRANDIAL PERIOD IS THE TIME, AFTER HAVING A MEAL WHEN THE BODY DIGESTS AND ABSORBS NUTRIENTS, CAUSING AN INCREASE IN BLOOD SUGAR LEVELS. MONITORING POSTPRANDIAL BLOOD SUGAR IS IMPORTANT FOR MANAGING DIABETES AND ASSESSING HOW WELL BLOOD SUGAR LEVELS ARE CONTROLLED AFTER EATING.
85	37	GLUCOSE (PP 1/2 HOURS)	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	2 Hours	THE POSTPRANDIAL PERIOD IS THE TIME, AFTER HAVING A MEAL WHEN THE BODY DIGESTS AND ABSORBS NUTRIENTS, CAUSING AN INCREASE IN BLOOD SUGAR LEVELS. MONITORING POSTPRANDIAL BLOOD SUGAR IS IMPORTANT FOR MANAGING DIABETES AND ASSESSING HOW WELL BLOOD SUGAR LEVELS ARE CONTROLLED AFTER EATING.
86	60	GLUCOSE (PP)	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	2 Hours	MONITORING POSTPRANDIAL GLUCOSE LEVELS IS IMPORTANT FOR ASSESSING HOW WELL THE BODY PROCESSES AND MANAGES BLOOD SUGAR AFTER EATING. THE POSTPRANDIAL GLUCOSE TEST SERVES SEVERAL PURPOSES LIKE DIABETES ASSESSMENT, MONITORING DIABETES MANAGEMENT, ETC.
87	38	GLUCOSE (PP 2 HOURS)	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	2 Hours	THE POSTPRANDIAL PERIOD IS THE TIME, AFTER HAVING A MEAL WHEN THE BODY DIGESTS AND ABSORBS NUTRIENTS, CAUSING AN INCREASE IN BLOOD SUGAR LEVELS. MONITORING POSTPRANDIAL BLOOD SUGAR IS IMPORTANT FOR MANAGING DIABETES AND ASSESSING HOW WELL BLOOD SUGAR LEVELS ARE CONTROLLED AFTER EATING.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
88	41	GLUCOSE (PP 3 HOURS)	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	2 Hours	THE POSTPRANDIAL PERIOD IS THE TIME, AFTER HAVING A MEAL WHEN THE BODY DIGESTS AND ABSORBS NUTRIENTS, CAUSING AN INCREASE IN BLOOD SUGAR LEVELS. MONITORING POSTPRANDIAL BLOOD SUGAR IS IMPORTANT FOR MANAGING DIABETES AND ASSESSING HOW WELL BLOOD SUGAR LEVELS ARE CONTROLLED AFTER EATING.
89	62	GLUCOSE (F)	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	2 Hours	THIS TEST IS COMMONLY USED TO ASSESS BLOOD SUGAR LEVELS AND SCREEN FOR DIABETES OR PREDIABETES.
90	34	GLUCOSE (R)	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	2 Hours	RANDOM BLOOD SUGAR, OR CASUAL GLUCOSE, MEASURES BLOOD SUGAR LEVELS ANYTIME, REGARDLESS OF MEALS. UNLIKE FASTING TESTS, IT OFFERS A SNAPSHOT OF YOUR CURRENT BLOOD SUGAR LEVEL.
91	63	GLUCOSE AFTER 1 HR	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	2 Hours	THE GLUCOSE CHALLENGE TEST, ALSO CALLED THE ONE-HOUR GLUCOSE TOLERANCE TEST, MEASURES THE BODY'S RESPONSE TO SUGAR (GLUCOSE). THE GLUCOSE CHALLENGE TEST IS DONE DURING PREGNANCY TO SCREEN FOR GESTATIONAL DIABETES, DIABETES THAT DEVELOPS DURING PREGNANCY.
92	35	GLUCOSE (PP 1 HOUR)	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	2 Hours	THE POSTPRANDIAL PERIOD IS THE TIME, AFTER HAVING A MEAL WHEN THE BODY DIGESTS AND ABSORBS NUTRIENTS, CAUSING AN INCREASE IN BLOOD SUGAR LEVELS. MONITORING POSTPRANDIAL BLOOD SUGAR IS IMPORTANT FOR MANAGING DIABETES AND ASSESSING HOW WELL BLOOD SUGAR LEVELS ARE CONTROLLED AFTER EATING.
93	60	GLUCOSE (PP)	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	2 Hours	MONITORING POSTPRANDIAL GLUCOSE LEVELS IS IMPORTANT FOR ASSESSING HOW WELL THE BODY PROCESSES AND MANAGES BLOOD SUGAR AFTER EATING. THE POSTPRANDIAL GLUCOSE TEST SERVES SEVERAL PURPOSES LIKE DIABETES ASSESSMENT, MONITORING DIABETES MANAGEMENT, ETC.
94	811	GTT	HEXOKINASE	PLASMA	SODIUM FLUORIDE CONTAINING CONTAINER	3 Hours	GLUCOSE TOLERANCE TEST, IS A DIAGNOSTIC TEST USED TO EVALUATE HOW YOUR BODY PROCESSES GLUCOSE (SUGAR) OVER TIME. IT'S COMMONLY USED TO DIAGNOSE DIABETES AND GESTATIONAL DIABETES DURING PREGNANCY.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
95	43	HBA1C GLYCATED HAEMOGLOBIN	ION-EXCHANGE- HPLC	WHOLE BLOOD	EDTA CONTAINER	2 Hours	HEMOGLOBIN A1C, IS A BLOOD TEST THAT MEASURES THE AVERAGE BLOOD SUGAR LEVELS OVER A PERIOD OF APPROXIMATELY TWO TO THREE MONTHS. THIS TEST IS COMMONLY USED TO ASSESS LONG-TERM BLOOD SUGAR CONTROL IN INDIVIDUALS WITH DIABETES AND TO MONITOR HOW WELL DIABETES MANAGEMENT STRATEGIES ARE WORKING.
96	42	HDL CHOLESTEROL (DIRECT)	ACCELERATOR SELECTIVE DETERGENT	SERUM	SERUM CONTAINER	2 Hours	THIS TEST PROVIDES AN ACCURATE MEASUREMENT OF THE AMOUNT OF HDL CHOLESTEROL PRESENT, WHICH IS IMPORTANT FOR ASSESSING AN INDIVIDUAL'S RISK OF HEART DISEASE AND EVALUATING THEIR LIPID PROFILE.
97	1042	HEMOGLOBIN ELECTRO PHORESIS-HPLC (QUANTITATIVE)	ION-EXCHANGE HPLC	BLOOD	EDTA CONTAINER	24 Hours	HEMOGLOBIN ELECTROPHORESIS IS COMMONLY USED TO DIAGNOSE AND DIFFERENTIATE HEMOGLOBINOPATHIES, WHICH ARE GENETIC CONDITIONS CHARACTERIZED BY ABNORMALITIES IN THE STRUCTURE OF HEMOGLOBIN.
98	799	HOMOCYSTEINE	CMIA	SERUM	SERUM CONTAINER	2 Hours	HOMOCYSTEINE IS AN AMINO ACID PRODUCED BY THE BODY AS A PART OF NORMAL METABOLISM. ELEVATED LEVELS OF HOMOCYSTEINE ARE ASSOCIATED WITH AN INCREASED RISK OF CERTAIN HEALTH CONDITIONS, PARTICULARLY CARDIOVASCULAR DISEASES.
99	1639	HUMAN GROWTH HORMONE (HGH)	ECLIA	SERUM	SERUM CONTAINER	3 Hours	A HUMAN GROWTH HORMONE (HGH) TEST IS USED TO MEASURE THE LEVELS OF GROWTH HORMONE IN THE BLOOD. TEST IS TYPICALLY USED TO DIAGNOSE GROWTH-RELATED DISORDERS, SUCH AS GROWTH HORMONE DEFICIENCY OR CONDITIONS THAT AFFECT LINEAR GROWTH. IT IS ALSO USED TO ASSESS THE FUNCTION OF PITUITARY GLAND.
100	159	IGA (IMMUNO GLOBULIN A)	NEPHELOMETRY	SERUM	SERUM CONTAINER	3 Hours	IGA LEVELS CAN GIVE INSIGHTS INTO THE BODY'S IMMUNE RESPONSE, PARTICULARLY IN MUCOSAL AREAS. CHANGES IN IGA LEVELS MIGHT INDICATE IMMUNE SYSTEM DYSFUNCTION OR SPECIFIC IMMUNE CHALLENGES. IGA TESTING MIGHT BE USED TO DIAGNOSE ALLERGIES, RESPIRATORY INFECTIONS, AND GASTROINTESTINAL INFECTIONS.
101	158	IGE	ECLIA	SERUM	SERUM CONTAINER	3 Hours	IGE TESTING IS COMMONLY USED TO DIAGNOSE ALLERGIES.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
102	161	IGG (IMMUNO GLOBULIN)	NEPHELOMETRY	SERUM	SERUM CONTAINER	3 Hours	IGG ANTIBODIES ARE PRODUCED IN RESPONSE TO EXPOSURE TO VARIOUS PATHOGENS, SUCH AS BACTERIA, VIRUSES, AND OTHER ANTIGENS. IGG TESTING CAN HELP DIAGNOSE CURRENT OR PAST INFECTIONS BY IDENTIFYING THE PRESENCE OF SPECIFIC ANTIBODIES AGAINST PARTICULAR PATHOGENS.
103	162	IGM (IMMUNO-GLOBULIN M)	NEPHELOMETRY	SERUM	SERUM CONTAINER	3 Hours	DETECTING IGM ANTIBODIES CAN INDICATE AN ONGOING OR RECENT INFECTION. IGM TESTING IS OFTEN USED TO DIAGNOSE ACUTE INFECTIONS SUCH AS VIRAL INFECTIONS (E.G., HEPATITIS, MEASLES, DENGUE) AND BACTERIAL INFECTIONS (E.G., CERTAIN TYPES OF STREPTOCOCCUS).IGM TESTING CAN ALSO BE USED TO DIAGNOSE CONGENITAL INFECTIONS.
104	699	INSULIN (F)	ONE STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	THIS TEST IS USED TO EVALUATE HOW WELL THE BODY IS REGULATING BLOOD SUGAR LEVELS, PARTICULARLY IN RELATION TO INSULIN PRODUCTION AND INSULIN RESISTANCE. THE FASTING INSULIN TEST SERVES SEVERAL PURPOSES, INCLUDING: DIABETES DIAGNOSIS, ASSESS INSULIN RESISTANCE, EVALUATING METABOLIC SYNDROME, ETC.
105	1145	INHIBIN A	CLIA	SERUM	SERUM CONTAINER	3 Hours	INHIBIN A LEVELS CAN BE USED AS A MARKER FOR MONITORING THE HEALTH AND PROGRESSION OF PREGNANCY. LEVELS OF INHIBIN A INCREASE DURING PREGNANCY, PARTICULARLY DURING THE SECOND TRIMESTER. INHIBIN A, ALONG WITH OTHER MARKERS SUCH AS HCG,AFP AND UE3 IS USED IN COMBINATION TO SCREEN FOR DOWN SYNDROME AND OTHER CHROMOSOMAL ABNORMALITIES DURING THE SECOND TRIMESTER OF PREGNANCY.
106	700	INSULIN (PP)	ONE STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	THIS TEST PROVIDES INFORMATION ABOUT HOW THE BODY RESPONDS TO THE INTAKE OF FOOD IN TERMS OF INSULIN PRODUCTION AND GLUCOSE REGULATION. THE INSULIN POSTPRANDIAL TEST SERVES SEVERAL PURPOSES, INCLUDING DIABETES MANAGEMENT, ASSESS BODY INSULIN SENSITIVITY FOR PERSONALIZED TREATMENT, ETC.
107	971	IONIZED CALCIUM	DIRECT-ION SELECTIVE ELECTRODE	SERUM	SERUM CONTAINER	2 Hours	IONIZED CALCIUM SPECIFICALLY REFERS TO THE BIOLOGICALLY ACTIVE FORM OF CALCIUM THAT IS NOT BOUND TO PROTEINS. IMBALANCES IN IONIZED CALCIUM LEVELS CAN LEAD TO VARIOUS HEALTH ISSUES, INCLUDING MUSCLE WEAKNESS, HEART RHYTHM ABNORMALITIES, BONE DISORDERS AND OVERACTIVITY OR UNDERACTIVITY OF THE PARATHYROID GLANDS.
108	44	IRON	FERENE	SERUM	SERUM CONTAINER	2 Hours	THESE TESTS HELP ASSESS IRON STATUS, DIAGNOSE IRON-RELATED DISORDERS, AND MONITOR THE EFFECTIVENESS OF TREATMENTS. IRON IS AN ESSENTIAL MINERAL REQUIRED FOR VARIOUS BODILY FUNCTIONS, INCLUDING THE PRODUCTION OF RED BLOOD CELLS AND THE TRANSPORT OF OXYGEN.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
109	670	IUI	CAPILLARY	SEMEN	STERILE AIR TIGHT CONTAINER	5 Hours	SEMEN SAMPLE, PROVIDED BY DONOR IS PROCESSED IN THE LABORATORY TO CONCENTRATE THE HEALTHY AND MOTILE SPERM FOR INSEMINATION.
110	1700	LACTATE	ENZYMATIC METHOD	PLASMA	SODIUM FLURIDE CONTAINING CONTAINER	2 Hours	TO DETECT HIGH LEVELS OF LACTATE IN BLOOD, WHICH MIGHT BE AN INDICATION OF HYPOXIA OR THE PRESENCE OF OTHER CONDITIONS THAT CAUSE EXCESS PRODUCTION OR INSUFFICIENT CLEARING OF LACTATE FROM THE BLOOD.
111	45	L.D.H (BLOOD)	LACTATE-PYRUVATE UV ABSORBANCE	SERUM	SERUM CONTAINER	2 Hours	LDH IS AN ENZYME THAT IS FOUND IN VARIOUS TISSUES THROUGHOUT THE BODY, INCLUDING THE HEART, LIVER, KIDNEYS, MUSCLES, AND BLOOD CELLS. THE LDH TEST IS COMMONLY USED AS A GENERAL INDICATOR OF TISSUE DAMAGE, INFLAMMATION, OR DISEASE AFFECTING THESE DIFFERENT BODY SYSTEMS.
112	763	LDH (ASCITIC FLUID)	LACTATE-PYRUVATE UV ABSORBANCE	FLUID	AIR TIGHT CONTAINER	2 Hours	LDH ASCITIC FLUID IS A GOOD BIOCHEMICAL MARKER FOR SPONTANEOUS BACTERIAL PERITONITIS. ELEVATED LDH ASCITIC FLUID (>400) OR ELEVATED ASCITIC/SERUM LDH RATIO (>0.6) HAS BEEN ASSOCIATED WITH MALIGNANCY.
113	177	LDH (PLEURAL FLUID)	LACTATE-PYRUVATE UV ABSORBANCE	FLUID	AIR TIGHT CONTAINER	2 Hours	LDH LEVELS IN PLEURAL FLUID HELP DISTINGUISH TRANSUDATIVE FROM EXUDATIVE EFFUSIONS. TRANSUDATIVE EFFUSIONS (E.G., HEART FAILURE) HAVE LOWER LDH, WHILE EXUDATIVE EFFUSIONS (E.G., INFLAMMATION, INFECTIONS, MALIGNANCIES) SHOW HIGHER LDH.
114	971	LDH (FLUID)	LACTATE-PYRUVATE UV ABSORBANCE	FLUID	AIR TIGHT CONTAINER	2 Hours	LDH (LACTATE DEHYDROGENASE) TEST MEASURES LDH LEVELS IN BODY FLUIDS LIKE PERITONEAL, CEREBROSPINAL (CSF), OR SYNOVIAL FLUID. ELEVATED LEVELS MAY INDICATE INFECTIONS, CANCER, OR LUNG-RELATED CONDITIONS.
115	46	LDL CHOLESTEROL	CALCULATION	SERUM	SERUM CONTAINER	2 Hours	THE LDL CHOLESTEROL TEST, ALSO KNOWN AS A LOW-DENSITY LIPOPROTEIN CHOLESTEROL TEST, MEASURES THE LEVEL OF LDL CHOLESTEROL IN THE BLOOD. THIS TEST IS A CRUCIAL PART OF ASSESSING AN INDIVIDUAL'S RISK OF CARDIOVASCULAR DISEASE AND OVERALL LIPID PROFILE.
116	895	LDL CHOLESTEROL DIRECT	MEASURED, LIQUID SELECTIVE DETERGENT	SERUM	SERUM CONTAINER	2 Hours	A DIRECT LDL TEST MEASURES LDL-C IN THE BLOOD, PROVIDING A MORE ACCURATE ASSESSMENT OF CHOLESTEROL LEVELS AND CARDIOVASCULAR DISEASE RISK, INCLUDING HEART ATTACKS AND STROKES.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
117	664	LDL/HDL RATIO	CALCULATION	SERUM	SERUM CONTAINER	2 Hours	THIS TEST IS USED AS A QUICK WAY TO ASSESS THE BALANCE BETWEEN "BAD" CHOLESTEROL (LDL) AND "GOOD" CHOLESTEROL (HDL) IN THE BLOOD, PROVIDING INSIGHT INTO CARDIOVASCULAR RISK.
118	119	LH (FEMALE < 16 YEARS)	TWO STEP IMMUNO ASSAY- CMIA	SERUM	SERUM CONTAINER	2 Hours	LH (LUTEINIZING HORMONE) IS A KEY HORMONE THAT PLAYS A CRUCIAL ROLE IN THE REGULATION OF THE MENSTRUAL CYCLE AND OVULATION. MONITORING LH LEVELS IN FEMALES CAN PROVIDE IMPORTANT INFORMATION ABOUT THE FUNCTIONING OF THE REPRODUCTIVE SYSTEM.
119	117	LH (MALE <16 YEARS)	TWO STEP IMMUNO ASSAY- CMIA	SERUM	SERUM CONTAINER	2 Hours	IN MALES, LUTEINIZING HORMONE (LH) IS RESPONSIBLE FOR STIMULATING THE PRODUCTION OF TESTOSTERONE IN THE TESTES. MONITORING LH LEVELS IN MALES CAN PROVIDE VALUABLE INFORMATION ABOUT TESTICULAR FUNCTION AND OVERALL REPRODUCTIVE HEALTH.
120	402	LIPASE	KINETIC	SERUM	SERUM CONTAINER	3 Hours	THE LIPASE BLOOD TEST HELPS DIAGNOSE AND MONITOR ACUTE AND CHRONIC PANCREATITIS, PANCREATIC CANCER, CYSTS, GALLBLADDER ISSUES (E.G., GALLSTONES OR BLOCKAGES), AND DIGESTIVE DISORDERS LIKE CELIAC OR CROHN'S DISEASE.
121	357	LIPOPROTEIN (a)	NEPHELOMETRY	SERUM	SERUM CONTAINER	3 Hours	THIS TEST ASSESSES THE RISK OF HEART DISEASE, STROKE, AND BLOOD VESSEL ISSUES, ESPECIALLY IN CASES LIKE FAMILIAL HYPERCHOLESTEROLEMIA OR HIGH LDL CHOLESTEROL.
122	710	LUPUS ANTI COAGULANT	COAGULANTION	PLASMA	3.2% SODIUM CITRATE CONTAINING CONTAINER	5 Days	LUPUS ANTICOAGULANT TESTING IS A SERIES OF TESTS USED TO DETECT LUPUS ANTICOAGULANT (LA) IN THE BLOOD. LA IS AN AUTOANTIBODY ASSOCIATED WITH EXCESS BLOOD CLOT FORMATION.
123	824	MAGNESIUM	CALMAGITE METHOD	SERUM	SERUM CONTAINER	3 Hours	A MAGNESIUM TEST MEASURES BLOOD MAGNESIUM, AN ESSENTIAL ELECTROLYTE NEEDED FOR MUSCLE, NERVE, AND HEART FUNCTION, AS WELL AS REGULATING BLOOD PRESSURE AND SUGAR.
124	1215	METANEPHRINES URINE 24 HOUR	COLUMN CHROMATOGRAPHY	24 HOUR URINE	URINE CONTAINER	24 Hours	THIS TEST IS PRIMARILY USED TO HELP DIAGNOSE CERTAIN ADRENAL GLAND TUMORS, SPECIFICALLY PHEOCHROMOCYTOMAS AND PARAGANGLIOMAS, WHICH ARE RARE BUT POTENTIALLY SERIOUS CONDITIONS THAT INVOLVE THE OVERPRODUCTION OF EPINEPHRINE AND NOREPINEPHRINE.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
125	1590	MICROALBUMIN, 24 HOUR URINE	IMMUNO TURBIDIMETRY	24 HOUR URINE	URINE CONTAINER	3 Hours	A 24-HOUR URINE MICROALBUMIN TEST MEASURES THE AMOUNT OF ALBUMIN IN A 24-HOUR COLLECTION OF URINE. ELEVATED LEVELS OF ALBUMIN IN THE URINE, KNOWN AS MICROALBUMINURIA, CAN BE AN EARLY SIGN OF KIDNEY DYSFUNCTION, PARTICULARLY IN INDIVIDUALS WITH DIABETES OR HYPERTENSION.
126	POO 113	MULTIPLE MYELOMA PANEL (SCREENING)	VARIOUS METHODS	SERUM	SERUM CONTAINER	3 Days	MULTIPLE MYELOMA IS A CANCER THAT FORMS IN A TYPE OF WHITE BLOOD CELL CALLED PLASMA CELL. IN THIS CONDITION CANCEROUS PLASMA CELLS BUILD UP IN THE BORN MARROW. PARAMETERS COVERED: SERUM PROTEIN ELECTROPHORESIS, IMMUNOTYPING, IMMUNOGLOBULINS AND FREE LIGHT CHAIN ASSAY.
127	POO 146	MULTIPLE MYELOMA PANEL (COMPREHENSIVE)	VARIOUS METHODS	SERUM	SERUM CONTAINER	3 Days	MULTIPLE MYELOMA IS A CANCER THAT FORMS IN A TYPE OF WHITE BLOOD CELL CALLED PLASMA CELL. IN THIS CONDITION CANCEROUS PLASMA CELLS BUILD UP IN THE BORN MARROW. PARAMETERS COVERED: SERUM PROTEIN ELECTROPHORESIS, IMMUNOTYPING, FREE LIGHT CHAIN ASSAY & BETA 2 MICROGLOBULIN.
128	POO 147	MULTIPLE MYELOMA PANEL (ADVANCED)	VARIOUS METHODS	SERUM	SERUM CONTAINER	3 Days	MULTIPLE MYELOMA IS A CANCER THAT FORMS IN A TYPE OF WHITE BLOOD CELL CALLED PLASMA CELL. IN THIS CONDITION CANCEROUS PLASMA CELLS BUILD UP IN THE BORN MARROW. PARAMETERS COVERED: SERUM PROTEIN ELECTROPHORESIS, IMMUNOTYPING,, FREE LIGHT CHAIN ASSAY, BETA 2 MICROGLOBULIN, CALCIUM, LDH,IGA, IGG, IGM,COMPLETE HAEMOGRAM, UREA, CREATININE & URINE BJP.
129	1751	MITO-CHONDRIAL ANTIBODY (AMA)	IMMUNO-FLUORESCENCE	SERUM	SERUM CONTAINER	24 Hours	AMA ARE SUBSTANCES (ANTIBODIES) THAT ARE FORMED AGAINST MITOCHONDRIA. THIS TEST IS MOST OFTEN USED TO DIAGNOSE PRIMARY BILIARY CHOLANGITIS, FORMERLY KNOWN AS PRIMARY BILIARY CIRRHOSIS (PBC).
130	1191	MYOSITIS PANEL (16 Antigens)	IMMUNOBLOT	SERUM	SERUM CONTAINER	24 Hours	A MYOSITIS PANEL IS A SET OF BLOOD TESTS THAT ARE USED TO DIAGNOSE AND DIFFERENTIATE VARIOUS TYPES OF MYOSITIS, WHICH ARE INFLAMMATORY MUSCLE DISEASES. SPECIFIC ANTIBODIES AND MARKERS ASSOCIATED WITH DIFFERENT FORMS OF MYOSITIS ARE DETECTED IN THIS TEST.
131	454	PAPP-A	CLIA	SERUM	SERUM CONTAINER	3 Hours	THE PAPP-A TEST IS COMMONLY USED IN COMBINATION WITH OTHER TESTS, SUCH AS NUCHAL TRANSLUCENCY ULTRASOUND, TO ASSESS THE RISK OF CERTAIN CHROMOSOMAL ABNORMALITIES, PARTICULARLY DOWN SYNDROME (TRISOMY 21) AND OTHER TRISOMIES.
132	47	PHOSPHOROUS	PHOSPHO MOLYBDATE FORMATION	SERUM	SERUM CONTAINER	2 Hours	THIS TEST IS ORDERED TO SEE HOW MUCH PHOSPHOROUS IS IN YOUR BLOOD. KIDNEY, LIVER, AND CERTAIN BONE DISEASES CAN CAUSE ABNORMAL PHOSPHOROUS LEVELS. IF PHOSPHATE LEVELS IN YOUR BLOOD ARE TOO HIGH OR TOO LOW, IT CAN BE A SIGN OF KIDNEY DISEASE OR OTHER SERIOUS DISORDER.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
133	1612	PIVKA-II (DES-CARBOXYLATED PROTHROMBIN)	CMIA	SERUM	SERUM CONTAINER	2 Hours	THE PIVKA-II TEST, ALSO KNOWN AS THE DES-GAMMA CARBOXY PROTHROMBIN (DCP) TEST, IS A BLOOD TEST USED TO DETECT THE PRESENCE OF A PROTEIN CALLED PIVKA-II. THIS PROTEIN IS AN ABNORMAL FORM OF PROTHROMBIN, A PROTEIN INVOLVED IN BLOOD CLOTting, PRODUCED IN THE LIVER WHEN THERE'S A DEFICIENCY OF VITAMIN K. THE TEST IS PRIMARILY USED AS A SCREENING TOOL FOR HEPATOCELLULAR CARCINOMA (HCC), A TYPE OF LIVER CANCER, ESPECIALLY IN INDIVIDUALS AT HIGH RISK DUE TO CHRONIC LIVER DISEASE.
134	1640	PIGF (PLACENTAL GROWTH FACTOR)	ECLIA	SERUM	SERUM CONTAINER	3 Hours	PLACENTAL GROWTH FACTOR (PIGF) IS A PROTEIN THAT IS PRODUCED BY THE PLACENTA DURING PREGNANCY. IT PLAYS A ROLE IN THE GROWTH OF BLOOD VESSELS AND IS INVOLVED IN THE DEVELOPMENT OF THE PLACENTA AND THE FETUS. THE LEVELS OF PIGF IN A PREGNANT WOMAN'S BLOOD CAN PROVIDE INFORMATION ABOUT THE HEALTH OF THE PLACENTA AND THE VASCULAR SYSTEM. THE TEST IS OFTEN USED AS PART OF THE ASSESSMENT OF A PREGNANT WOMAN'S RISK FOR DEVELOPING PREECLAMPSIA.
135	48	POTASSIUM	ION-SELECTIVE ELECTRODE (INDIRECT)	SERUM	SERUM CONTAINER	2 Hours	A POTASSIUM TEST, PART OF A BASIC METABOLIC PANEL, CHECKS FOR ELECTROLYTE IMBALANCES, MONITORS MEDICATIONS LIKE DIURETICS AND HEART DRUGS, AND HELPS DIAGNOSE HEART ISSUES, KIDNEY DISEASE, OR METABOLIC ACIDOSIS.
136	1663	POTASSIUM (FLUID)	ION-SELECTIVE ELECTRODE (INDIRECT)	FLUID	AIR TIGHT CONTAINER	2-3 Days	POTASSIUM IS AN ESSENTIAL ELECTROLYTE THAT PLAYS A VITAL ROLE IN MAINTAINING PROPER NERVE AND MUSCLE FUNCTION, FLUID BALANCE, AND HEART RHYTHM. MONITORING POTASSIUM LEVELS IS CRUCIAL FOR OVERALL HEALTH.
137	1409	POTASSIUM, RANDOM URINE	ION-SELECTIVE ELECTRODE (INDIRECT)	URINE	URINE CONTAINER	3 Hours	POTASSIUM IS A KEY ELECTROLYTE VITAL FOR MUSCLE CONTRACTIONS, NERVE IMPULSES, FLUID BALANCE, AND HEART RHYTHM. URINE POTASSIUM LEVELS REVEAL POTASSIUM BALANCE AND KIDNEY HEALTH.
138	1631	PREECLAMPSIA PREDICTION PANEL (SFLT 1:PIGF RATION)	ECLIA	SERUM	SERUM CONTAINER	24 Hours	THE SFIT-1/PIGF RATIO IS USED AS A MARKER FOR ASSESSING THE RISK OF PREECLAMPSIA. A HIGH RATIO IS OFTEN ASSOCIATED WITH AN INCREASED RISK OF DEVELOPING PREECLAMPSIA, AND THIS CAN BE AN INDICATION FOR FURTHER EVALUATION AND MANAGEMENT.
139	731	PROGESTERONE	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	PROGESTERONE IS A HORMONE THAT PLAYS A CRUCIAL ROLE IN VARIOUS PROCESSES IN THE FEMALE REPRODUCTIVE SYSTEM, PARTICULARLY DURING THE MENSTRUAL CYCLE AND PREGNANCY. THE PROGESTERONE TEST IS USED FOR SEVERAL PURPOSES LIKE EVALUATING OVULATION, MONITORING PREGNANCY, MONITORING HORMONE REPLACEMENT THERAPY AND ASSESSING FERTILITY ISSUES.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
140	103	PROLACTIN	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	PROLACTIN IS OFTEN DETECTED IN HIGH LEVELS IN INFANTS AND PREGNANT WOMEN. WOMEN WHO ARE NOT PREGNANT USUALLY HAVE LOW LEVELS OF PROLACTIN IN THE BLOOD. THE TEST CAN HELP TO DIAGNOSE CERTAIN HEALTH CONDITIONS AND MONITOR PATIENTS WITH A TYPE OF PITUITARY TUMOR CALLED PROLACTINOMA.
141	1540	PROSTATE HEALTH INDEX(PHI), SERUM	CLIA	SERUM	SERUM CONTAINER	3 Hours	THE PROSTATE HEALTH INDEX (PHI) TEST ASSESSES PROSTATE CANCER RISK BY COMBINING PSA, FREE PSA, AND P2PSA LEVELS FOR GREATER ACCURACY THAN TRADITIONAL PSA TESTING.
142	175	PROTEINS (PERITONEAL FLUID)	BIURET	FLUID	AIR TIGHT CONTAINER	2 Hours	MEASURING PROTEIN LEVELS IN PERITONEAL FLUID CAN SERVE VARIOUS DIAGNOSTIC PURPOSES LIKE ASCITES. ELEVATION IS OFTEN SEEN IN CONDITIONS SUCH AS LIVER CIRRHOSIS, HEART FAILURE, CANCER, AND CERTAIN INFECTIONS AND IN CASE OF MALIGNANCIES.
143	4	PROTEINS (ASCITIC FLUID)	BIURET	FLUID	AIR TIGHT CONTAINER	2 Hours	THE TEST DETERMINES THE TOTAL PROTEIN LEVEL IN ASCITIC FLUID. ASCITIC FLUID TOTAL PROTEIN IS A USEFUL MARKER IN NON-PORTAL HYPERTENSIVE ASCITES.
144	141	PROTEINS(CSF)	BENZETHONIUM CHLORIDE	CSF	AIR TIGHT CONTAINER	2 Hours	CSF PROTEIN TESTING IS GENERALLY USED IN DIAGNOSING NEUROLOGICAL DISORDERS. ELEVATED PROTEIN LEVELS MIGHT BE SEEN IN CONDITIONS SUCH AS MULTIPLE SCLEROSIS, GUILLAIN-BARRÉ SYNDROME, CERTAIN INFECTIONS (LIKE VIRAL OR BACTERIAL MENINGITIS), AND OTHER INFLAMMATORY OR AUTOIMMUNE DISORDERS AFFECTING THE CENTRAL NERVOUS SYSTEM.
145	178	PROTEINS (PLEURAL FLUID)	BIURET	FLUID	AIR TIGHT CONTAINER	2 Hours	THE PROTEIN COMPOSITION CAN HELP DISTINGUISH BETWEEN DIFFERENT TYPES OF PLEURAL EFFUSION, SUCH AS TRANSUDATES AND EXUDATES. THE PROTEIN CONTENT OF PLEURAL FLUID CAN CHANGE IN RESPONSE TO INFECTIONS SUCH AS BACTERIAL OR FUNGAL PNEUMONIA.
146	149	PROTHROMBIN TIME	COAGULATION	PLASMA	3.2% SODIUM CITRATE CONTAINING CONTAINER	2 Hours	IT'S A COMMON TEST USED TO EVALUATE THE EXTRINSIC AND COMMON PATHWAYS OF THE COAGULATION CASCADE. PT IS USED TO ASSESS THE FUNCTION OF VARIOUS CLOTTING FACTORS AND IS OFTEN ORDERED TO MONITOR ANTICOAGULANT THERAPY, DIAGNOSE BLEEDING DISORDERS, AND EVALUATE LIVER FUNCTION.
147	90	PSA (PROSTATE SPECIFIC ANTIGEN)	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	THE PSA TEST IS DONE TO HELP SCREEN FOR AND FOLLOW PROSTATE CANCER IN MEN. IT IS ALSO USED TO TRACK PEOPLE AFTER PROSTATE CANCER TREATMENT TO SEE IF THE CANCER HAS COME BACK.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
148	772	PTH, INTACT	CLIA	SERUM	SERUM CONTAINER	2 Hours	THE PTH TEST IS USED FOR SEVERAL PURPOSES, LIKE DIAGNOSE AND MONITOR HYPERPARATHYROIDISM, HYPOPARATHYROIDISM, ETC.
149	1144	QUADRUPLE TEST	CLIA	SERUM	SERUM CONTAINER	24 Hours	THIS PRENATAL SCREENING, DONE BETWEEN WEEKS 15-21 WEEKS AND 6 DAYS, MEASURES AFP, BETA-HCG, ESTRIOL, AND INHIBIN-A IN MATERNAL BLOOD TO ESTIMATE THE RISK OF CHROMOSOMAL ABNORMALITIES AND NEURAL TUBE DEFECTS.
150	672	RENAL STONE ANALYSIS		STONE	AIR TIGHT CONTAINER	3 Days	THIS ANALYSIS HELPS IDENTIFY THE COMPOSITION OF THE STONE, WHICH IS CRUCIAL FOR DETERMINING THE UNDERLYING CAUSES OF KIDNEY STONE FORMATION AND GUIDING APPROPRIATE TREATMENT AND PREVENTION STRATEGIES. COMMON STONE COMPONENTS INCLUDE CALCIUM OXALATE, CALCIUM PHOSPHATE, URIC ACID, STRUVITE, AND CYSTINE.
151	49	S.G.O.T. (AST)	NADH (WITHOUT P-5'-P)	SERUM	SERUM CONTAINER	2 Hours	SGOT IS FOUND IN VARIOUS TISSUES THROUGHOUT THE BODY, INCLUDING THE LIVER, HEART, MUSCLES, AND OTHER ORGANS. THE TEST IS USED TO ASSESS THE HEALTH OF THE LIVER AND OTHER ORGANS AND TO DIAGNOSE AND MONITOR VARIOUS MEDICAL CONDITIONS.
152	50	S.G.P.T. (ALT)	NADH (WITHOUT P-5'-P)	SERUM	SERUM CONTAINER	2 Hours	SGPT IS FOUND PRIMARILY IN THE LIVER AND TO A LESSER EXTENT IN OTHER TISSUES. THE TEST IS COMMONLY USED TO ASSESS LIVER HEALTH AND TO DIAGNOSE AND MONITOR VARIOUS LIVER-RELATED CONDITIONS.
153	130	SERUM PROTEIN ELECTROPHORESIS	CAPILLARY ELECTROPHORESIS	SERUM	SERUM CONTAINER	24 Hours	THIS TEST DETECTS AND DIAGNOSES CONDITIONS AFFECTING BLOOD PROTEIN LEVELS. GAMMA GLOBULIN SPIKES MAY INDICATE MULTIPLE MYELOMA, WHILE OTHER PROTEIN CHANGES CAN SIGNAL LIVER DISEASE, KIDNEY DISORDERS, OR IMMUNE ABNORMALITIES.
154	58	SERUM PROTEINS	VARIOUS METHODS	SERUM	SERUM CONTAINER	2 Hours	SERUM PROTEIN TESTING IS USED FOR ASSESSING LIVER FUNCTION, MONITORING NUTRITIONAL STATUS, DETECTING INFLAMMATIONS, ETC.
155	1653	SFLT-1 (SOLUBLE FMS-LIKE TYROSINE KINASE-1)	ECLIA	SERUM	SERUM CONTAINER	3 Hours	SFLT-1 IS A PROTEIN THAT PLAYS A ROLE IN REGULATING BLOOD VESSEL GROWTH. DURING PREGNANCY, PARTICULARLY IN CASES OF PREECLAMPSIA, THERE CAN BE AN IMBALANCE IN THE PRODUCTION OF SFIT-1.
156	1750	SMOOTH MUSCLE ANTIBODY	IMMUNO-FLUORESCENCE	SERUM	SERUM CONTAINER	24 Hours	HIGH LEVELS OF SMAS ARE USUALLY A SIGN OF CHRONIC LIVER CONDITION CALLED AUTO IMMUNE HEPATITIS.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
157	51	SODIUM (Na+)	ION-SELECTIVE ELECTRODE (INDIRECT)	SERUM	SERUM CONTAINER	2 Hours	SODIUM IS AN ESSENTIAL ELECTROLYTE THAT PLAYS A VITAL ROLE IN MAINTAINING FLUID BALANCE, NERVE FUNCTION, AND OVERALL CELLULAR FUNCTION IN THE BODY. THIS TEST IS USED TO ASSESS ELECTROLYTE BALANCE AND DIAGNOSE VARIOUS HEALTH CONDITIONS.
158	1407	SODIUM 24-HR URINE	ION-SELECTIVE ELECTRODE (INDIRECT)	24 HOUR URINE	URINE CONTAINER	3 Hours	SODIUM IS AN ESSENTIAL ELECTROLYTE THAT PLAYS A VITAL ROLE IN MAINTAINING FLUID BALANCE, NERVE FUNCTION, AND OVERALL CELLULAR FUNCTION IN THE BODY. THIS TEST IS USED TO ASSESS ELECTROLYTE BALANCE AND DIAGNOSE VARIOUS HEALTH CONDITIONS.
159	1662	SODIUM (FLUID)	ION-SELECTIVE ELECTRODE (INDIRECT)	FLUID	AIR TIGHT CONTAINER	2 Hours	SODIUM IS AN ESSENTIAL ELECTROLYTE THAT PLAYS A VITAL ROLE IN MAINTAINING FLUID BALANCE, NERVE FUNCTION, AND MUSCLE CONTRACTIONS. ANALYZING SODIUM LEVELS IN BODILY FLUIDS CAN PROVIDE IMPORTANT INFORMATION ABOUT A PERSON'S HEALTH AND ELECTROLYTE BALANCE.
160	1408	SODIUM, RANDOM URINE	ION-SELECTIVE ELECTRODE (INDIRECT)	URINE	URINE CONTAINER	3 Hours	SODIUM IS AN ESSENTIAL ELECTROLYTE IN THE BODY THAT PLAYS A CRUCIAL ROLE IN MAINTAINING FLUID BALANCE, BLOOD PRESSURE, NERVE FUNCTION, AND MUSCLE CONTRACTIONS. THE LEVEL OF SODIUM IN THE URINE CAN PROVIDE VALUABLE INFORMATION ABOUT THE BODY'S SODIUM BALANCE AND ITS OVERALL FLUID STATUS.
161	5	SUGAR (ASCITIC FLUID)	HEXOKINASE	FLUID	AIR TIGHT CONTAINER	2 Hours	DETERMINES THE GLUCOSE LEVEL IN ASCITIC FLUID. THIS TEST IS USEFUL IN DIFFERENTIATING PATIENTS WITH SPONTANEOUSLY INFECTED ASCITIC FLUID FROM THOSE WITH GUT PERFORATION.
162	82	SUGAR (FLUID)	HEXOKINASE	FLUID	AIR TIGHT CONTAINER	2 Hours	DETERMINES THE GLUCOSE LEVEL IN FLUID.
163	144	SUGAR (GLUCOSE) (CSF)	HEXOKINASE	CSF	AIR TIGHT CONTAINER	2 Hours	MEASURING CSF GLUCOSE LEVELS IS A VALUABLE DIAGNOSTIC TOOL IN ASSESSING VARIOUS NEUROLOGICAL CONDITIONS. ABNORMAL CSF GLUCOSE LEVELS CAN INDICATE INFECTIONS AFFECTING THE CENTRAL NERVOUS SYSTEM, SUCH AS BACTERIAL OR FUNGAL MENINGITIS. IN THESE CASES, GLUCOSE LEVELS ARE OFTEN LOWER THAN NORMAL AS PATHOGENS CAN CONSUME GLUCOSE.
164	174	SUGAR (PERICARDIAL)	HEXOKINASE	FLUID	AIR TIGHT CONTAINER	2 Hours	THE PRESENCE OF SUGAR (GLUCOSE) IN PERICARDIAL FLUID MIGHT BE ASSESSED FOR VARIOUS REASONS LIKE CONDITIONS AFFECTING THE HEART, SUCH AS PERICARDITIS (INFLAMMATION OF THE PERICARDIUM) OR INFECTIONS OF THE PERICARDIUM.
165	75	SUGAR (PERITONEAL)	HEXOKINASE	FLUID	AIR TIGHT CONTAINER	2 Hours	GLUCOSE MEASUREMENT IN BODY FLUID MAY BE USEFUL, WITH OTHER LABORATORY TESTS, TO EVALUATE EFFUSIONS. DECREASED CONCENTRATIONS ARE ASSOCIATED WITH BACTERIAL INFECTIONS, INFLAMMATION SUCH AS RHEUMATOID ARTHRITIS, AND OCCASIONALLY MALIGNANCY.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
166	183	SUGAR (SYNOVIAL FLUID)	HEXOKINASE	FLUID	AIR TIGHT CONTAINER	2 Hours	CHANGES IN SYNOVIAL FLUID SUGAR LEVELS CAN PROVIDE DIAGNOSTIC INSIGHTS INTO JOINT INFECTIONS, SUCH AS SEPTIC ARTHRITIS, CRYSTAL ARTHROPATHIES, INFLAMMATORY ARTHRITIS, JOINT INJURIES, ETC.
167	176	SUGAR (PLEURAL FLUID)	HEXOKINASE	FLUID	AIR TIGHT CONTAINER	2 Hours	PLEURAL FLUID SUGAR LEVEL CAN HELP TO DIFFERENTIATE BETWEEN DIFFERENT CAUSES OF PLEURAL EFFUSION. CERTAIN INFECTIONS, SUCH AS BACTERIAL PNEUMONIA OR TUBERCULOSIS, CAN CAUSE GLUCOSE LEVELS IN PLEURAL FLUID TO BE LOWER THAN NORMAL. LOW PLEURAL GLUCOSE LEVELS IS A CHARACTERISTIC FEATURE OF EMPYEMA.
168	52	T.I.B.C.	CALCULATION	SERUM	SERUM CONTAINER	2 Hours	TIBC, OR TOTAL IRON BINDING CAPACITY TEST MAY BE USED ALONG WITH OTHER IRON TESTS TO ASSESS THE AMOUNT OF IRON CIRCULATING IN THE BLOOD, THE TOTAL CAPACITY OF THE BLOOD TO TRANSPORT IRON, AND THE AMOUNT OF STORED IRON IN THE BODY. TESTING MAY ALSO HELP DIFFERENTIATE VARIOUS CAUSES OF ANEMIA.
169	107	TESTOSTERONE	DELAYED ONE STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	THE TESTOSTERONE TEST CAN HELP DIAGNOSE CONDITIONS RELATED TO HORMONE IMBALANCES, SUCH AS HYPOGONADISM (LOW TESTOSTERONE LEVELS) OR HYPERGONADISM (HIGH TESTOSTERONE LEVELS). IN CASES WHERE TESTOSTERONE REPLACEMENT THERAPY IS PRESCRIBED, REGULAR TESTING HELPS MONITOR THE EFFECTIVENESS OF TREATMENT AND ENSURE APPROPRIATE LEVELS.
170	127	TESTOSTERONE (FEMALE 17 YEARS)	DELAYED ONE STEP IMMUNO ASSAY CMIA	SERUM	SERUM CONTAINER	2 Hours	IN FEMALES, TESTOSTERONE IS PRIMARILY PRODUCED BY THE OVARIES AND ADRENAL GLANDS. IT CONTRIBUTES TO THE REGULATION OF THE MENSTRUAL CYCLE, HELPS MAINTAIN BONE DENSITY, AND HAS AN IMPACT ON LIBIDO AND OVERALL ENERGY LEVELS. TESTOSTERONE IS TESTED IN FEMALES TO ASSESS HORMONAL IMBALANCES, FERTILITY ISSUES, ANDROGEN-RELATED DISORDERS, ETC.
171	125	TESTOSTERONE (MALE-17 YEARS)	DELAYED ONE STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	TESTING TESTOSTERONE LEVELS IN MALES IS A COMMON MEDICAL PRACTICE USED TO ASSESS THE FUNCTIONING OF THE MALE REPRODUCTIVE SYSTEM AND OVERALL HEALTH. TESTOSTERONE TESTING IN MALES MIGHT BE CONSIDERED IN CASES OF HORMONE IMBALANCES, DELAYED OR PRECOCIOUS PUBERTY, INFERTILITY, MONITORING HORMONE REPLACEMENT THERAPY. GYNACOMASTIA, ETC.
172	972	THYROGLOBULIN	CLIA	SERUM	SERUM CONTAINER	3 Hours	THE THYROGLOBULIN TEST IS OFTEN USED TO MONITOR PATIENTS WHO HAVE BEEN TREATED FOR THYROID CANCER. AFTER THE THYROID GLAND IS REMOVED (THYROIDECTOMY) OR TREATED WITH RADIOACTIVE IODINE, THERE SHOULD BE MINIMAL THYROGLOBULIN IN THE BLOOD.
173	25	TOTAL CHOLESTEROL	ENZYMATIC (CHOLESTROL ESTERASE AND CHOLESTROL OXIDASE)	SERUM	SERUM CONTAINER	2 Hours	A TOTAL CHOLESTEROL TEST MEASURES THE SUM OF GOOD AND BAD CHOLESTEROL. WHILE TOTAL CHOLESTEROL CAN BE TESTED ON ITS OWN, IT IS MORE OFTEN INTEGRATED INTO THE LIPID PANEL TEST THAT ALSO SHOWS THE LEVELS OF EACH TYPE OF CHOLESTEROL..

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
174	653	TOTAL CHOLESTEROL/ HDL RATIO	CALCULATION	SERUM	SERUM CONTAINER	2 Hours	TEST IS USED TO ASSESS CARDIOVASCULAR RISK, AS HIGHER RATIOS ARE GENERALLY ASSOCIATED WITH A HIGHER RISK OF HEART DISEASE.
175	142	TOTAL PROTEIN	BIURET REACTION END POINT	SERUM	SERUM CONTAINER	2 Hours	TOTAL PROTEIN TESTING MEASURES THE OVERALL CONCENTRATION OF PROTEINS IN A BIOLOGICAL FLUID SAMPLE. TEST IS USUALLY DONE AS PART OF A COMPREHENSIVE METABOLIC PANEL TO ASSESS LIVER AND KIDNEY FUNCTION. CONDITIONS, SUCH AS INFECTIONS OR INFLAMMATORY DISORDERS, CAN LEAD TO CHANGES IN PROTEIN LEVELS.
176	91	TOTAL T3	TWO STEP IMMUNO ASSAY CMIA	SERUM	SERUM CONTAINER	2 Hours	A TEST THAT MEASURES BOTH BOUND AND FREE T3 IS CALLED A TOTAL T3 TEST. A T3 TEST IS MOST OFTEN USED TO DIAGNOSE HYPERTHYROIDISM. T3 TESTS ARE FREQUENTLY ORDERED WITH T4 AND TSH TESTS. A T3 TEST MAY ALSO BE USED TO MONITOR TREATMENT FOR THYROID DISEASE.
177	96	TOTAL T4	TWO STEP IMMUNO ASSAY CMIA	SERUM	SERUM CONTAINER	2 Hours	THYROXINE TEST IS A BLOOD TEST THAT HELPS DIAGNOSE THYROID CONDITIONS. THYROXINE, ALSO KNOWN AS T4, IS A TYPE OF THYROID HORMONE. A T4 TEST MEASURES THE LEVEL OF T4 IN YOUR BLOOD. TOO MUCH OR TOO LITTLE T4 CAN BE A SIGN OF THYROID DISEASE.
178	53	TRIGLYCERIDES	GLYCEROL PHOSPHATE OXIDASE	SERUM	SERUM CONTAINER	2 Hours	TRIGLYCERIDES ARE A TYPE OF FAT (LIPID) FOUND IN THE BLOODSTREAM AND STORED IN FAT CELLS. TRIGLYCERIDE TESTING IS USED FOR SEVERAL PURPOSES, INCLUDING-ASSESSING CARDIOVASCULAR RISK, HEART DISEASES AND STROKE, METABOLIC HEALTH, INSULIN RESISTANCE AND METABOLIC SYNDROME.
179	87	TRIGLYCERIDES FLUID	GLYCEROL PHOSPHATE OXIDASE	FLUID	AIR TIGHT CONTAINER	2 Hours	MEASUREMENT OF TRIGLYCERIDES COMBINED WITH DETECTION OF CHYLOMICRONS IN BODY FLUIDS IS USEFUL FOR DIAGNOSING CHYLOUS EFFUSION OR DIFFERENTIATING FROM PSEUDOCYLOUS EFFUSION.
180	881	TRIPLE MARKER SCREENING	CLIA	SERUM	SERUM CONTAINER	24 Hours	TRIPLE MARKER SCREENING, ALSO KNOWN AS TRIPLE TEST OR TRIPLE SCREEN, IS A PRENATAL SCREENING TEST PERFORMED DURING PREGNANCY TO ASSESS THE RISK OF CHROMOSOMAL ABNORMALITIES AND NEURAL TUBE DEFECTS IN THE DEVELOPING FETUS IN THE SECOND TRIMESTER. IT INVOLVES MEASURING THREE SPECIFIC SUBSTANCES IN THE MOTHER'S BLOOD: AFP,BETA-HCG AND ESTRIOL. THE TRIPLE MARKER SCREENING DOES NOT PROVIDE A DEFINITIVE DIAGNOSIS BUT RATHER HELPS IDENTIFY PREGNANCIES THAT MIGHT BE AT A HIGHER RISK.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
182	696	TROPONIN I	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	1 Hours	TROPONIN I, IS A SPECIFIC PROTEIN FOUND IN CARDIAC MUSCLE CELLS. THIS TEST IS PRIMARILY USED TO DIAGNOSE AND ASSESS THE SEVERITY OF HEART-RELATED CONDITIONS, PARTICULARLY HEART ATTACKS (MYOCARDIAL INFARCTIONS) AND OTHER FORMS OF CARDIAC DAMAGE.
183	95	TSH	TWO STEP IMMUNO ASSAY-CMIA	SERUM	SERUM CONTAINER	2 Hours	THE PRIMARY PURPOSE OF THE TSH TEST IS TO ASSESS THE FUNCTION OF THE THYROID GLAND. TSH LEVELS HELP DETERMINE WHETHER THE THYROID GLAND IS PRODUCING ADEQUATE AMOUNTS OF THYROID HORMONES (T3 AND T4). THE TEST IS USED TO DIAGNOSE THYROID DISORDERS SUCH AS HYPOTHYROIDISM AND HYPERTHYROIDISM AND MONITORING THYROID TREATMENT.
184	1035	U.I.B.C	FERENE	SERUM	SERUM CONTAINER	2 Hours	UNSATURATED IRON BINDING CAPACITY (UIBC) MEASURES THE AMOUNT OF IRON THAT CAN BE BOUND TO TRANSFERRIN. UIBC IS USED AS PART OF AN IRON STUDIES PANEL TO ASSESS IRON METABOLISM AND RELATED CONDITIONS.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
185	1121	UNCONJUGATED ESTRIOL (UE3)	CLIA	SERUM	SERUM CONTAINER	3 Hours	MATERNAL SERUM UNCONJUGATED ESTRIOL (MSUE3), IS A PRENATAL SCREENING TEST THAT MEASURES THE LEVEL OF UNCONJUGATED ESTRIOL IN THE MOTHER'S BLOOD DURING PREGNANCY. THIS TEST IS OFTEN PART OF THE QUAD SCREEN OR QUADRUPLE TEST.
186	54	UREA	UREASE; UV ABSORBANCE	SERUM	SERUM CONTAINER	2 Hours	THE BLOOD UREA TEST REVEALS IMPORTANT INFORMATION ABOUT HOW WELL YOUR KIDNEYS ARE WORKING. UREA IS ROUTINELY MEASURED AS PART OF A "UREA AND ELECTROLYTES" (U&E) SCREEN TO ASSESS KIDNEY FUNCTION. IT MAY ALSO BE TESTED IF YOU HAVE SYMPTOMS OF ACUTE OR CHRONIC KIDNEY DISEASE.
187	55	URIC ACID	URICASE	SERUM	SERUM CONTAINER	2 Hours	URIC ACID IS CREATED WHEN THE BODY BREAKS DOWN SUBSTANCES CALLED PURINES. PURINES ARE NORMALLY PRODUCED IN THE BODY AND ARE ALSO FOUND IN SOME FOODS AND DRINKS. A URIC ACID BLOOD TEST, ALSO KNOWN AS A SERUM URIC ACID MEASUREMENT, DETERMINES HOW MUCH URIC ACID IS PRESENT IN YOUR BLOOD. THE TEST CAN HELP DETERMINE HOW WELL YOUR BODY PRODUCES AND REMOVES URIC ACID.
188	1595	URIC ACID (FLUID)	URICASE	FLUID	AIR TIGHT CONTAINER	2 Hours	MEASURING URIC ACID LEVELS IN BODILY FLUIDS, CAN PROVIDE INSIGHTS INTO VARIOUS HEALTH CONDITIONS LIKE GOUT, WHICH IS CHARACTERIZED BY PAINFUL JOINT INFLAMMATION. HYPERURICEMIA IS ALSO ASSOCIATED WITH AN INCREASED RISK OF KIDNEY STONES.
189	135	URINE ALBUMIN CREATININE RATIO	VARIOUS METHODS	URINE	URINE CONTAINER	3 Hours	ALBUMIN IS NORMALLY FOUND IN THE BLOOD AND PLAYS A CRITICAL ROLE IN MAINTAINING THE BALANCE OF FLUIDS BETWEEN THE BLOOD VESSELS AND TISSUES. UNDER CERTAIN CONDITIONS, ALBUMIN CAN LEAK INTO THE URINE, WHICH IS TYPICALLY A SIGN OF KIDNEY DYSFUNCTION OR DAMAGE TO THE KIDNEY'S FILTERING UNITS (GLOMERULI). URINE ALBUMIN LEVELS ARE COMMONLY MEASURED TO ASSESS KIDNEY HEALTH AND FUNCTION.
190	72	URINE CALCIUM CREATININE RATIO	VARIOUS METHODS	URINE	URINE CONTAINER	3 Hours	THE RATIO IS CALCULATED AND EXPRESSED FOR THE AMOUNT OF CALCIUM AGAINST CREATINE RELEASED IN THE URINE BY THE KIDNEYS. IT IS THE TEST USED TO DETECT THE PRESENCE OF DISORDERS THAT LEAD TO EXCESS CALCIUM IN URINE.
191	81	URINE CALCIUM	ARSENazo III	URINE	URINE CONTAINER	3 Hours	A CALCIUM IN URINE TEST MEASURES THE AMOUNT OF CALCIUM IN YOUR URINE. IF YOUR URINE CALCIUM LEVELS ARE TOO HIGH OR TOO LOW, IT MAY BE A SIGN OF KIDNEY DISEASE, KIDNEY STONES, BONE DISEASE, A PARATHYROID GLAND DISORDER, OR OTHER CONDITIONS.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
192	84	URINE CREATININE	KINETIC ALKALINE PICRATE	URINE	URINE CONTAINER	3 Hours	CREATININE LEVELS CAN BE MEASURED THROUGH A URINE TEST. CREATININE IS A NORMAL WASTE PRODUCT THAT THE BODY PRODUCES EVERY DAY DURING MUSCLE MOVEMENTS AND WHEN DIGESTING MEAT. HEALTHY KIDNEYS REMOVE CREATININE FROM THE BLOOD, THROUGH URINE. ABNORMAL LEVELS OF CREATININE IN A PERSON'S URINE CAN INDICATE MEDICAL ISSUES, SUCH AS KIDNEY INFECTION, KIDNEY STONES, AND DIABETES.
193	77	URINE MICROALBUMIN	IMMUNOTURBIDIMETRY	URINE	URINE CONTAINER	3 Hours	A MICROALBUMIN URINE TEST CHECKS FOR SMALL (OR "MICRO") AMOUNTS OF ALBUMIN IN YOUR URINE, AT LEVELS SO SMALL A REGULAR URINE TEST MIGHT NOT FIND THEM. IT CAN BE A SIGN OF KIDNEY DISEASE.
194	85	URINE PHOSPHOROUS	PHOSPHO MOLYBDATE FORMATION	URINE	URINE CONTAINER	3 Hours	THE PHOSPHORUS URINE TEST IS USED TO CHECK THE AMOUNT OF PHOSPHATE PRESENT IN OUR BODY. IT IS DONE BECAUSE BOTH THE HIGH LEVEL AND LOW LEVEL OF PHOSPHORUS CAN BE DANGEROUS FOR OUR HEALTH.
195	69	URINE PROTEIN CREATININE RATIO	VARIOUS METHODS	URINE	URINE CONTAINER	3 Hours	THE UPCR TEST IS USED TO ASSESS HOW MUCH PROTEIN IS BEING FILTERED THROUGH THE KIDNEYS AND EXCRETED IN THE URINE. ELEVATED PROTEIN LEVELS IN THE URINE CAN BE INDICATIVE OF KIDNEY DAMAGE OR DYSFUNCTION.
196	131	URINE VMA (SPOT)	COLUMN-CHROMATOGRAPHY	URINE	URINE CONTAINER	24 Hours	URINE VMA SPOT TEST IS USED TO MEASURE THE LEVELS OF VANILLYLMADELIC ACID, A BREAKDOWN PRODUCT OF CERTAIN CATECHOLAMINES (EPINEPHRINE, NOREPINEPHRINE, AND DOPAMINE), IN THE URINE. THIS TEST IS PRIMARILY USED TO HELP DIAGNOSE CONDITIONS RELATED TO THE ADRENAL GLANDS AND CERTAIN TYPES OF TUMORS, SUCH AS NEUROBLASTOMA AND PHEOCHROMOCYTOMA.
197	726	VITAMIN B 12	TWO STEP IMMUNO ASSAY CMIA	SERUM	SERUM CONTAINER	2 Hours	THE B12 TEST IS USED FOR SEVERAL PURPOSES, INCLUDING: DIAGNOSING VITAMIN B12 DEFICIENCY, AND TO MONITOR THE B12 LEVELS IN SUPPLEMENTATION THERAPY. B12 LEVELS CAN BE TESTED TO ASSESS UNDERLYING CAUSES OF CONDITIONS, SUCH AS ANEMIA, NEUROPATHY, OR GASTROINTESTINAL DISORDERS.
198	133	VITAMIN D (25 HYDROXY)	DELAYED ONE STEP IMMUNO ASSAY CMIA	SERUM	SERUM CONTAINER	2 Hours	VITAMIN D IS A FAT-SOLUBLE VITAMIN THAT PLAYS A CRUCIAL ROLE IN MAINTAINING BONE HEALTH, SUPPORTING THE IMMUNE SYSTEM, AND AIDING IN THE ABSORPTION OF CALCIUM AND PHOSPHORUS. THE TEST HELPS TO UNDERSTAND WHETHER YOU HAVE SUFFICIENT LEVELS OF VITAMIN D TO SUPPORT VARIOUS BODILY FUNCTIONS.
199	57	VLDL CHOLESTEROL	CALCULATION	SERUM	SERUM CONTAINER	2 Hours	VLDL IS A TYPE OF LIPOPROTEIN THAT IS RESPONSIBLE FOR TRANSPORTING TRIGLYCERIDES, A TYPE OF FAT, FROM THE LIVER TO VARIOUS TISSUES IN THE BODY. ELEVATED VLDL CHOLESTEROL LEVELS ARE ASSOCIATED WITH AN INCREASED RISK OF CARDIOVASCULAR DISEASE. VLDL IS A PRECURSOR TO LDL (LOW-DENSITY LIPOPROTEIN) CHOLESTEROL, OFTEN REFERRED TO AS "BAD CHOLESTEROL. "VLDL CHOLESTEROL TESTING IS OFTEN PART OF A COMPREHENSIVE LIPID PROFILE.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
200	1738	17 ALPHA-HYDROXY PROGESTERONE (17-ALPHA OHP)	ELISA	SERUM	SERUM CONTAINER	24 Hours	17 OHP TEST HELP TO DIAGNOSE A GROUP OF UNCOMMON, INHERITED GENETIC DISORDERS THAT AFFECT HOW WELL THE ADRENAL GLANDS MAKE CORTISOL. THESE GENETIC ADRENAL GLAND DISORDERS ARE CALLED CONGENITAL ADRENAL HYPERPLASIA (CAH) .
201	704	24 HOURS URINE CALCIUM	ARSENazo III	24 HOUR URINE	URINE CONTAINER	3 Hours	THE 24-HOUR URINE CALCIUM TEST IS USED FOR VARIOUS PURPOSES, LIKE ASSESSING CALCIUM BALANCE, DIAGNOSING HYPERCALCEMIA, MONITORING KIDNEY STONES, ETC.
202	1676	24 HOURS URINE CORTISOL	CMIA	24 HOUR URINE	URINE CONTAINER	3 Hours	A CORTISOL TEST MEASURES THE LEVEL OF HORMONE- CORTISOL IN A 24-HOUR SAMPLE OF URINE. ABNORMAL CORTISOL LEVELS INDICATE PROBLEMS WITH THE ADRENAL GLANDS OR THE PITUITARY GLAND.
203	705	24 HOURS URINE CREATININE	KINETIC ALKALINE PICRATE	24 HOUR URINE	URINE CONTAINER	3 Hours	A 24-HOUR URINE CREATININE TEST IS A DIAGNOSTIC TEST THAT MEASURES THE AMOUNT OF CREATININE EXCRETED IN URINE OVER A 24-HOUR PERIOD. THE TEST HELPS ASSESS KIDNEY FUNCTION, GFR, CHRONIC KIDNEY DISEASE, ETC.
204	184	24 HOURS URINE PHOSPHOROUS	PHOSPHO MOLYBDATE FORMATION	24 HOUR URINE	URINE CONTAINER	3 Hours	THE 24-HOUR URINE PHOSPHORUS TEST IS PARTICULARLY USEFUL IN ASSESSING DISORDERS RELATED TO PHOSPHORUS BALANCE, SUCH AS KIDNEY DYSFUNCTION DISORDERS LIKE HYPERPARATHYROIDISM AND CERTAIN BONE DISEASES, METABOLIC CONDITIONS THAT CAN AFFECT PHOSPHORUS METABOLISM ETC.
205	188	24 HOURS URINE PROTEINS	BENZETHONIUM CHLORIDE	24 HOUR URINE	URINE CONTAINER	3 Hours	DETECTING ABNORMAL LEVELS OF PROTEIN IN THE URINE CAN PROVIDE IMPORTANT INFORMATION ABOUT KIDNEY FUNCTION AND POTENTIAL UNDERLYING MEDICAL CONDITIONS LIKE KIDNEY DISEASE, KIDNEY DAMAGE, OR OTHER SYSTEMIC DISORDERS.
206	185	24 HOURS URINE URIC ACID	URICASE	24 HOUR URINE	URINE CONTAINER	3 Hours	THE 24-HOUR URINE URIC ACID TEST IS PARTICULARLY USEFUL IN ASSESSING CONDITIONS RELATED TO URIC ACID IMBALANCE, INCLUDING GOUT, ASSESS THE RISK OF KIDNEY STONE FORMATION, KIDNEY DYSFUNCTION, ETC.
207	186	24 HOURS URINE V.M.A.	COLUMN-CHROMATOGRAPHY	24 HOUR URINE	URINE CONTAINER	24 Hours	VMA IS OF VALUE FOR EVALUATION OF HYPERTENSION, DIAGNOSIS AND FOLLOW UP OF NEUROBLASTOMA, GANGLIONEUROMA, AND GANGLIO NEUROBLASTOMA. MOST NEUROBLASTOMA PATIENTS EXCRETE EXCESS VMA IN 24-HOUR COLLECTIONS.

DEPT NAME: CLINICAL BIOCHEMISTRY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
208	1410	24 HR URINE POTASSIUM	ION-SELECTIVE ELECTRODE (INDIRECT)	24 HOUR URINE	URINE CONTAINER	3 Hours	POTASSIUM IS AN ESSENTIAL ELECTROLYTE THAT PLAYS A VITAL ROLE IN VARIOUS BODILY FUNCTIONS, INCLUDING MUSCLE CONTRACTIONS, NERVE IMPULSES, FLUID BALANCE, AND MAINTAINING PROPER HEART RHYTHM. MONITORING URINE POTASSIUM LEVELS CAN PROVIDE INSIGHTS INTO THE BODY'S POTASSIUM BALANCE AND KIDNEY FUNCTION.
209	1824	PROCALCITONIN	CLIA	SERUM	SERUM CONTAINER	4 Hours	PROCALCITONIN (PCT) IS A PROTEIN PRECURSOR OF CALCITONIN, MAINLY PRODUCED BY THYROID C-CELLS. DURING BACTERIAL INFECTIONS, ITS LEVELS RISE, MAKING IT A VALUABLE BIOMARKER FOR DETECTING BACTERIAL INFECTIONS, SEPSIS, OR RISK OF SEPSIS.
210	1826	NT-PRO BNP	ELFA	SERUM/ SODIUM OR LITHIUM HEPARIN PLASMA	SERUM CONTAINER/ SODIUM/ LITHIUM HEPARIN CONTAINER	4 Hours	NT-PROBNP IS A PROTEIN RELEASED BY THE HEART UNDER STRESS, COMMONLY DUE TO HEART FAILURE. ELEVATED LEVELS IN THE BLOOD HELP DIAGNOSE AND MONITOR HEART CONDITIONS, INDICATING THE HEART IS STRUGGLING TO PUMP EFFECTIVELY.
211	1831	MDMA	IMMUNO CHROMATOGRAPHY	URINE	URINE CONTAINER	3 Hours	MDMA (ECSTASY) IS A STIMULANT DRUG THAT ENHANCES CNS ACTIVITY AND SUPPRESSES APPETITE. COMMONLY ABUSED, HIGH DOSES CAN CAUSE HIGH BLOOD PRESSURE, DILATED PUPILS, OVERHEATING, SEIZURES, AND PSYCHOSIS. IT IS USUALLY DETECTABLE IN URINE FOR 1-3 DAYS.
212	1865	FREE TESTOSTERONE	CLIA	SERUM	SERUM CONTAINER	4 Hours	MEASUREMENT OF FREE TESTOSTERONE PERMITS THE ESTIMATION OF THE BIOLOGICALLY ACTIVE HORMONE. FREE TESTOSTERONE DETERMINATION IS RECOMMENDED TO OVERCOME THE INFLUENCES CAUSED BY VARIATIONS IN TRANSPORT PROTEINS ON THE TOTAL TESTOSTERONE CONCENTRATION.

CLINICAL PATHOLOGY

DEPT NAME: CLINICAL PATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
1	253	BENCE-JONES PROTEINS	MANUAL	URINE	URINE CONTAINER	3 Hours	BENCE-JONES PROTEINS ARE IMMUNOGLOBULIN LIGHT CHAINS FOUND IN URINE, IN CASE OF MULTIPLE MYELOMA, AMYLOIDOSIS AND SYSTEMIC LUPUS ERYTHEMATOSUS.
2	197	URINE COLOUR	MANUAL	URINE	URINE CONTAINER	3 Hours	NORMAL URINE COLOUR RANGES FROM CLEAR TO PALE YELLOW. CERTAIN CONDITIONS LIKE JAUNDICE, RENAL DISEASES, ETC, CAN CHANGE THE COLOUR OF URINE. URINE COLOUR IS ALSO INFLUENCED BY DIET AND MEDICATION.
3	201	APPEARANCE (URINE)	MANUAL	URINE	URINE CONTAINER	2 Hours	THE TECHNICIAN VISUALLY EXAMINES THE URINE'S COLOR AND CLARITY. THIS WILL PROVIDE INFORMATION ABOUT SUBSTANCES THAT MAY BE PRESENT IN URINE. THEY ARE CLINICALLY CORRELATED WITH RESULTS OBTAINED DURING THE CHEMICAL AND MICROSCOPIC EXAMINATIONS FOR CONFIRMATION.
4	1427	REACTION (PH)	REGNT STRIP (PH INDICATOR)	URINE	URINE CONTAINER	2 Hours	A URINE TEST STRIP OR DIPSTICK TEST IS A BASIC DIAGNOSTIC TOOL USED TO DETERMINE PATHOLOGICAL CHANGES IN A PATIENT'S URINE IN STANDARD URINE ANALYSIS. A STANDARD URINE TEST STRIP MAY COMPRISE OF UP TO 10 DIFFERENT CHEMICAL PADS OR REAGENTS WHICH REACT (CHANGE COLOR) WHEN IMMERSERD IN A URINE SAMPLE. URINE IS USUALLY SLIGHTLY ACIDIC (PH RANGE 4.7 – 7.5). IF PH IS LESS THAN 4.7, IT IS MORE ACIDIC AND IF PH IS MORE THAN 7.5, IT IS MORE ALKALINE.
5	193	SPECIFIC GRAVITY (URINE)	REGNT STRIP (IONEXCHANGE)	URINE	URINE CONTAINER	2 Hours	URINE SPECIFIC GRAVITY (SG) IS A MEASURE OF CONCENTRATION OF SOLUTES IN THE URINE. MEASUREMENT OF SPECIFIC GRAVITY PROVIDES INFORMATION REGARDING A PATIENT'S STATE OF HYDRATION OR DEHYDRATION.THIS TEST SIMPLY INDICATES URINE CONCENTRATION.
6	256	URINE ALBUMIN	RGT STRIP (PROTEIN ERROR OF INDICATOR)	URINE	URINE CONTAINER	2 Hours	ALBUMIN IS A PROTEIN FOUND IN THE BLOOD. BUT A DAMAGED KIDNEY REACH TO MORETHAN NORMAL AMOUNT OF ALBUMIN PASS INTO THE URINE.
7	257	URINE UROBILINOGEN	RGTSTRIP (AZO DYE)	URINE	URINE CONTAINER	2 Hours	UROBILINOGEN IS A COLORLESS PIGMENT THAT IS A BYPRODUCT OF BILIRUBIN BREAKDOWN IN THE INTESTINES AND IS FOUND IN NORMAL URINE. THE AMOUNT OF UROBILINOGEN IN URINE CAN INDICATE LIVER FUNCTION AND OVERALL HEALTH.MORE THAN NORMAL AMOUNT OF UROBILINOGEN INDICATE ABNORMAL LIVERFUNCTION&HEMOLYSIS.

DEPT NAME: CLINICAL PATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
8	239	BILE SALTS (URINE)	RGTSTRIP (AZO DYE)	URINE	URINE CONTAINER	2 Hours	THE PRESENCE OF BILE SALTS IN URINE IS AN INDICATOR OF LIVER PROBLEMS/OBSTRUCTIVE JAUNDICE.
9	979	BILE PIGMENTS (URINE)	RGTSTRIP (AZO DYE)	URINE	URINE CONTAINER	2 Hours	THIS TEST IS PERFORMED ON A SAMPLE OF URINE TO DETECT DETECT BILE PIGMENTS. A POSITIVE RESULT INDICATES LIVER DYSFUNCTION&JAUNDICE.
10	1560	KETONE BODIES (URINE)	RGTSTRIP (NITROPRUSIDE)	URINE	URINE CONTAINER	2 Hours	KETONURIA IS A MEDICAL CONDITION IN WHICH KETONE BODIES ARE RELEASED IN THE URINE. IT IS SEEN IN CONDITIONS WHERE THE BODY PRODUCES EXCESS KETONES AS AN INDICATION THAT IT IS USING AN ALTERNATIVE SOURCE OF ENERGY. IT IS SEEN DURING STARVATION OR MORE COMMONLY, IN TYPE I DIABETES MELLITUS.
11	1561	MICROSCOPY (URINE)	MANUAL MICROSCOPY	URINE	URINE CONTAINER	1 Hours	THIS TEST LOOKS AT A SAMPLE OF URINE UNDER THE MICROSCOPE. TO IDENTIFY THE CELLS FROM URINARY TRACT, BLOOD CELLS, CRYSTALS, BACTERIA, PARASITES, ETC. THIS TEST IS OFTEN USED TO CONFIRM THE FINDINGS OF OTHER TESTS OR TO ADD INFORMATION TO THE DIAGNOSIS.
12	235	PORPHOBILINOGEN (URINE)	SHWATS AND WATSONS	URINE	URINE CONTAINER	2 Hours	URINARY PORPHOBILINOGEN LEVELS ARE MEASURED IN THE EVALUATION OF NEUROVISCERAL PORPHYRIAS, NAMELY, ACUTE INTERMITTENT PORPHYRIA, VARIEGATE PORPHYRIA, AND HEREDITARY COPROPORPHYRIA.
13	689	URINE FOR EOSINOPHILS	MANUAL MICROSCOPY	URINE	URINE CONTAINER	3 Hours	EOSINOPHILS ARE WHITE BLOOD CELLS THAT NORMALLY DO NOT APPEAR IN URINE. THE PRESENCE OF EOSINOPHILS IN URINE IS OBSERVED IN ACUTE INTERSTITIAL NEPHRITIS, WHICH IS CAUSED BY AN ALLERGIC REACTION, TYPICALLY TO DRUGS.
14	203	URINE FOR FAT GLOBULES	SUDAN DYE	URINE	URINE CONTAINER	1 Hours	THIS TEST IS DONE TO DETERMINE IF FAT GLOBULES ARE RELEASED IN THE URINE, WHICH MAY BE CAUSED DUE TO KIDNEY DISORDERS.
15	695	URINE FOR PREGNANCY TEST	CARD METHOD	URINE	URINE CONTAINER	1 Hours	TO CHECK FOR PREGNANCY.

DEPT NAME: CLINICAL PATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
16	222	URINE ROUTINE EXAMINATION	RGTSTRIP(AZO DYE,NITROPRUSIDE,PH INDICATOR , IONEXCHANGE , PROTEIN ERROR OF INDICATOR , GLUCOSEOXIDASE)AND MICROSCOPY	URINE	URINE CONTAINER	2 Hours	A URINE ROUTINE EXAMINATION, ALSO KNOWN AS A URINALYSIS, IS A SIMPLE TEST THAT CHECKS YOUR URINE FOR SIGNS OF HEALTH ISSUES. IT CAN HELP DETECT MANY DISEASES EARLY, SUCH AS INFECTIONS, KIDNEY PROBLEMS, DIABETES, AND LIVER DISEASE
17	236	URINE COMPLETE EXAMINATION	RGTSTRIP(AZO DYE,NITROPRUSIDE,PH INDICATOR , IONEXCHANGE , PROTEIN ERROR OF INDICATOR , GLUCOSEOXIDASE)AND MICROSCOPY	URINE	URINE CONTAINER	2 Hours	A URINE COMPLETE EXAMINATION, ALSO KNOWN AS A URINALYSIS, IS A SIMPLE TEST THAT CHECKS YOUR URINE FOR SIGNS OF HEALTH ISSUES. IT CAN HELP DETECT MANY DISEASES EARLY, SUCH AS INFECTIONS, KIDNEY PROBLEMS, DIABETES, AND LIVER DISEASE
18	249	STOOL ROUTINE ANALYSIS	MANUAL MICROSCOPY	STOOL	STERILE AIR TIGHT CONTAINER	2 Hours	MICROSCOPIC EXAMINATION OF STOOL SAMPLES FOR DIFFERENT CELLS, ORGANISMS, CYSTS OR OVA ETC.
19	218	STOOL: REDUCING SUBSTANCES	MANUAL	STOOL	STERILE AIR TIGHT CONTAINER	2 Hours	THIS TEST IS USED TO DETECT THE UNABSORBED SUGAR IN STOOL. SUGARS LIKE FRUCTOSE, GLUCOSE, LACTOSE, AND GALACTOSE CAN BE IDENTIFIED IN STOOL SAMPLES WITH THIS TEST. THIS TEST IS USUALLY DONE TO CHECK FOR LACTOSE INTOLERANCE.
20	260	MOTION - OCCULT BLOOD	CARD METHOD	STOOL	STERILE AIR TIGHT CONTAINER	1 Hours	FECAL OCCULT BLOOD REFERS TO THE BLOOD IN FECES THAT IS NOT VISIBLY APPARENT. A FECAL OCCULT BLOOD TEST CHECKS FOR HIDDEN BLOOD IN STOOL.
21	1074	MOTION FAT GLOBULES	MANUAL	STOOL	STERILE AIR TIGHT CONTAINER	1 Hours	THIS TEST MEASURES THE AMOUNT OF FAT IN YOUR STOOL. HAVING EXCESSIVE FAT IN YOUR STOOL IS CALLED STEATORRHEA. IF YOU HAVE LARGE QUANTITIES OF FAT IN YOUR STOOL, IT MAY BE A SIGN THAT THE FOOD IS MOVING THROUGH YOUR DIGESTIVE SYSTEM WITHOUT BEING BROKEN DOWN AND ABSORBED PROPERLY.

DEPT NAME: CLINICAL PATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
22	254	MOTION MICROSCOPY	MANUAL MICROSCOPY	STOOL	STERILE AIR TIGHT CONTAINER	2 Hours	STOOL MICROSCOPY IS A DIAGNOSTIC TOOL FOR IDENTIFYING PARASITIC ORGANISMS INCLUDING PROTOZOA AND HELMINTHS. IT IS ALSO USEFUL FOR QUANTIFICATION OF FECAL LEUKOCYTES.
23	711	SEMEN ANALYSIS	MANUAL	SEMEN	STERILE AIR TIGHT CONTAINER	4 Hours	IT IS DONE TO HELP EVALUATE MALE FERTILITY. THIS ANALYSIS MEASURES THE SPERM COUNT, MOTILITY, MORPHOLOGY, VOLUME, FRUCTOSE LEVEL AND PH.
24	207	PCV (HAEMATOCRIT) - FLUID	CELL COUNTER (CAL. VALUE)	FLUID	STERILE AIR TIGHT CONTAINER	3 Hours	IT IS THE PERCENTAGE (VOL %) OF CELLS IN FLUID.
25	818	RBC COUNT (FLUID)	FLOW CYTOMETRY	FLUID	STERILE AIR TIGHT CONTAINER	3 Hours	TO DETERMINE THE RBC COUNT IN FLUID.
26	820	TC, DC (SYNOVIAL FLUID)	MANUAL	FLUID	STERILE AIR TIGHT CONTAINER	3 Hours	DETECTS THE TOTAL CELL COUNT AND DIFFERENTIAL COUNT IN SYNOVIAL FLUID.
27	194	BREAST MILK - DC	MANUAL	BREAST MILK	AIR TIGHT CONTAINER	3 Hours	FOR DETERMINING THE DIFFERENTIAL COUNT IN BREAST MILK.
28	221	CELL COUNT (CSF)	MANUAL	CSF	STERILE AIR TIGHT CONTAINER	3 Hours	DETERMINES THE TOTAL CELL COUNT AND DIFFERENTIAL COUNT IN CSF.
29	232	CELL COUNT (PERICARDIAL)	MANUAL	PERICARDIAL FLUID	STERILE AIR TIGHT CONTAINER	3 Hours	DETERMINES THE TOTAL CELL COUNT AND DIFFERENTIAL COUNT IN PERICARDIAL FLUID.

DEPT NAME: CLINICAL PATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
30	225	CELL COUNT (PLEURAL FLUID)	MANUAL	PLEURAL FLUID	STERILE AIR TIGHT CONTAINER	3 Hours	DETERMINES THE TOTAL CELL COUNT AND DIFFERENTIAL COUNT IN PLEURAL FLUID.
31	689	FLUID FOR CRYSTALS	MANUAL	FLUID	STERILE AIR TIGHT CONTAINER	3 Hours	DETECTS THE PRESENCE OF ANY TYPE OF CRYSTALS IN BODY FLUIDS.
32	691	FLUID PH	REGNT STRIP (PH INDICATOR)	FLUID	STERILE AIR TIGHT CONTAINER	2 Hours	FOR DETERMINING THE PH OF FLUID.
33	252	FLUID - TC, DC	MANUAL METHOD	FLUID	STERILE AIR TIGHT CONTAINER	3 Hours	DETERMINES TOTAL WBC COUNT AND DIFFERENTIAL COUNT IN BODY FLUIDS.
34	228	FLUID SPECIFIC GRAVITY	REGNT STRIP (IONEXCHANGE)	FLUID	STERILE AIR TIGHT CONTAINER	2 Hours	FOR DETERMINING THE SPECIFIC GRAVITY OF FLUID.
35	240	URINE FOR FAT GLOBULES	SUDAN DYE	URINE	URINE CONTAINER	1 Hours	THIS TEST IS DONE TO DETERMINE IF FAT GLOBULES ARE RELEASED IN THE URINE, WHICH MAY BE CAUSED DUE TO KIDNEY DISORDERS.
36	204	URINE FOR PREGNANCY TEST	CARD METHOD	URINE	URINE CONTAINER	1 Hours	TO CHECK FOR PREGNANCY.
37	353	URINE KETONES (ACETONE)	URINE REAGENT STRIP (NITROPRUSSIDE)	URINE	URINE CONTAINER	2 Hours	TESTING FOR THE PRESENCE OF KETONES IN URINE (KETONURIA) IS ASSISTIVE IN THE DIAGNOSIS OF DIABETES MELLITUS, AS WELL AS IN EVALUATING CONDITIONS ASSOCIATED WITH KETOACIDOTIC STATES, SUCH AS STARVATION.

DEPT NAME: CLINICAL PATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
38	227	URINE MICROSCOPY	MANUAL METHOD	URINE	URINE CONTAINER	1 Hours	THIS TEST LOOKS AT A SAMPLE OF URINE UNDER THE MICROSCOPE. IT CAN SEE CELLS FROM URINARY TRACT, BLOOD CELLS, CRYSTALS, BACTERIA, PARASITES, ETC. THIS TEST IS OFTEN USED TO CONFIRM THE FINDINGS OF OTHER TESTS OR TO ADD INFORMATION TO THE DIAGNOSIS.
39	247	URINE OCCULT BLOOD	BENZIDINE TEST	URINE	URINE CONTAINER	1 Hours	OCCULT BLOOD IS OBSERVED WHEN BLOOD IN URINE IS NOT VISIBLE TO THE NAKED EYE, AND IS ONLY VISIBLE UNDER A MICROSCOPE. IT IS SEEN IN CASE OF CANCERS, KIDNEY STONES, RENAL DISEASES, URINARY TRACT INFECTIONS, AND BENIGN PROSTATIC HYPERPLASIA.
40	231	URINE PH	URINE REAGENT STRIP (BENZIDINE METHOD)	URINE	URINE CONTAINER	1 Hours	THE DETERMINATION OF URINE PH PROVIDES INFORMATION REGARDING THE ACID-BASE STATUS OF THE PATIENT. LOWER URINE PH IS ASSOCIATED WITH DIABETIC KETOACIDOSIS, DIARRHEA, STARVATION, ETC. HIGHER URINE PH IS ASSOCIATED WITH URINARY TRACT INFECTIONS, METABOLIC ALKALOSIS, CHRONIC RENAL FAILURE, ETC.
41	243	URINE ROUTINE ANALYSIS	URINE REAGENT STRIP/MANUAL MICROSCOPY	URINE	URINE CONTAINER	2 Hours	THIS TEST IS USED TO DETECT AND MANAGE A WIDE RANGE OF DISORDERS, SUCH AS, URINARY TRACT INFECTIONS, KIDNEY DISEASES AND DIABETES. A URINE ANALYSIS INVOLVES CHECKING THE APPEARANCE, CONCENTRATION AND CONTENT OF URINE.
42	666	URINE SUGAR (0.5 HOUR)	URINE REAGENT STRIP (GLUCOSE OXIDASE METHOD)	URINE	URINE CONTAINER	1 Hours	NORMALLY THERE IS VERY LITTLE TO NO GLUCOSE IN URINE. DETECTABLE LEVELS OF GLUCOSE IN URINE MAY BE A SIGN OF DIABETES, THE MOST COMMON CAUSE OF HIGH GLUCOSE IN URINE.
43	211	URINE SUGAR (1 HOUR)	URINE REAGENT STRIP (GLUCOSE OXIDASE METHOD)	URINE	URINE CONTAINER	1 Hours	NORMALLY THERE IS VERY LITTLE TO NO GLUCOSE IN URINE. DETECTABLE LEVELS OF GLUCOSE IN URINE MAY BE A SIGN OF DIABETES, THE MOST COMMON CAUSE OF HIGH GLUCOSE IN URINE.
44	667	URINE SUGAR (1.5 HOURS)	URINE REAGENT STRIP (GLUCOSE OXIDASE METHOD)	URINE	URINE CONTAINER	1 Hours	NORMALLY THERE IS VERY LITTLE TO NO GLUCOSE IN URINE. DETECTABLE LEVELS OF GLUCOSE IN URINE MAY BE A SIGN OF DIABETES, THE MOST COMMON CAUSE OF HIGH GLUCOSE IN URINE.
45	212	URINE SUGAR (2 HOURS)	URINE REAGENT STRIP (GLUCOSE OXIDASE METHOD)	URINE	URINE CONTAINER	1 Hours	NORMALLY THERE IS VERY LITTLE TO NO GLUCOSE IN URINE. DETECTABLE LEVELS OF GLUCOSE IN URINE MAY BE A SIGN OF DIABETES, THE MOST COMMON CAUSE OF HIGH GLUCOSE IN URINE.

DEPT NAME: CLINICAL PATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
46	213	URINE SUGAR (2.5 HOURS)	URINE REAGENT STRIP (GLUCOSE OXIDASE METHOD)	URINE	URINE CONTAINER	1 Hours	NORMALLY THERE IS VERY LITTLE TO NO GLUCOSE IN URINE. DETECTABLE LEVELS OF GLUCOSE IN URINE MAY BE A SIGN OF DIABETES, THE MOST COMMON CAUSE OF HIGH GLUCOSE IN URINE.
47	215	URINE SUGAR (3 HOURS)	URINE REAGENT STRIP (GLUCOSE OXIDASE METHOD)	URINE	URINE CONTAINER	1 Hours	NORMALLY THERE IS VERY LITTLE TO NO GLUCOSE IN URINE. DETECTABLE LEVELS OF GLUCOSE IN URINE MAY BE A SIGN OF DIABETES, THE MOST COMMON CAUSE OF HIGH GLUCOSE IN URINE.
48	251	URINE SUGAR (4 HOURS)	URINE REAGENT STRIP (BENZIDINE METHOD)	URINE	URINE CONTAINER	1 Hours	NORMALLY THERE IS VERY LITTLE TO NO GLUCOSE IN URINE. DETECTABLE LEVELS OF GLUCOSE IN URINE MAY BE A SIGN OF DIABETES, THE MOST COMMON CAUSE OF HIGH GLUCOSE IN URINE.
49	209	URINE SUGAR (FASTING)	URINE REAGENT STRIP/MANUAL MICROSCOPY	URINE	URINE CONTAINER	1 Hours	NORMALLY THERE IS VERY LITTLE TO NO GLUCOSE IN URINE. DETECTABLE LEVELS OF GLUCOSE IN URINE MAY BE A SIGN OF DIABETES, THE MOST COMMON CAUSE OF HIGH GLUCOSE IN URINE.
50	220	URINE SUGAR (RANDOM)	URINE REAGENT STRIP (GLUCOSE OXIDASE METHOD)	URINE	URINE CONTAINER	1 Hours	NORMALLY THERE IS VERY LITTLE TO NO GLUCOSE IN URINE. DETECTABLE LEVELS OF GLUCOSE IN URINE MAY BE A SIGN OF DIABETES, THE MOST COMMON CAUSE OF HIGH GLUCOSE IN URINE.
51	217	UROBILINOGEN	URINE REAGENT STRIP (AZO DYE)	URINE	URINE CONTAINER	2 Hours	A SMALL AMOUNT OF UROBILINOGEN IS NORMALLY EXCRETED IN THE URINE. AN INCREASE IN UROBILINOGEN IS AN INDICATION OF HEPATIC DYSFUNCTION, OR A HEMOLYTIC PROCESS.
52	1633	VOMITUS MICROSCOPY	MANUAL/ MICROSCOPY	URINE	URINE CONTAINER	2 Hours	THE VOMITUS IS MICROSCOPICALLY EXAMINED FOR THE PRESENCE OF RBCS, WBCS, EPITHELIAL CELLS, PARASITES, ETC.



HEMATOLOGY



DEPT NAME: HEMATOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
1	752	L.E CELL EXAMINATION	MANUAL/ MICROSCOPY	WHOLE BLOOD	STERILE BOTTLE WITH GLASS BEEDS.	24 Hours	THE LE CELL TEST IS USED TO DIAGNOSE AN AUTOIMMUNE DISEASE CALLED SLE.
2	280	MANTOUX TEST	MANUAL			2 Hours	THE MANTOUX TUBERCULIN SKIN TEST (TST) IS ONE METHOD OF DETERMINING WHETHER A PERSON IS INFECTED WITH M. TUBERCULOSIS
3	262	OSMOTIC FRAGILITY TEST		WHOLE BLOOD	LITHIUM HEPARIN.	24 Hours	OSMOTIC FRAGILITY (OF) REFERS TO THE DEGREE OR PROPORTION OF HEMOLYSIS THAT OCCURS WHEN A SAMPLE OF RED BLOOD CELLS ARE SUBJECTED TO OSMOTIC STRESS BY PLACING IN A HYPOTONIC SOLUTION. IT IS HELPFUL IN DETECTING OR CONFIRMING SPHEROCYTOSIS .
4	746	BONE MARROW (RECEIVED FROM OUTSIDE)	MANUAL/ MICROSCOPY	BONE MARROW	SLIDE/AIR TIGHT CONTAINER	4- 5 Days	TO DETECT INFECTIONS, INFILTRATIVE NEOPLASTIC LESIONS OF THE MARROW, AND ABNORMALITIES OF HEMATOPOIESIS.
5	751	BONE MARROW (COLLECTED AT ASWINI)	MANUAL/ MICROSCOPY	BONE MARROW	SLIDE/AIR TIGHT CONTAINER	4- 5 Days	TO DETECT INFECTIONS, INFILTRATIVE NEOPLASTIC LESIONS OF THE MARROW, AND ABNORMALITIES OF HEMATOPOIESIS.
6	303	BONE MARROW (IMPRINT)	MANUAL/ MICROSCOPY	BONE MARROW	SLIDE/AIR TIGHT CONTAINER	4- 5 Days	TO DETECT INFECTIONS, INFILTRATIVE NEOPLASTIC LESIONS OF THE MARROW, AND ABNORMALITIES OF HEMATOPOIESIS.
7	769	SECOND OPINION BONE MARROW	MANUAL/ MICROSCOPY	BONE MARROW	SLIDE/AIR TIGHT CONTAINER	4- 5 Days	TO DETECT INFECTIONS, INFILTRATIVE NEOPLASTIC LESIONS OF THE MARROW, AND ABNORMALITIES OF HEMATOPOIESIS.
8	293	SICKLING TEST	MANUAL/ MICROSCOPY	WHOLE BLOOD	EDTA CONTAINER	24 Hours	A SICKLE CELL TEST IS A BLOOD TEST USED TO DETERMINE IF YOU HAVE SICKLE CELL DISEASE (SCD). PEOPLE WITH SCD HAVE RED BLOOD CELLS (RBCS) THAT ARE ABNORMALLY SHAPED.

DEPT NAME: HEMATOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
9	1157	BLOOD GROUP&RH	GEL CARD	WHOLE BLOOD	EDTA CONTAINER	2 Hours	THE BLOOD TYPING IS FOR THE DETECTION OF ABO AND RH ANTIGENS ON THE RED BLOOD CELLS IN AN INDIVIDUAL'S BLOOD.
10	286	RETICULOCYTE COUNT	MANUAL/ MICROSCOPY	WHOLE BLOOD	EDTA CONTAINER	3 Hours	RETICULOCYTES ARE RED BLOOD CELLS THAT ARE STILL DEVELOPING. THEY ARE ALSO KNOWN AS IMMATURE RED BLOOD CELLS. A RETICULOCYTE COUNT MEASURES THE NUMBER OF RETICULOCYTES IN THE BLOOD. IF THE COUNT IS TOO HIGH OR TOO LOW, IT CAN MEAN A SERIOUS HEALTH PROBLEM, INCLUDING ANEMIA AND DISORDERS OF THE BONE MARROW, LIVER, AND KIDNEYS.
11	715	PERIPHERAL SMEAR	MANUAL/ MICROSCOPY	WHOLE BLOOD	EDTA CONTAINER	24 Hours	THE PURPOSE OF EXAMINING A BLOOD SMEAR IS TO CHECK THE SIZE, SHAPE, AND THE NUMBER OF THREE TYPES OF BLOOD CELLS: RED BLOOD CELLS, WHITE BLOOD CELLS AND PLATELETS. A BLOOD SMEAR IS USED TO HELP DIAGNOSE AND MONITOR MANY CONDITIONS, SUCH AS BLOOD DISORDERS, SUDDEN KIDNEY FAILURE, AND TREATMENT FOR CERTAIN CANCERS.
12	719	PERIPHERAL SMEAR FOR MP	MANUAL/ MICROSCOPY	WHOLE BLOOD	EDTA CONTAINER	24 Hours	LIGHT MICROSCOPY OF THICK AND THIN BLOOD SMEARS REMAIN THE STANDARD METHOD FOR DIAGNOSING MALARIA. IT INVOLVES COLLECTION OF A BLOOD SMEAR, ITS STAINING WITH ROMANOWSKY STAIN AND EXAMINATION OF RED BLOOD CELLS FOR INTRACELLULAR MALARIAL PARASITES.
13	114	PERIPHERAL SMEAR FOR MICROFILARIA	MANUAL/ MICROSCOPY	WHOLE BLOOD	EDTA CONTAINER	24 Hours	THE TEST HELPS IN THE DETECTION OF MICROFILARIA IN THE BLOOD AND LYMPHATIC TISSUES. FILARIASIS IS A PARASITIC INFECTION RESULTING FROM THREAD LIKE WORMS.
14	291	BLEEDING TIME	MANUAL	WHOLE BLOOD		2 Hours	BLEEDING TIME IS A SCREENING TEST FOR DETECTING DISORDERS INVOLVING PLATELET FUNCTION AND VASCULAR (CAPILLARY) DEFECTS THAT INTERFERE WITH THE CLOTTING PROCESS.
15	285	CLOTTING TIME	MANUAL	WHOLE BLOOD		2 Hours	CLOTTING TIME IS THE TIME REQUIRED FOR A SAMPLE OF BLOOD TO COAGULATE IN VITRO UNDER STANDARD CONDITIONS.
16	281	COOMBS TEST - DIRECT	GEL CARD	WHOLE BLOOD	EDTA CONTAINER	2 Hours	DIRECT COOMBS TEST (ALSO CALLED DIRECT ANTIGLOBULIN TEST) DETECTS THE ANTIBODIES ATTACHED TO RBCS IN THE BLOOD. ITS OFTEN USED TO DIAGNOSE BLOOD-RELATED CONDITIONS SUCH AS AUTOIMMUNE HEMOLYTIC ANEMIA.

DEPT NAME: HEMATOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
17	289	COOMBS TEST - INDIRECT	GEL CARD	SERUM	SERUM CONTAINER	3 Hours	AN INDIRECT COOMBS TEST CAN BE USED TO DETERMINE WHETHER THERE ARE ANTIBODIES TO THE RH FACTOR IN THE MOTHER'S BLOOD. A NORMAL (NEGATIVE) RESULT MEANS THAT THE MOTHER HAS NOT DEVELOPED ANTIBODIES AGAINST THE FETUS BLOOD.
18	1567	ICT TITRE	GEL CARD	SERUM	SERUM CONTAINER	3 Hours	AN INDIRECT COOMBS TEST CAN BE USED TO DETERMINE WHETHER THERE ARE ANTIBODIES TO THE RH FACTOR IN THE MOTHER'S BLOOD. A NORMAL (NEGATIVE) RESULT MEANS THAT THE MOTHER HAS NOT DEVELOPED ANTIBODIES AGAINST THE FETUS BLOOD.
19	12	COMPLETE HAEMOGRAM(ESR, HAEMOGLOBIN,RB C COUNT, MCH,MCHC,MCV,M PV,PCT,PCV,PDW,R DW-CV,DIFFERENTIAL COUNT,TOTAL WBC COUNT)	WESTERGREN'S , COLOMETRIC , FLOW CYTOMETRY, CELL COUNTER CAL.VALUE ,AUTOMATION & DERIVED.	WHOLE BLOOD	EDTA CONTAINER	2 Hours	A COMPLETE HEMOGRAM TEST CAN HELP DIAGNOSE CONDITIONS SUCH AS: ANEMIA, HEMATOLOGICAL CANCERS, INFECTIONS, ACUTE HEMORRHAGIC STATES, ALLERGIES, AND IMMUNODEFICIENCIES.
20	57	COMPLETE CELL COUNT(C.C.C) (HAEMOGLOBIN,RB C COUNT, MCH,MCHC,MCV,M PV,PCT,PCV,PDW,R DW-CV,DIFFERENTIAL COUNT,TOTAL WBC COUNT)	COLOMETRIC , FLOW CYTOMETRY, CELL COUNTER CAL.VALUE ,AUTOMATION & DERIVED.	WHOLE BLOOD	EDTA CONTAINER	2 Hours	A COMPLETE BLOOD COUNT (CBC) IS USED TO FIND A WIDE RANGE OF CONDITIONS, INCLUDING ANEMIA, INFECTION AND LEUKEMIA.
21	287	ESR	WESTERGREN'S	WHOLE BLOOD	3.8% SODIUM CITRATE CONTAINER	2 Hours	AN ERYTHROCYTE SEDIMENTATION RATE (ESR) IS A BLOOD TEST THAT CAN INDICATE IF YOU HAVE AN INFLAMMATION IN YOUR BODY. INFLAMMATION IS YOUR IMMUNE SYSTEM'S RESPONSE TO INJURY, INFECTION, AND MANY TYPES OF CONDITIONS, INCLUDING IMMUNE SYSTEM DISORDERS, CERTAIN CANCERS, AND BLOOD DISORDERS.

DEPT NAME: HEMATOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
22	277	HAEMOGLOBIN	COLOMETRIC METHOD,(NON-CYANMETHHB)	WHOLE BLOOD	EDTA CONTAINER	2 Hours	A HEMOGLOBIN TEST MEASURES THE LEVEL OF HEMOGLOBIN IN YOUR BLOOD. HEMOGLOBIN IS A PROTEIN IN RED BLOOD CELLS, THAT CARRY OXYGEN FROM THE LUNGS TO THE REST OF YOUR BODY. LOW HEMOGLOBIN LEVELS MAY BE A SIGN OF: DIFFERENT TYPES OF ANEMIAS, THALASSEMIA, IRON DEFICIENCY, LIVER DISEASE, CANCER AND OTHER DISEASES. HIGH HEMOGLOBIN LEVELS MAY BE A SIGN OF: LUNG DISEASE, HEART DISEASE, POLYCYTHEMIA VERA, ETC.
23	283	TOTAL WBC COUNT	FLOW CYTOMETRY	WHOLE BLOOD	EDTA CONTAINER	2 Hours	A WHITE BLOOD CELL COUNT MEASURES THE NUMBER OF WHITE CELLS IN YOUR BLOOD. WHITE BLOOD CELLS ARE PART OF THE IMMUNE SYSTEM. THEY HELP YOUR BODY FIGHT OFF INFECTIONS AND OTHER DISEASES. WHEN YOU GET SICK, YOUR BODY MAKES MORE WHITE BLOOD CELLS TO FIGHT THE BACTERIA, VIRUSES, OR OTHER FOREIGN SUBSTANCES CAUSING YOUR ILLNESS. THIS INCREASES YOUR WHITE BLOOD COUNT. SOME DISEASES CAN CAUSE A REDUCTION IN YOUR WBC COUNT, FOR E.G: SOME TYPES OF CANCER AND HIV/AIDS, A VIRAL DISEASE THAT ATTACKS WHITE BLOOD CELLS.
24	276	RBC COUNT	FLOW CYTOMETRY	WHOLE BLOOD	EDTA CONTAINER	2 Hours	A RED BLOOD CELL (RBC) COUNT MEASURES THE NUMBER OF RED BLOOD CELLS, ALSO KNOWN AS ERYTHROCYTES, IN YOUR BLOOD. A LOW RED BLOOD CELL COUNT CAN BE A SIGN OF: ANEMIA, LEUKEMIA, MALNUTRITION, MULTIPLE MYELOMA, OR KIDNEY FAILURE. A HIGH RED BLOOD CELL COUNT CAN BE A SIGN OF: DEHYDRATION, HEART DISEASE, POLYCYTHEMIA VERA, ETC.
25	278	PCV	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	THE PACKED CELL VOLUME (PCV) IS A MEASUREMENT OF THE PROPORTION OF BLOOD THAT IS MADE UP OF CELLS. THE PCV INCREASES WHEN THE NUMBER OF RED BLOOD CELLS INCREASE OR WHEN THE TOTAL BLOOD VOLUME IS REDUCED, AS IN DEHYDRATION. A FALL IN PCV BELOW NORMAL RANGE INDICATES ANAEMIA, WHEN YOUR BODY DECREASES ITS PRODUCTION OF RED BLOOD CELLS OR INCREASES ITS DESTRUCTION OF RED BLOOD CELLS. PCV IS ALSO LOWERED IF THE BLOOD CELLS ARE DILUTED BY AN INCREASE IN TOTAL VOLUME (HAEMODILUTION, OCCURRING DURING PREGNANCY).
26	288	PLATELET COUNT	FLOW CYTOMETRY	WHOLE BLOOD	EDTA CONTAINER	2 Hours	A PLATELET COUNT IS MOST OFTEN USED TO MONITOR OR DIAGNOSE CONDITIONS THAT CAUSE EXCESSIVE BLEEDING OR ABNORMAL CLOTTING.

DEPT NAME: HEMATOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
27	156	DIFFERENTIAL COUNT(NEUTROPHILS,LYMPHOCYTES, MONOCYTES,EOSINOPHILS,BASOPHILS & ABSOLUTE NEUTROPHIL COUNT,ABSOLUTE LYMPHOCYTE COUNT, ABSOLUTE MONOCYTE COUNT,ABSOLUTE EOSINOPHIL COUNT & ABSOLUTE BASOPHILL COUNT)	FLOW CYTOMETRY+LEISHMAN'S STAIN+ CELL COUNTER CAL.VALUE	WHOLE BLOOD	EDTA CONTAINER	2 Hours	A DIFFERENTIAL COUNT TEST MEASURES THE PERCENTAGE OF EACH TYPE OF WHITE BLOOD CELL (WBC) IN YOUR BLOOD. IT CAN HELP DIAGNOSE OR MONITOR CONDITIONS THAT AFFECT YOUR IMMUNE SYSTEM, SUCH AS INFECTIONS, ANEMIA, OR LEUKEMIA. ALONG WITH THE PERCENTAGE, WE ALSO CALCULATE THE ABSOLUTE COUNT OF EACH TYPE OF WBC TO GET MORE DETAILS.
28	669	MCH	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	MEAN CORPUSCULAR HEMOGLOBIN (MCH) IS A MEASURE OF THE AVERAGE AMOUNT OF HEMOGLOBIN IN EACH RED BLOOD CELL. THE PURPOSE OF MCH TEST IS TO CALCULATE THE AMOUNT OF HEMOGLOBIN IN AN INDIVIDUAL'S RED BLOOD CELLS. THE MCH TEST IS ONE OF THE SEVERAL TESTS THAT ARE USED TO DIAGNOSE AND CLASSIFY DIFFERENT TYPES OF ANEMIA.
29	282	MCHC	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC) MEASURES THE AVERAGE HEMOGLOBIN CONCENTRATION IN A GIVEN VOLUME OF RED BLOOD CELLS. THE PURPOSE OF AN MCHC TEST IS TO EVALUATE WHETHER RBCS ARE CARRYING AN APPROPRIATE AMOUNT OF HEMOGLOBIN. MCHC IS ONE OF THE SEVERAL TESTS USED TO ASSESS THE FUNCTION AND HEALTH OF RBCS TO CHECK FOR SIGNS OF ANEMIA AND OTHER BLOOD DISORDERS.
30	272	MCV	DERIVED	WHOLE BLOOD	EDTA CONTAINER	2 Hours	MCV STANDS FOR MEAN CORPUSCULAR VOLUME. AN MCV BLOOD TEST MEASURES THE AVERAGE SIZE OF YOUR RED BLOOD CELLS, ALSO KNOWN AS ERYTHROCYTES. MCV IS THE MOST USEFUL VALUE USED TO CLASSIFY THE TYPE OF ANEMIA BASED ON RED CELL MORPHOLOGY.
31	270	RDW - CV	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	THE RDW VALUE INDICATES WHETHER ENOUGH RED BLOOD CELLS ARE OF NORMAL SIZE AND SHAPE.THE RDW-CV IS A CALCULATION BASED ON BOTH THE WIDTH OF THE DISTRIBUTION CURVE AND THE MEAN CELL SIZE.

DEPT NAME: HEMATOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
32	278	MPV	DERIVED	WHOLE BLOOD	EDTA CONTAINER	2 Hours	MPV STANDS FOR MEAN PLATELET VOLUME. AN MPV BLOOD TEST MEASURES THE AVERAGE SIZE OF YOUR PLATELETS. THE TEST CAN HELP DIAGNOSE BLEEDING DISORDERS AND DISEASES OF THE BONE MARROW.
33	273	PDW	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	PDW REFLECTS HOW UNIFORM THE PLATELETS ARE IN SIZE. A NORMAL PDW INDICATES PLATELETS THAT ARE MOSTLY THE SAME SIZE, WHILE A HIGH PDW MEANS THAT PLATELET SIZE VARIES GREATLY, WHICH MAY RESULT FROM INCREASED ACTIVATION, DESTRUCTION AND/OR CONSUMPTION OF PLATELETS.
34	274	PCT	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	PCT IS THE VOLUME OCCUPIED BY PLATELETS IN THE BLOOD.
35	78	DIFF.WBCCOUNT	FLOW CYTOMETRY+LEISHMAN'S STAIN+CELL COUNTER CAL.VALUE	WHOLE BLOOD	EDTA CONTAINER	2 Hours	INCLUDES TOTAL WBC COUNT AND DIFFERENTIAL COUNT. IT IS USED TO HELP DIAGNOSE CONDITIONS LIKE: INFECTIONS, IMMUNODEFICIENCY DISORDERS, ETC.
36	292	ABSOLUTE EOSINOPHIL COUNT	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	AN ABSOLUTE EOSINOPHIL COUNT IS A BLOOD TEST THAT MEASURES THE EXACT NUMBER OF EOSINOPHILS, WHICH IS A TYPE OF WBC. EOSINOPHILS BECOMES ACTIVE WHEN YOU HAVE CERTAIN ALLERGIC DISEASES, INFECTIONS, AND OTHER MEDICAL CONDITIONS.
37	296	ABSOLUTE LYMPHOCYTE COUNT	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	AN ABSOLUTE LYMPHOCYTE COUNT TELLS YOU THE NUMBER OF LYMPHOCYTES AS AN ABSOLUTE NUMBER INSTEAD OF PERCENTAGE.
38	232	ABSOLUTE MONOCYTE COUNT	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	AN ABSOLUTE MONOCYTE COUNT IDENTIFIES HOW MANY MONOCYTES ARE PRESENT IN A SAMPLE OF BLOOD.
39	1731	ABSOLUTE BASOPHIL COUNT	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	AN ABSOLUTE BASOPHIL COUNT (ABC) IDENTIFIES HOW MANY BASOPHILS SPECIFICALLY ARE PRESENT IN A BLOOD SAMPLE.
40	1731	ABSOLUTE NEUTROPHIL COUNT	AUTOMATED CELL COUNTER	WHOLE BLOOD	EDTA CONTAINER	2 Hours	AN ABSOLUTE NEUTROPHIL COUNT DETERMINES HOW MANY NEUTROPHILS ARE PRESENT IN A SAMPLE OF BLOOD.

**HISTOPATHOLOGY
AND
CYTOPATHOLOGY**

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
1	858	ASCITIC FLUID CYTOLOGY	PAP STAINING/ MICROSCOPY	ASCITIC FLUID	STERILE CONTAINER	24-48 Hours	
2	337	BONE MARROW TREPINE	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN BOUIN'S FLUID CONTAINING CONTAINER	4-5 Days	
3	865	BRONCHIAL LAVAGE FOR CYTOLOGY	PAP,H&E STAINING/ MICROSCOPY	BRONCHIAL LAVAGE	STERILE CONTAINER	24-48 Hours	
4	1276	BRUSHINGS FOR CYTOLOGY	MANUAL/ MICROSCOPY	BRUSHINGS FLUID	TIGHT CONTAINER	24-48 Hours	
5	866	BUCCAL SMEAR	PAP STAINING/ MICROSCOPY	BUCCAL MUCOSA SMEAR	STERILE CONTAINER	24-48 Hours	
6	1334	CELL BLOCK PREPARATION	H&E STAINING/ MICROSCOPY	FLUID/ TISSUE	TIGHT CONTAINER	4-5 Days	
7	867	CSF CYTOLOGY	PAP STAINING/ MICROSCOPY	CSF	STERILE CONTAINER	24-48 Hours	
8	859	CYTOLOGY (FLUID)	PAP, H&E STAINING/ MICROSCOPY	FLUID	STERILE CONTAINER	24-48 Hours	

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
9	860	CYTOLOGY (PLEURAL FLUID)	PAP STAINING/ MICROSCOPY	PLEURAL FLUID	STERILE CONTAINER	24-48 Hours	
10	861	CYTOLOGY FOR MALIGNANT CELL (PERICARDIAL)	PAP STAINING/ MICROSCOPY	PERICARDIAL FLUID	STERILE CONTAINER	24-48 Hours	
11	862	CYTOLOGY FOR MALIGNANT CELL (PERITONEAL FLUID)	PAP STAINING/ MICROSCOPY	PERITONEAL FLUID	STERILE CONTAINER	24-48 Hours	
12	863	CYTOLOGY FOR MALIGNANT CELL (SYNOVIAL)	PAP STAINING/ MICROSCOPY	SYNOVIAL FLUID	STERILE CONTAINER	24-48 Hours	
13	868	DISCHARGE FOR MALIGNANT CELLS	PAP,H&E STAINING/ MICROSCOPY	SMEAR	STERILE CONTAINER	24-48 Hours	
14	313	FNAC (RECEIVED OUTSIDE)	PAP,H&E STAINING/ MICROSCOPY	ASPIRATE	CYTO FIX JAR/ DRIED SMEAR	24-48 Hours	
15	314	FNAC (TAKEN FROM ASWINI)	PAP,H&E, MGG STAINING/ MICROSCOPY	ASPIRATE	CYTO FIX JAR/ DRIED SMEAR	24-48 Hours	
16	316	HPE (2 SAMPLES/ MEDIUM)	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
17	1480	HPE (4-8 SAMPLES)	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	
18	340	HPE (LARGE)	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	
19	311	HPE (S)	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 Days	
20	1468	HPE ENDOSCOPY	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 Days	
21	1477	HPE EYEBALL	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	
22	1475	HPE FIBROADENOMA	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 Days	
23	1472	HPE GALL BLADDER	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 Days	
24	1482	HPE GASTRECTOMY WITH LYMPH NODE	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
25	1483	HPE HEMICOLECTOMY WITH LYMPHNODE	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	
26	1476	HPE LIPOMA	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 Days	
27	1485	HPE LUNG WITH LYMPH NODE	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	
28	1479	HPE LUNGS	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	
29	1474	HPE LYMPH NODE	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 Days	
30	1484	HPE MAXILLA/ MANDIBLE	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	Upto 20 Days	
31	1478	HPE NEPHRECTOMY	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	
32	1486	HPE RADICAL NECK DISSECTION	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
33	1487	HPE RADICAL NEPHRECTOMY WITH MALIGNANCY	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	
34	1469	HPE SKIN BIOPSY	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 Days	
35	1497	HPE SPECIAL STAIN ALCIAN BLUE	ALCIAN BLUE STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	24-48 Hours	THE ALCIAN BLUE STAIN IS MOST COMMONLY USED ON TISSUE SAMPLES OBTAINED FROM THE GASTROINTESTINAL (GI) TRACT AND IS USEFUL IN DIAGNOSING PATHOLOGICAL PROCESSES SUCH AS BARRETT'S ESOPHAGUS.
36	1491	HPE SPECIAL STAIN CONGO RED	CONGO RED STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	24-48 Hours	STAINING WITH CONGO RED (CR) IS A QUALITATIVE METHOD USED FOR THE IDENTIFICATION OF AMYLOIDS IN VITRO AND IN TISSUE SECTIONS.
37	1490	HPE SPECIAL STAIN GIEMSA	GIEMSA STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	24-48 Hours	GIEMSA STAIN IS PERFORMED ON PARAFFIN SECTIONS TO STAIN THE BLOOD CELLS OF HEMATOPOIETIC TISSUES. IT CAN ALSO BE APPLIED TO ALL TISSUE SECTIONS IN WHICH THE PRESENCE OF MICROORGANISMS IS SUSPECTED, ESPECIALLY PLASMODIUM.
38	1488	HPE SPECIAL STAIN MAY GRUNWALD	MAY GRUNWALD GIEMSA STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	24-48 Hours	MAY GRUNWALD-GIEMSA (MGG) STAINING METHOD IS USED FOR MORPHOLOGICAL INSPECTION AND DIFFERENTIAL COUNTING OF BLOOD CELLS IN AIR DRIED SMEARS.
39	1496	HPE SPECIAL STAIN METHYLENE BLUE	METHYLENE BLUE STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	24-48 Hours	METHYLENE BLUE IS USED FOR STAINING ACIDIC CELL COMPONENTS SUCH AS THE NUCLEUS, BACTERIA, AND BLOOD TISSUE SPECIMENS. IT IS ALSO USED TO EXAMINE THE FORMS, STRUCTURES, AND LOCATIONS OF DEAD CELLS AND OTHER TISSUES.
40	1489	HPE SPECIAL STAIN PAS	PAS STAINING/ MICROSCOPY	TISSUE/ SMEAR	10% FORMALIN CONTAINING CONTAINER	24-48 Hours	PERIODIC ACID SCHIFF (PAS) IS A SPECIAL STAIN PRIMARILY USED TO DETERMINE IF A SUBSTANCE INSIDE THE CELL IS GLYCOGEN OR A TYPE OF PROTEIN CALLED MUCIN.

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
41	1494	HPE SPECIAL STAIN PERL'S PRUSSIAN BLUE	PERL'S PRUSSIAN BLUE STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	24-48 Hours	THIS STAINING IS WIDELY USED TO DETECT THE PRESENCE OF IRON AND ANALYSE IRON DEFICIENCY IN BIOPSY SPECIMENS, ESPECIALLY IN TISSUES SUCH AS BONE MARROW AND SPLEEN.
42	1495	HPE SPECIAL STAIN TOLUEDENE BLUE	TOLUEDENE BLUE STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	24-48 Hours	TOLUIDINE BLUE HAS BEEN USED IN VIVO TO IDENTIFY DYSPLASIA AND CARCINOMA OF THE ORAL CAVITY. TOLUIDINE BLUE CAN HIGHLIGHT COMPONENTS, SUCH AS MAST CELLS GRANULES, MUCINS, AND CARTILAGE.
43	1493	HPE SPECIAL STAIN WADE FITE	WADE FITE STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	24-48 Hours	THE WADE FITE SPECIAL STAIN IS A MODIFICATION OF ZIEHL-NEELSEN METHOD. THIS IS USED TO DEMONSTRATE M LEPRAE WHICH IS MUCH LESS ACID-AND ALCOHOL-FAST THAN THE TUBERCLE BACILLUS.
44	1492	HPE SPECIAL STAIN ZIEHL NEILSON	ZIEHL-NEILSON STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	24-48 Hours	THE ZIEHL-NEELSEN METHOD USES A CARBOL FUCHSIN STAIN, ACID ALCOHOL DECOLORIZER, AND METHYLENE BLUE COUNTERSTAIN. ACID-FAST ORGANISMS STAIN RED, WHILE THE BACKGROUND OF DEBRIS STAINS BLUE. THE ZN STAIN CONFIRMS THE ACID-FAST PROPERTY OF MYCOBACTERIA.
45	1473	HPE TONSILS	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 Days	
46	1470	HPE TRUCUT	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 Days	
47	1471	HPE TURP	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 Days	
48	1481	HPE UTERUS (LARGE)	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 Days	

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
49		IHC-ACTIN	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. ACTIN IS A SMOOTH MUSCLE MARKER AND IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF SMOOTH MUSCLE TUMOR.
50	1041	IHC-ANY 1 MARKER	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING.
51	1416	IHC-ANY 2 MARKER	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING.
52	903	IHC-ANY 3 MARKER	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING.
53	1403	IHC-ANY 4 MARKER	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING.
54	1500	IHC-ANY 5 MARKER	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING.

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
55	1404	IHC-ANY 6 MARKER	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING.
56		IHC-BCL-2	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. BCL-2 STANDS FOR B-CELL LYMPHOMA 2 AND IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
57		IHC-BCL-6	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. BCL-6 STANDS FOR B-CELL LYMPHOMA 6 AND IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
58		IHC-CALDESMON	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CALDESMON IS A SMOOTH MUSCLE MARKER AND IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF SMOOTH MUSCLE TUMOR.
59		IHC-CALRETININ	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIALIZED STAINING TECHNIQUE FOR TISSUE SAMPLES, AIDING IN PRECISE DISEASE DIAGNOSIS, ESPECIALLY CANCERS. IT IDENTIFIES SPECIFIC CELLULAR MARKERS TO DISTINGUISH DISEASE TYPES AND SUBTYPES. CRUCIAL FOR TREATMENT PLANNING, CALRETININ IS KEY IN ANALYZING AND SELECTING TREATMENT FOR MESOTHELIOMA.

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
60		IHC-CD-1A	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD1A IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
61		IHC-CD-3	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD3 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
62		IHC-CD-4	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD4 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
63		IHC-CD-5	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD5 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
64		IHC-CD-8A	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD8A IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
65		IHC-CD-10	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD10 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
66		IHC-CD-15	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD15 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
67		IHC-CD-20	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD20 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
68		IHC-CD-23	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD23 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
69		IHC-CD-30	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSAN SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD30 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA,

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
70		IHC-CD-34	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD34 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
71		IHC-CD-45	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD45 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA,
72		IHC-CD-68	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD68 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LYMPHOMA.
73		IHC-CD-99	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD99 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF EWING SARCOMA (CANCER THAT OCCURS IN BONES OR IN SOFT TISSUES AROUND THE BONES).
74		IHC-CD-117	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CD117 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF GASTRO-INTESTINAL STROMAL TUMOR.
75		IHC-CD-138	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC STAINS TISSUE SAMPLES TO DIAGNOSE DISEASES, ESPECIALLY CANCERS, BY IDENTIFYING CELL MARKERS. CD138 HELPS ANALYZE AND TREAT MYELOMA.

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
76		IHC-CDX2	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PRECESS PERFORMED EN TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CDX2 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF COLORECTAL CANCER.
77		IHC-CK-5/6	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CK5/6 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LUNG CANCER.
78		IHC-CK-7	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CK7 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LUNG CANCER AND MESOTHELIOMA.
79		IHC-CK-20	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MERE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CK20 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF COLORECTAL CANCER.
80		IHC-CYCLIN D1	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISOLY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CYCLIN D1 IN USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF COLORECTAL CANCER AND LYMPHARMA.

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
81		IHC CYTOKERATIN AE1/3	IMMUNO- HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY, IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. CYTOKERATIN AE1/AE3 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF CANCERS WITH EPITHELIAL ORIGIN.
82		IHC-DESMIN	IMMUNO- HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. DESMIN IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF SMOOTH MUSCLE TUMOR.
83		IHC-DOG-1	IMMUNO- HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. DOG1 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF GASTRO INTESTINAL STROMAL TUMOR.
84		IHC-E-CADHERIN	IMMUNO- HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. E-CADHERIN IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF BREAST CANCER.
85		IHC-ER	IMMUNO- HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL TISSUE STAINING PROCESS FOR PRECISE DISEASE DIAGNOSIS. IHC HELPS DISTINGUISH DISEASE TYPES, ESPECIALLY CANCERS, BY IDENTIFYING CELLULAR MARKERS. IT PLAYS A KEY ROLE IN TREATMENT PLANNING, SUCH AS USING THE ESTROGEN RECEPTOR (ER) FOR BREAST CANCER ANALYSIS AND THERAPY SELECTION.

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
86		IHC-GFAP	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY.IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS.IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. GFAP(GLIAL FIBRILLARY ACIDIC PROTEIN) IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF GLIOMA (BRAIN TUMOR).
87		IHC-HER2	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. HER2(HUMAN EPIDERMAL GROWTH FACTOR RECEPTOR 2) IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF BREAST CANCER.
88		IHC HMB-45	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. HMB-45(HUMAN MELANOMA BLACK-45) IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF MELANOMA.
89		IHC-IgG4	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY, IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. IGG40MMUNOGLOBULIN 64) IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF IGG4 RELATED SYSTEMIC DISEASES.
90		IHC-INHIBIN	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNESE DISEASE MORE PRECISELY, IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. INHIBIN IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF OVARIAN CANCER.

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
91		IHC-KI67	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. K167 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF BREAST CANCER AND LYMPHOMA.
92		IHC-MYO D1	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A STAINING PROCESS FOR PRECISE DISEASE DIAGNOSIS, DISTINGUISHING CANCER TYPES BY IDENTIFYING CELLULAR MARKERS. IT AIDS TREATMENT PLANNING, WITH MYOD1 USED FOR ANALYZING AND GUIDING THERAPY FOR RHABDOMYOSARCOMA, A CANCER AFFECTING SKELETAL MUSCLE OR HOLLOW ORGANS.
93		IHC-MYOGENIN	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. MYOGENIN IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF RHABDOMY OSARCOMA (CANCER THAT FORMS IN SOFT TISSUE, SPECIFICALLY SKELETAL MUSCLE TISSUE OR SOMETIMES HOLLOW ORGANS SUCH AS UTERUS OR BLADDER).
94		IHC-P-53	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS.IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. P53 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF PROSTATE. CANCER, LUNG CANCER AND OVARIAN CANCER.
95		IHC-P-63	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNESE DISEASE MORE PRECISELY, IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. INHIBIN IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF OVARIAN CANCER.

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
96		IHC-PR	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. PR (PROGESTERONE RECEPTOR) IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF BREAST CANCER.
97		IHC-S-100	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. \$100 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF MELANOMA.
98		IHC-SYNAPTO- PHYSIN	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. SYNAPTOPHYSIN IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF NEUROENDOCRINE TUMORS.
99		IHC-TLE	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. TLE1 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF SYNOVIAL SARCOMA.
100		IHC-TTF-1	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC HELPS TO DISTINGUISH BETWEEN DIFFERENT TYPES AND SUBTYPES OF A DISEASE, PARTICULARLY CANCERS, BY IDENTIFYING SPECIFIC MOLECULES WITHIN CELLS, KNOWN AS MARKERS. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. TTF-1 IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF LUNG CANCER.

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
101		IHC-VIMENTIN	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	IHC IS A SPECIAL STAINING PROCESS PERFORMED ON TISSUE SAMPLES TO DIAGNOSE DISEASE MORE PRECISELY. IHC PLAYS A CRUCIAL ROLE IN TREATMENT PLANNING. VIMENTIN IS USED FOR SELECTING APPROPRIATE TREATMENT AND DETAILED ANALYSIS OF RENAL CELL CARCINOMA AND COLORECTAL CANCER.
102	869	KNEE ASPIRATE CYTOLOGY	PAP STAINING/ MICROSCOPY	SMEAR/ ASPIRATE	STERILE CONTAINER	24-48 hours	
103	1678	LIQUID BASED CYTOLOGY (LBC)	PAP, LBC STAINING/ MICROSCOPY	SMEAR	LBC VIAL	24-48 hours	
104	1773	MICRO SATELLITE INSTABILITY PANEL (MSI)	IMMUNO-HISTOCHEMISTRY	TISSUE/ PARAFFIN EMBEDDED TISSUE BLOCK	10% FORMALIN CONTAINING CONTAINER	7-10 Days	MSI TESTING EVALUATES THE LIKELIHOOD OF POSSESSING A PARTICULAR INHERITED FORM OF COLORECTAL CANCER KNOWN AS HEREDITARY NON-POLYPOSIS COLORECTAL CANCER (HNPCC) OR LYNCH SYNDROME. MARKERS: MLH1, MSH2, MSH6 & PMS2.
105	876	NIPPLE DISCHARGE CYTOLOGY	PAP, H&E STAINING/ MICROSCOPY	NIPPLE DISCHARGE/ SMEAR	STERILE CONTAINER/ SMEAR	24-48 hours	

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
106	335	PAP SMEAR FROM ASWINI	PAP STAINING/ MICROSCOPY	SMEAR	CYTO FIX JAR	24-48 Hours	
107	336	PAP SMEAR FROM OUTSIDE	PAP,H&E STAINING/ MICROSCOPY	SMEAR	CYTO FIX JAR/ DRIED SMEAR	24-48 Hours	
108	870	PUS FOR CYTOLOGY	PAP,H&E STAINING/ MICROSCOPY	SMEAR	STERILE CONTAINER	24-48 Hours	
109	871	SECOND OPINION- CYTOLOGY	MICROSCOPY	SLIDE	SLIDE	24-48 Hours	
110	872	SECOND OPINION-HISTO PATHOLOGY	H&E STAINING/ MICROSCOPY	SLIDE/BLOCK/ SPECIMEN	SLIDE/BLOCK	2-4 days	
111	873	SEMEN CYTOLOGY	PAP STAINING/ MICROSCOPY	SEMEN	STERILE CONTAINER	24-48 Hours	
112	874	SKIN BIOPSY (TAKEN FROM ASWINI)	H&E STAINING/ MICROSCOPY	PUNCH BIOPSY SKIN	10% FORMALIN CONTAINING CONTAINER	4-5 days	
113	864	SPUTUM CYTOLOGY	PAP,H&E STAINING/ MICROSCOPY	SPUTUM	STERILE CONTAINER	24-48 Hours	

DEPT NAME: HISTOPATHOLOGY AND CYTOPATHOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
114	1510	TESTICULAR BIOPSY	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	4-5 days	
115	1532	THYROID WITH LYMPH NODE	H&E STAINING/ MICROSCOPY	TISSUE	10% FORMALIN CONTAINING CONTAINER	5-7 days	
116	322	TZANCH SMEAR	LEISHMAN STAINING/ MICROSCOPY	DISCHARGE SMEAR	DRY SMEAR	24-48 Hours	
117	778	URETHRAL WASHINGS FOR CYTOLOGY	PAP.H&E STAINING/ MICROSCOPY	URETHRAL WASH	STERILE CONTAINER	24-48 Hours	
118	321	URINE CYTOLOGY	PAP STAINING/ MICROSCOPY	URINE	STERILE CONTAINER	24-48 Hours	
119	309	VAULT SMEAR RECEIVED FROM OUTSIDE	PAP,H&E STAINING/ MICROSCOPY	SMEAR	CYTO FIX JAR/ DRIED SMEAR	24-48 Hours	
120	315	VAULT SMEAR TAKEN FROM ASWINI	PAP STAINING/ MICROSCOPY	SMEAR	CYTO FIX JAR	24-48 Hours	



DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
1	532	AFB CULTURE CSF	LJ SLANT	CSF	STERILE CONTAINER	8 Weeks	ACID-FAST BACILLUS (AFB) IS A TYPE OF BACTERIA THAT CAUSES <u>TUBERCULOSIS</u> (MYCOBACTERIUM TUBERCULOSIS), <u>OTHER TYPES OF MYCOBACTERIAL INFECTIONS</u> , SUCH AS LEPROSY (HANSEN'S DISEASE) AND ATYPICAL MYCOBACTERIAL INFECTIONS. A CEREBROSPINAL FLUID (CSF) ACID-FAST BACILLUS (AFB) CULTURE IS USED TO DETECT THE PRESENCE OF ACID-FAST BACILLI, PARTICULARLY MYCOBACTERIUM TUBERCULOSIS (THE BACTERIA RESPONSIBLE FOR TUBERCULOSIS), IN THE CSF, AIDING IN THE DIAGNOSIS AND TREATMENT OF TUBERCULOUS MENINGITIS
2	539	AFB CULTURE URINE	LJ SLANT	URINE	STERILE CONTAINER	8 Weeks	ACID-FAST BACILLUS (AFB) IS A TYPE OF BACTERIA THAT CAUSES <u>TUBERCULOSIS</u> (MYCOBACTERIUM TUBERCULOSIS), <u>OTHER TYPES OF MYCOBACTERIAL INFECTIONS</u> , SUCH AS LEPROSY (HANSEN'S DISEASE) AND OTHER ATYPICAL MYCOBACTERIAL INFECTIONS. AFB CULTURE CAN POSITIVELY CONFIRM A DIAGNOSIS OF TB OR OTHER MYCOBACTERIAL INFECTION, EXCEPT IN CASE OF MYCOBACTERIUM LEPRAE .
3	523	AFB CULTURE	LJ SLANT-MANUAL	ALL SAMPLES EXCEPT BLOOD	STERILE CONTAINER	8 Weeks	ACID-FAST BACILLUS (AFB) IS A TYPE OF BACTERIA THAT CAUSES <u>TUBERCULOSIS</u> (MYCOBACTERIUM TUBERCULOSIS) AND <u>OTHER TYPES OF MYCOBACTERIAL INFECTIONS</u> , SUCH AS LEPROSY (HANSEN'S DISEASE) AND OTHER ATYPICAL MYCOBACTERIAL INFECTIONS. AFB CULTURE CAN POSITIVELY CONFIRM A DIAGNOSIS OF TB OR OTHER MYCOBACTERIAL INFECTION, EXCEPT IN CASE OF MYCOBACTERIUM LEPRAE .
4	1405	AFB STAIN	ZN STAIN/MICROSCOPY	ALL SAMPLES	SLIDE	4 Hours	THE ACID-FAST STAIN, ALSO KNOWN AS THE ZIEHL-NEELSEN STAIN, IS A MICROSCOPIC TECHNIQUE USED TO DETECT THE PRESENCE OF ACID-FAST BACTERIA, PARTICULARLY THOSE IN THE GENUS MYCOBACTERIUM LIKE M. TUBERCULOSIS (THE CAUSE OF TUBERCULOSIS), M. LEPRAE (THE CAUSE OF LEPROSY) AND OTHER ATYPICAL MYCOBACTERIAL SPECIES.
5	248	AUTOMATED ASCITIC FLUID CULTURE & SUSCEPTIBILITY WITH MIC	AUTOMATED	ASCITIC FLUID	STERILE CONTAINER	3 Days	THE PERITONEAL CAVITY IS THE SPACE WITHIN THE ABDOMEN THAT CONTAINS THE INTESTINES, THE STOMACH, AND THE LIVER. IT IS LINED BY THE PERITONEUM, A THIN LAYER OF TISSUE. ASCITES IS THE ACCUMULATION OF A FLUID IN THE ABDOMINAL CAVITY, IT'S ACCUMULATION MAY BE DUE TO VARIOUS CONDITIONS, INCLUDING INFECTION. CULTURE AND SENSITIVITY OF PERITONEAL OR ASCITIC FLUID IS AN ESSENTIAL LABORATORY TEST THAT HELPS IN THE DIAGNOSIS AND TREATMENT OF INFECTIONS IN THE ABDOMINAL CAVITY.
6	1101	AUTOMATED BLOOD ANAEROBIC CULTURE	AUTOMATED ANAEROBIC CULTURE	BLOOD	BACT ALERT BLOOD CULTURE BOTTLE-ORANGE CAP	5 Days	AUTOMATED BLOOD ANAEROBIC CULTURE AND SUSCEPTIBILITY TESTING IDENTIFIES THE PRESENCE OF ANAEROBIC BACTERIA THE BLOOD, AIDING IN THE DIAGNOSIS AND TREATMENT OF BLOODSTREAM INFECTIONS (BACTEREMIA/SEPSIS) BY DETERMINING THE CAUSATIVE AGENT.
7	543	AUTOMATED C.S.F. CULTURE & SUSCEPTIBILITY WITH MIC	AUTOMATED	CSF	STERILE CONTAINER	3 Days	A CSF CULTURE AND SENSITIVITY (CSF C/S) TEST DETECTS INFECTIONS IN THE FLUID AROUND THE BRAIN AND SPINAL CORD, IDENTIFYING BACTERIAL OR FUNGAL ORGANISMS AND THEIR ANTIBIOTIC SENSITIVITY, INCLUDING MIC VALUES. IT IS THE GOLD STANDARD FOR DIAGNOSING CNS INFECTIONS AND GUIDES EFFECTIVE TREATMENT.

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
8	847	AUTOMATED CULTURE & SUSCEPTIBILITY	AUTOMATED	CSF	STERILE CONTAINER	3 Days	AUTOMATED CULTURE AND SENSITIVITY (C&S) TESTS, USE AUTOMATED SYSTEMS TO DETECT AND IDENTIFY BACTERIA/FUNGI (YEAST), AND THEN DETERMINE THEIR SUSCEPTIBILITY TO VARIOUS ANTIBIOTICS, PROVIDING ACCURATE IDENTIFICATION WITH SUSCEPTIBILITY PATTERN WITH MIC VALUES , WHICH ARE MORE EFFICIENT THAN TRADITIONAL METHODS.
9	837	AUTOMATED FUNGAL CULTURE (YEAST)	AUTOMATED	ALL SAMPLES	STERILE CONTAINER	3 Days	AUTOMATED FUNGAL CULTURE AND SENSITIVITY (C&S) TESTS, USE AUTOMATED SYSTEMS TO DETECT AND IDENTIFY FUNGI (YEAST), AND THEN DETERMINE THEIR SUSCEPTIBILITY TO VARIOUS ANTIBIOTICS, PROVIDING ACCURATE IDENTIFICATION WITH SUSCEPTIBILITY PATTERN WITH MIC VALUES .
10	850	AUTOMATED SYNOVIAL FLUID CULTURE & SUSCEPTIBILITY WITH MIC	AUTOMATED	SYNOVIAL FLUID	STERILE CONTAINER	3 Days	SYNOVIAL FLUID CULTURE AND SENSITIVITY TESTING IS USED TO IDENTIFY BACTERIA/FUNGI (YEAST) CAUSING JOINT INFECTIONS (SEPTIC ARTHRITIS) AND DETERMINE WHICH ANTIBIOTICS ARE EFFECTIVE, AIDING IN TIMELY AND APPROPRIATE TREATMENT
11	514	C.S.F. CULTURE & SENSITIVITY	MANUAL	CSF	STERILE CONTAINER	3 Days	A CEREBROSPINAL FLUID CULTURE AND SENSITIVITY TEST, ALSO KNOWN AS A CSF C/S, IS A LABORATORY TEST THAT EXAMINES THE FLUID SURROUNDING THE BRAIN AND SPINAL CORD FOR SIGNS OF INFECTION(BACTERIAL/FUNGAL ETIOLOGY) IDENTIFYING THE CAUSATIVE ORGANISM AND ITS ANTIBIOTIC SENSITIVITY BY CONVENTIONAL METHODS.CEREBROSPINAL FLUID (CSF) CULTURE IS THE GOLD STANDARD FOR CENTRAL NERVOUS SYSTEM (CNS) INFECTIONS AND CAN PROVIDE CRITICAL INFORMATION FOR PATIENT MANAGEMENT.
13	1099	FLUID ANAEROBIC CULTURE	ANAEROBIC CULTURE AUTOMATED	ALL STERILE BODY FLUIDS	BACT ALERT BLOOD CULTURE BOTTLE/STERILE CONTAINER	5 Days	AUTOMATED FLUID ANAEROBIC CULTURE AND SUSCEPTIBILITY TESTING IDENTIFIES THE PRESENCE OF ANAEROBIC BACTERIA THE BODY FLUIDS, AIDING IN THE DIAGNOSIS AND TREATMENT OF THESE INFECTIONS, BY DETERMINING THE CAUSATIVE AGENT.
14	540	FUNGAL CULTURE	MANUAL	A SAMPLES EXCEPT SWAB	STERILE CONTAINER	30 Days	THE MOST DIRECT AND CONCLUSIVE WAY OF IDENTIFYING THE FUNGUS CAUSING AN INFECTION IS TO GROW IT FROM A PATIENT SAMPLE, SUCH AS BLOOD, CSF, PUS, URINE, TISSUE, RESPIRATORY SAMPLES (SPUTUM, BAL), PLEURAL, PERICARDIAL OR PERITONEAL FLUID, SKIN SCRAPING, HAIR, NAIL CLIPPINGS, ORAL OR VAGINAL SAMPLES, AND IDENTIFYING THE ISOLATE BY COLONY CHARACTERS AND MICROSCOPIC MORPHOLOGY.
15	541	FUNGAL STAIN	MICROSCOPY	ALL SAMPLES EXCEPT SWAH	AIR TIGHT CONTAINER	1 Day	A FUNGAL STAIN TEST IS A LABORATORY TEST USED TO DETECT THE PRESENCE OF FUNGI IN A BIOLOGICAL SPECIMEN, SUCH AS TISSUE, CORNEAL SCRAPING, ASPIRATES OR BODY FLUIDS

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
16	953	GENEXPERT MTB RIF ULTRA	AUTOMATED REAL TIME PCR	SPUTUM TISSUE URINE (3 DAYS EARLY MORNING SAMPLES)	AIR TIGHT CONTAINER	4 Hours	A FUNGAL STAIN TEST IS A LABORATORY TEST USED TO DETECT THE PRESENCE OF FUNGI IN A BIOLOGICAL SPECIMEN, SUCH AS TISSUE, CORNEAL SCRAPPING, ASPIRATES OR BODY FLUIDS
17	509	GRAMS STAIN	MICROSCOPY	ALL SAMPLES EXCEPT BLOOD AND STOOL	STERILE CONTAINER	3 Hours	GRAM STAIN IS OFTEN THE INITIAL DIAGNOSTIC TEST FOR THE EVALUATION OF INFECTIONS, THE GRAM STAIN DIFFERENTIATES BACTERIA INTO TWO FUNDAMENTAL VARIETIES OF CELLS, "GRAM-POSITIVE" & "GRAM-NEGATIVE." POSITIVE GRAM STAIN REPORT INDICATES TYPE OF MICROORGANISM & THEIR MORPHOLOGY BASED ON THE TYPE OF CELLS SEEN, MICROBIAL LOAD IN THE GIVEN SPECIMEN CAN BE ASSESSED , IT CAN BE USED TO DETECT THE PRESENCE OF FUGAL ELEMENTS AND IS ALSO USED TO SCREEN SPUTUM SPECIMENS TO ESTABLISH ACCEPTABILITY FOR BACTERIAL CULTURE.
18	510	KOH SMEAR FOR FUNGUS	MICROSCOPY	SKIN/ HAIR /NAIL	AIR TIGHT CONTAINER	3 Hours	KOH (POTASSIUM HYDROXIDE) SMEAR TEST, IS A QUICK AND SIMPLE DIAGNOSTIC TEST USED TO DEMONSTRATE THE EVIDENCE OF FUNGAL INFECTION IN SKIN, HAIR, AND NAILS BY DISSOLVING CELLULAR DEBRIS AND ALLOWING VISUALIZATION OF FUNGAL ELEMENTS UNDER THE MICROSCOPE.
19	841	NASAL SWAB CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	NASAL SWAB	STERILE SWAB	3 Days	THE CULTURE AND SENSITIVITY, NASAL SWAB TEST IS A DIAGNOSTIC PROCEDURE PERFORMED TO IDENTIFY AND ANALYZE MICROORGANISMS PRESENT IN THE NASAL PASSAGES.
20	533	NASAL SWAB CULTURE & SENSITIVITY	MANUAL	NASAL SWAB	STERILE SWAB	3 Days	THE CULTURE AND SENSITIVITY, NASAL SWAB TEST IS A DIAGNOSTIC PROCEDURE PERFORMED TO IDENTIFY AND ANALYZE MICROORGANISMS PRESENT IN THE NASAL PASSAGES. THE TEST CAN ALSO BE USED TO SCREEN MRSA , MRSA SCREENING HELPS IDENTIFY INDIVIDUALS WHO ARE CARRYING MRSA BACTERIA, EVEN WITHOUT SHOWING SYMPTOMS, TO PREVENT THE SPREAD OF THE INFECTION.
21	829	PERITONEAL DIALYSIS FLUID & AUTOMATED SUSCEPTIBILITY	AUTOMATED	FLUID	STERILE CONTAINER	3 Days	PERITONEAL DIALYSIS FLUID CULTURE,IS A CRITICAL TEST FOR DIAGNOSING AND MANAGING PERITONITIS (INFECTION OF THE PERITONEAL CAVITY), INVOLVES COLLECTING A SAMPLE OF THE DIALYSIS FLUID, CULTURING IT TO IDENTIFY ANY BACTERIA OR FUNGI, THEN DETERMINE THEIR SUSCEPTIBILITY TO VARIOUS ANTIBIOTICS, PROVIDING ACCURATE IDENTIFICATION WITH SUSCEPTIBILITY PATTERN WITH MIC VALUES , WHICH ARE MORE EFFICIENT THAN TRADITIONAL METHODS.
22	825	PERITONEAL DIALYSIS FLUID C/S	MANUAL	FLUID	STERILE CONTAINER	3 Days	PERITONEAL DIALYSIS FLUID CULTURE,IS A CRITICAL TEST FOR DIAGNOSING AND MANAGING PERITONITIS (INFECTION OF THE PERITONEAL CAVITY), INVOLVES COLLECTING A SAMPLE OF THE DIALYSIS FLUID, CULTURING IT TO IDENTIFY ANY BACTERIA OR FUNGI, AND DETERMINE ITS ANTIBIOTIC SENSITIVITY BY CONVENTIONAL METHODS.

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
23	825	PERITONEAL DIALYSIS FLUID C/S	MANUAL	FLUID	STERILE CONTAINER	3 Days	AUTOMATED PLEURAL FLUID ANAEROBIC CULTURE AND SUSCEPTIBILITY TESTING IDENTIFIES THE PRESENCE OF ANAEROBIC BACTERIA AIDING IN THE DIAGNOSIS AND TREATMENT BY DETERMINING THE CAUSATIVE AGENT.
24	1104	PUS ANAEROBIC CULTURE	ANAEROBIC CULTURE SENSITIVITY	PUS	STERILE CONTAINER IN ANAEROBIC CONDITION	5 Days	AN ANAEROBIC PUS CULTURE TEST IDENTIFIES BACTERIA IN PUS THAT THRIVE IN LOW-OXYGEN ENVIRONMENTS, HELPING DIAGNOSE INFECTIONS IN DEEP WOUNDS AND ABSCESES, PARTICULARLY THOSE CAUSED BY ANAEROBIC BACTERIA.
25	830	PUS CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	PUS	STERILE CONTAINER	3 Days	THE <u>PUS CULTURE AND SENSITIVITY TEST</u> , ALSO KNOWN AS THE PUS C&S TEST, IS A DIAGNOSTIC TOOL USED TO IDENTIFY AND DETERMINE THE CAUSE OF AN INFECTION. ONE PRIMARY REASON FOR CONDUCTING A PUS C&S TEST IS TO PINPOINT THE SPECIFIC BACTERIA OR MICROORGANISM RESPONSIBLE FOR AN INFECTION. THIS KNOWLEDGE HELPS DOCTORS SELECT THE MOST EFFECTIVE ANTIBIOTIC TREATMENT, ENSURING TARGETED THERAPY THAT CAN COMBAT THE INFECTION MORE EFFICIENTLY.
26	524	SEMEN CULTURE & SENSITIVITY	MANUAL	SEMEN	STERILE CONTAINER	3 Days	SEMEN CULTURE AND SENSITIVITY TESTING IS DONE TO RULE OUT THE PRESENCE OF ANY INFECTIONS IN THE SEMEN. THIS IS IMPORTANT AS VARIOUS BACTERIAL INFECTIONS CAN LEAD TO ABNORMAL MORPHOLOGY OF THE SPERMS AND CAN ALSO AFFECT THE MOTILITY OF THE SPERMS. SEMEN CULTURE IS ESSENTIAL WHEN THE MAN SUDDENLY PRESENTS WITH SYMPTOMS OF THE LOWER URINARY TRACT (LUTS) AND GENITAL SYSTEM -SUCH AS PAINFUL EJACULATION, HAEMOSPERMIA (PRESENCE OF BLOOD IN THE SEMEN)- AND ALSO IN CASES OF URINARY FREQUENCY OR URGENCY, DYSURIA ETC
27	838	SPUTUM CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	SPUTUM	STERILE CONTAINER	3 Days	SPUTUM CULTURE HELPS DETERMINE THE SPECIFIC TYPE OF BACTERIA OR FUNGUS CAUSING A RESPIRATORY INFECTION, WHICH IS CRUCIAL FOR EFFECTIVE TREATMENT. SPUTUM CULTURE CAN BE USED TO MONITOR THE EFFECTIVENESS OF TREATMENT FOR RESPIRATORY INFECTIONS. METHOD INVOLVES COLLECTING A SAMPLE OF THE SPUTUM, CULTURING IT TO IDENTIFY ANY BACTERIA OR FUNGI, THEN DETERMINE THEIR SUSCEPTIBILITY TO VARIOUS ANTIBIOTICS, PROVIDING ACCURATE IDENTIFICATION WITH SUSCEPTIBILITY PATTERN WITH MIC VALUES , WHICH ARE MORE EFFICIENT THAN TRADITIONAL METHODS.
28	527	SPUTUM FOR AFB STAIN	ZN STAIN MICROSCOPY	SPUTUM	STERILE CONTAINER	4 Hours	A SPUTUM AFB (ACID-FAST BACILLUS) STAIN IS A TEST THAT EXAMINES A SPUTUM SAMPLE UNDER A MICROSCOPE TO DETECT THE PRESENCE OF ACID-FAST BACTERIA, PARTICULARLY MYCOBACTERIUM TUBERCULOSIS, THE BACTERIA THAT CAUSES TUBERCULOSIS (TB). SPUTUM AFB SMEAR IS GRADED AS PER NTEP GUIDELINES

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
29	835	STOOL CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	STOOL CIS	STERILE CONTAINER	3 Days	A STOOL CULTURE, ALSO KNOWN AS A FECAL CULTURE, IS USED TO DETECT PATHOGENIC BACTERIA IN A STOOL SAMPLE TO DIAGNOSE GASTROINTESTINAL INFECTIONS AND DETERMINE THE APPROPRIATE TREATMENT. STOOL CULTURES CAN HELP INVESTIGATE ETIOLOGICAL AGENTS OF DIARRHEA, ABDOMINAL DISCOMFORT, OR OTHER GASTROINTESTINAL ISSUES. BY IDENTIFYING THE SPECIFIC PATHOGEN, DOCTORS CAN CHOOSE THE MOST EFFECTIVE TREATMENT, SUCH AS ANTIBIOTICS FOR BACTERIAL INFECTIONS.
30	576	STOOL HANGING DROP	HANGING DROP-MIS	STOOL	STERILE CONTAINER	2 Hours	THE HANGING DROP TEST FOR STOOL IS PERFORMED TO VISUALIZE THE MOTILITY OR MOVEMENT PATTERN OF BACTERIA IN THE STOOL SAMPLE AND IDENTIFY THEM FROM THESE PATTERNS TO HELP IN THE DIAGNOSIS. THE TEST PRINCIPALLY HELPS IN PRESUMPTIVE DIAGNOSIS OF CHOLERA CAUSED BY THE INFECTION OF VIBRIO CHOLERAE.
31	506	SWAB CULTURE & SENSITIVITY	MANUAL	SWAB	STERILE SWAB	3 Days	INCLUDES BACTERIAL CULTURE OF SUPERFICIAL WOUND SWABS, SKIN SWABS, ULCER SWABS, ABSCESS SWABS, PUS SWABS, EAR SWABS, EYE SWABS, AND THROAT SWABS. SAMPLES OF EXUDATE AND PUS IS PREFERABLE TO SWABS IF PRESENT. SAMPLE A REPRESENTATIVE PART OF THE LESION OR SITE. SWABBING DRY CRUSTED AREAS IS UNLIKELY TO YIELD THE CAUSATIVE PATHOGEN. IF SPECIMENS ARE TAKEN FROM ULCERS, THE DEBRIS ON THE ULCER SHOULD BE REMOVED AND THE ULCER SHOULD BE CLEANED WITH SALINE PRIOR TO SAMPLING. USED TO DETERMINE IF THE SITE IS INFECTED, BY IDENTIFYING THE SPECIFIC PATHOGEN, DOCTORS CAN CHOOSE THE MOST EFFECTIVE TREATMENT, SUCH AS ANTIBIOTICS FOR BACTERIAL INFECTIONS.
32	845	THROAT SWAB CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	THROAT SWAB	STERILE SWAB	3 Days	THE SWAB THROAT CULTURE TEST IS USED TO DIAGNOSE BACTERIAL OR FUNGAL INFECTIONS IN THE THROAT, SUCH AS ACUTE SORE THROAT (PHARYNGITIS) OR TONSILLITIS. IT HELPS PINPOINT THE EXACT CAUSE OF INFECTION FOR EFFECTIVE TREATMENT.
33	528	THROAT SWAB FOR GRAM'S STAIN	GRAM STAIN/MICROSCOPY	THROAT SWAB	STERILE SWAB	4 Hours	A THROAT SWAB GRAM STAIN IS A MICROSCOPIC EXAMINATION OF A SAMPLE TAKEN FROM THE THROAT, USED TO IDENTIFY AND CLASSIFY BACTERIA, TO DETECT THE PRESENCE OF FUNGI ESPECIALLY YEAST, WHICH HELPS GUIDE TREATMENT DECISIONS.
34	552	THROAT SWAB KLB	ALBERT'S STAIN/M/S	STERILE SWAB SMEAR FROM THROAT	STERILE CONTAINER	24 Hours	A THROAT SWAB FOR KLB (ALBERT) STAIN IS A TEST USED TO PRESUMPTIVELY DETECT CORYNEBACTERIUM DIPHTHERIAE, THE BACTERIA THAT CAUSES DIPHTHERIA, BY EXAMINING A THROAT SWAB FOR THE PRESENCE OF CHARACTERISTIC GRANULES CALLED METACHROMATIC GRANULES. DIPHTHERIA IS A SERIOUS BACTERIAL INFECTION THAT CAN CAUSE A SORE THROAT, FEVER, AND A THICK, GRAY MEMBRANE IN THE THROAT, WHICH CAN OBSTRUCT BREATHING.

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
35	831	TIP CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	CATHETER TIP	STERILE CONTAINER	3 Days	A "TIP CULTURE AND SENSITIVITY" TEST, OFTEN PERFORMED ON CATHETER TIPS, HELPS DETERMINE IF A CATHETER-RELATED INFECTION IS SUSPECTED AND IDENTIFIES THE BACTERIA CAUSING THE INFECTION, AS WELL AS WHICH ANTIBIOTICS ARE EFFECTIVE AGAINST THEM.
36	849	URETHRAL SWAB CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	URETHRAL SWAB	STERILE SWAB	3 Days	THIS TEST IS OFTEN CARRIED OUT TO DETERMINE THE CAUSATIVE FACTORS BEHIND THE URETHRAL INFECTION AND TO DETERMINE THE RIGHT COURSE OF TREATMENT.
37	581	URINE AFB STAIN	ZN STAIN/MICROSCOPY	URINE-3 CONSECUTIVE DAY SAMPLES, PREFERABLY EARLY MORNING SAMPLES	URINE CONTAINER	5 Days	THE AFB-SMEAR EXAMINATION ZN STAIN ON URINE IS A MICROSCOPIC TEST USED TO DETECT THE PRESENCE OF ACID-FAST BACILLI (AFB) BACTERIA IN THE URINARY TRACT, FOR THIS THREE CONSECUTIVE DAYS, THREE COMPLETE URINE SPECIMENS SHOULD BE TAKEN IN THE EARLY MORNING (FIRST PASSED). EACH COLLECTION IS HOUSED IN A STERILE CONTAINER, AND SEND TO THE LABORATORY.
38	826	URINE CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	URINE	STERILE CONTAINER	3 Days	A URINE CULTURE TEST IS USED TO DETECT AND IDENTIFY BACTERIA OR OTHER MICROORGANISMS (YEAST) IN A URINE SAMPLE, PRIMARILY TO DIAGNOSE URINARY TRACT INFECTIONS AND DETERMINE THE MOST EFFECTIVE ANTIBIOTIC TREATMENT. OTHER IMPORTANT USES INCLUDE. CONFIRM ANTIBIOTIC EFFECTIVENESS: A REPEAT URINE CULTURE AFTER <u>ANTIBIOTIC TREATMENT</u> SHOWS IF THE PRESCRIBED MEDICATIONS SUCCESSFULLY CURED THE UTI. INVESTIGATE RECURRENT UTIS: IDENTIFYING BACTERIAL STRAINS CAUSING REPEATED UTIS GUIDES PREVENTION APPROACHES TO STOP REINFECTION.
39	1063	AUTOMATED BLOOD FUNGAL CULTURE *	AUTOMATED	BLOOD	BLOOD CULTURE BOTTLE	14 Days	A FUNGAL CULTURE TEST IS USED TO FIND OUT WHETHER YOU HAVE A FUNGAL INFECTION. THIS TEST HELPS TO IDENTIFY THE TYPE OF FUNGUS PRESENT IN THE SAMPLE. IT IS ALSO USED TO GUIDE TREATMENT AND TO SEE IF TREATMENT IS WORKING.
40	827	AUTOMATED BLOOD CULTURE & SUSCEPTIBILITY	AUTOMATED	BLOOD	BACT ALERT BLOOD CULTURE BOTTLE (YELLOW-PEDIATRIC AND GREEN ADULT)	5 Days	A LABORATORY PROCESS THAT USES AUTOMATED SYSTEMS TO DETECT AND IDENTIFY MICROORGANISMS IN BLOOD SAMPLES, AND THEN DETERMINE WHICH ANTIBIOTICS ARE EFFECTIVE AGAINST THEM.

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
41	843	AUTOMATED BONE MARROW CULTURE & SUSCEPTIBILITY WITH MIC	AUTOMATED	BLOOD	BACT ALERT CULTURE BOTTLE STERILE CONTAINER	5 Days	AUTOMATED BLOOD CULTURE AND SUSCEPTIBILITY TESTING SYSTEMS PROVIDE AUTOMATED INOCULATION, READING AND INTERPRETATION. THESE SYSTEMS HAVE THE ADVANTAGE OF BEING RAPID AND CONVENIENT. THE RESULTS USUALLY HELP TO IDENTIFY THE SPECIFIC BACTERIA OR FUNGI THAT'S CAUSING THE INFECTION. DEPENDING ON THE TYPE OF ORGANISM DISCOVERED, ANOTHER TEST CALLED SENSITIVITY OR SUSCEPTIBILITY TEST IS PERFORMED. THIS HELPS TO DETERMINE WHICH MEDICATION WILL WORK BEST AGAINST THAT ORGANISM.
42	836	AUTOMATED C.S.F. CULTURE & SUSCEPTIBILITY WITH MIC	AUTOMATED	CSF	BACT ALERT CULTURE BOTTLE STERILE CONTAINER	3 Days	CEREBROSPINAL FLUID (CSF) IS A CLEAR, COLORLESS, WATERY FLUID THAT FLOWS IN AND AROUND YOUR BRAIN AND SPINAL CORD. CSF CULTURE IS A LABORATORY TEST USED TO LOOK FOR BACTERIA, FUNGI, AND VIRUSES IN THE CSF. AUTOMATED CULTURE AND SUSCEPTIBILITY TESTING SYSTEMS PROVIDE AUTOMATED INOCULATION, READING AND INTERPRETATION. THE RESULTS USUALLY HELP TO IDENTIFY THE SPECIFIC BACTERIA OR FUNGI THAT'S CAUSING THE INFECTION. DEPENDING ON THE TYPE OF ORGANISM DISCOVERED, ANOTHER TEST CALLED SENSITIVITY OR SUSCEPTIBILITY TEST IS PERFORMED. THIS HELPS TO DETERMINE WHICH MEDICATION WILL WORK BEST AGAINST THAT ORGANISM.
43	951	AUTOMATED CULTURE & SUSCEPTIBILITY	AUTOMATED	ALL SAMPLES	BACT ALERT BLOOD CULTURE BOTTLE-ORANGE CAP	3 Days	AUTOMATED CULTURE AND SUSCEPTIBILITY TESTING SYSTEMS PROVIDE AUTOMATED INOCULATION, READING AND INTERPRETATION. THESE SYSTEMS HAVE THE ADVANTAGE OF BEING RAPID AND CONVENIENT. THE RESULTS USUALLY HELP TO IDENTIFY THE SPECIFIC BACTERIA OR FUNGI THAT'S CAUSING THE INFECTION.
44	833	AUTOMATED FLUID CULTURE & SUSCEPTIBILITY WITH MIC	AUTOMATED	FLUID	BACT ALERT BLOOD CULTURE BOTTLE/STERILE CONTAINER	3-5 Days	AUTOMATED CULTURE AND SUSCEPTIBILITY TESTING SYSTEMS PROVIDE AUTOMATED INOCULATION, READING AND INTERPRETATION. THESE SYSTEMS HAVE THE ADVANTAGE OF BEING RAPID AND CONVENIENT. THE RESULTS USUALLY HELP TO IDENTIFY THE SPECIFIC BACTERIA OR FUNGI THAT'S CAUSING THE INFECTION.
45	837	AUTOMATED FUNGAL CULTURE (YEAST)	AUTOMATED	ALL SAMPLES	STERILE CONTAINER	3 Days	AUTOMATED FUNGAL CULTURE AND SENSITIVITY (C&S) TESTS, USE AUTOMATED SYSTEMS TO DETECT AND IDENTIFY FUNGI (YEAST), AND THEN DETERMINE THEIR SUSCEPTIBILITY TO VARIOUS ANTIBIOTICS, PROVIDING ACCURATE IDENTIFICATION WITH SUSCEPTIBILITY PATTERN WITH MIC VALUES .
46	583	CULTURE & SENSITIVITY	MANUAL	ALL SAMPLES	STERILE CONTAINER	3 Days	CHECK WHETHER THE SAMPLE HAS ANY BACTERIAL INFECTION AND ALSO ABOUT THE DRUG SUSCEPTIBILITY OF THE ISOLATED ORGANISM.

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
47	840	EAR SWAB CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	STERILE SWAB	STERILE CONTAINER	3 Days	THE CULTURE AND SENSITIVITY EAR SWAB IS A TEST THAT IS PERFORMED ON A SAMPLE OF SWAB TAKEN FROM THE EAR. IT IS PERFORMED TO CONFIRM EAR INFECTION. AUTOMATED CULTURE AND SUSCEPTIBILITY TESTING SYSTEMS PROVIDE AUTOMATED INOCULATION, READING AND INTERPRETATION. DEPENDING ON THE TYPE OF ORGANISM DISCOVERED, ANOTHER TEST CALLED SENSITIVITY OR SUSCEPTIBILITY TEST IS PERFORMED. THIS HELPS TO DETERMINE WHICH MEDICATION WILL WORK BEST AGAINST THAT ORGANISM.
48	872	EAR SWAB CULTURE & SENSITIVITY	MANUAL	STERILE SWAB	STERILE CONTAINER	3 Days	THE CULTURE AND SENSITIVITY EAR SWAB IS A TEST THAT IS PERFORMED ON A SAMPLE OF SWAB TAKEN FROM THE EAR. IT IS PERFORMED TO CONFIRM EAR INFECTION. THE CULTURE IS THEN TESTED TO CHECK AND CONFIRM THE CAUSATIVE AGENTS BEHIND YOUR CURRENT INFECTION AS WELL AS TO DETERMINE THE SENSITIVITY OF THE BACTERIA TO ANTIBIOTICS. THIS SENSITIVITY TEST IS THEN USED TO DETERMINE THE BEST COURSE OF TREATMENT AGAINST THE INFECTION.
49	569	FLUID CULTURE & SENSITIVITY	MANUAL	FLUID	STERILE CONTAINER	3 Hours	CULTURE AND SENSITIVITY TESTING IS DONE TO HELP DIAGNOSE AN INFECTION. IT ALSO HELP YOUR HEALTH CARE PROVIDER TO DECIDE WHICH MEDICINES TO USE FOR TREATING YOUR INFECTION.
50	505	INDIA INK PREPARATION FOR FUNGUS (CRYPTO-COCCUS)	WET FILM-M/S	CSF	STERILE CONTAINER	4 Hours	INDIA INK IS USED TO DETECT THE PRESENCE OF ENCAPSULATED YEAST IN CSF, PARTICULARLY CRYPTOCOCCUS NEOFORMANS WHICH CAUSES CRYPTOCOCCAL MENINGITIS.
51	567	NAIL CLIPPING FOR FUNGUS	WET FILM-M/S	NAIL PIECE	AIR TIGHT CONTAINER	24 Hours	DIAGNOSIS OF FUNGAL INFECTION IN NAIL.
52	577	NIPPLE DISCHARGE C/S	MANUAL	NIPPLE DISCHARGE	STERILE CONTAINER	3 Days	NIPPLE DISCHARGE IS A COMMON COMPLAINT IN WOMEN WHO ARE NOT PREGNANT OR BREASTFEEDING, ESPECIALLY DURING THE REPRODUCTIVE YEARS. SPONTANEOUS UNILATERAL NIPPLE DISCHARGE, REGARDLESS OF COLOR IS CONSIDERED ABNORMAL. NIPPLE DISCHARGE MAY BE BREAST MILK OR AN EXUDATE PRODUCED BY A NUMBER OF CONDITIONS. IT CAN BE SEROUS (YELLOW), MUCINOUS (CLEAR AND WATERY), MILKY, BLOODY, PURULENT, MULTICOLORED AND STICKY, OR PINK. CULTURE AND SENSITIVITY OF NIPPLE DISCHARGE IS DONE TO DETECT THE PRESENCE OF ANY PATHOGENIC ORGANISMS IF PRESENT.

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
53	828	NIPPLE DISCHARGE CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	NIPPLE DISCHARGE	STERILE CONTAINER	3 Days	NIPPLE DISCHARGE IS COMMON IN NON-PREGNANT, NON-BREASTFEEDING WOMEN, BUT SPONTANEOUS UNILATERAL DISCHARGE IS ABNORMAL. IT CAN VARY IN COLOR AND CONSISTENCY, INCLUDING MILKY, BLOODY, OR PURULENT. CULTURE AND SENSITIVITY TESTING DETECT PATHOGENIC ORGANISMS, WITH AUTOMATED SYSTEMS AIDING ANALYSIS. IF BACTERIA OR FUNGI ARE FOUND, A SUSCEPTIBILITY TEST DETERMINES THE BEST TREATMENT.
54	816	OP (OPERATION THEATRE SWAB)	MANUAL	STERILE SWAB	AIR TIGHT CONTAINER	14 Days	THE OT SWAB CULTURE TEST IS DONE TO CHECK FOR AND IDENTIFY AEROBIC BACTERIA ON DIFFERENT EQUIPMENT AND SURFACES IN AN OPERATION THEATRE. ITS PURPOSE IS TO DETERMINE IF THE OT IS FREE FROM HARMFUL MICROBES.
55	1151	PARASITE IDENTIFICATION	AUTOMATED	STOOL	STERILE CONTAINER	5 Days	THIS TEST LOOKS FOR PARASITES AND THEIR EGGS (OVA) IN A SAMPLE OF YOUR STOOL. PARASITES CAN LIVE IN YOUR DIGESTIVE SYSTEM AND CAUSE ILLNESS.
56	1103	PLEURAL FLUID ANAEROBIC CULTURE	AUTOMATED ANAEROBIC CULTURE	PLEURAL FLUID	BACT ALERT BLOOD CULTURE BOTTLE ORANGE CAP	5 Days	AUTOMATED PLEURAL FLUID ANAEROBIC CULTURE AND SUSCEPTIBILITY TESTING IDENTIFIES THE PRESENCE OF ANAEROBIC BACTERIA AIDING IN THE DIAGNOSIS AND TREATMENT BY DETERMINING THE CAUSATIVE AGENT.
57	516	PLEURAL FLUID CULTURE & SENSITIVITY	MANUAL	FLUID	AIR TIGHT CONTAINER	5 Days	PLEURAL FLUID CULTURE IS A TEST THAT EXAMINES A SAMPLE OF FLUID THAT HAS COLLECTED IN THE PLEURAL SPACE TO SEE IF YOU HAVE AN INFECTION OR UNDERSTAND THE CAUSE OF BUILDUP OF FLUID IN THIS SPACE. THE PLEURAL SPACE IS THE AREA BETWEEN THE LINING OF THE OUTSIDE OF THE LUNGS (PLEURAL AND THE WALL OF THE CHEST).
58	517	PUS CULTURE & SENSITIVITY	MANUAL	PUS	STERILE CONTAINER	3 Days	THIS TEST IS DONE ON A PUS SAMPLE TO IDENTIFY THE PRESENCE OF PATHOGENIC ORGANISMS. A POSITIVE RESULT MEANS THAT BACTERIA OR OTHER ORGANISMS DID GROW AND THAT YOUR WOUND IS INFECTED. FROM YOUR TEST RESULTS, YOUR HEALTHCARE PROVIDER CAN FIGURE OUT THE BEST ANTIBIOTIC TO TREAT IT.
59	670	PUS FOR AFB CULTURE	LJ SLANT	PUS	STERILE CONTAINER	56 Days	ISOLATION OF VIABLE MYCOBACTERIUM TUBERCULOSIS FROM THE SAMPLE.
60	544	SCRAPING FOR FUNGUS	WET FILM M/S	SKIN SCRAPINGS	AIR TIGHT CONTAINER	3 Hours	DIRECT EXAMINATION OF SKIN SCRAPINGS FOR FUNGUS.

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
61	582	SCRAPINGS FOR PARASITES	MICROSCOPY	SKIN SCRAPINGS	AIR TIGHT CONTAINER	2 Hours	DIRECT EXAMINATION OF SKIN SCRAPPINGS FOR PARASITES.
62	834	SEMEN CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	SEMEN	STERILE CONTAINER	3 Days	USED TO IDENTIFY AEROBIC BACTERIAL INFECTION PRESENT IN THE SEMEN SAMPLE AND SELECTION OF APPROPRIATE ANTIBIOTICS AGAINST THE PATHOGEN CAUSING THE INFECTION.
63	570	SKIN SLIT SMEAR LEPROSY	ZN STAIN/ MICROSCOPY	SMEAR	SMEAR IN SLIDE	24 Hours	SLIT SKIN SMEAR (SSS) IS A TECHNIQUE FOR LEPROSY DIAGNOSIS. LEPROSY IS A CHRONIC INFECTIOUS DISEASE CAUSED BY MYCOBACTERIUM LEPRAE.
64	564	SPUTUM AFB CULTURE	LJ SLANT MANUAL	3 SPUTUM SAMPLES (3 CONSECUTIV E DAYS)	STERILE AIR TIGHT CONTAINER	8 Weeks	DETECTS THE PRESENCE OF VIABLE MYCOBACTERIUM TUBERCULOSIS FROM THE SPUTUM SAMPLE.
65	518	SPUTUM CULTURE & SENSITIVITY	MANUAL	SPUTUM	STERILE CONTAINER	3 Days	A SPUTUM CULTURE AND SENSITIVITY IS A TEST THAT CHECKS FOR BACTERIA OR ANOTHER TYPE OF ORGANISM THAT MAY BE CAUSING AN INFECTION IN YOUR LUNGS OR THE AIRWAYS LEADING TO THE LUNGS AND ALSO TO CHECK WHAT KIND OF MEDICINE WILL INHIBIT THE GROWTH OF THE ORGANISM AND HELP IN TREATING THE INFECTION.
66	554	SPUTUM FOR GRAM STAIN	GRAM STAIN/ MICROSCOPY	SPUTUM	STERILE CONTAINER	3 Hours	A SPUTUM GRAM STAIN IS A LABORATORY TEST USED TO DETECT BACTERIA IN A SPUTUM SAMPLE. THE GRAM STAIN METHOD IS ONE OF THE MOST COMMONLY USED METHODS TO RAPIDLY IDENTIFY THE CAUSE OF A BACTERIAL INFECTION, INCLUDING PNEUMONIA. IT ALSO DETERMINES WHAT TYPE OF BACTERIA (GRAM POSITIVE OR GRAM NEGATIVE) ARE CAUSING IT.
67	525	STOOL CULTURE & SENSITIVITY	MANUAL	STOOL	STERILE CONTAINER	3 Days	ISOLATION OF PATHOGENIC MICROBES FROM THE GIVEN STOOL SAMPLE AND SELECTION OF APPROPRIATE ANTIBIOTICS AGAINST THE PATHOGEN CAUSING THE INFECTION.
68	832	SWAB CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	SWAB	STERILE SWAB	3 Days	A SWAB CULTURE IS A QUICK, PAINLESS TEST USED TO DETECT INFECTIONS BY GROWING MICROORGANISMS LIKE BACTERIA, FUNGI, OR VIRUSES IN A LAB. AUTOMATED SYSTEMS SPEED UP THIS PROCESS, OFFERING FAST AND CONVENIENT RESULTS. ONCE THE ORGANISM IS IDENTIFIED, A SUSCEPTIBILITY TEST MAY BE DONE TO FIND THE MOST EFFECTIVE TREATMENT.

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
69	519	SYNOVIAL FLUID CULTURE & SENSITIVITY	MANUAL	SYNOVIAL FLUID	STERILE CONTAINER	3 Days	SYNOVIAL FLUID, ALSO KNOWN AS JOINT FLUID, IS A THICK LIQUID LOCATED. BETWEEN YOUR JOINTS. THE FLUID CUSHIONS THE ENDS OF BONES AND REDUCES FRICTION WHEN YOU MOVE YOUR JOINTS. SYNOVIAL FLUID CULTURE IS A LABORATORY TEST TO DETECT INFECTION-CAUSING PATHOGENS IN A SAMPLE OF FLUID SURROUNDING A JOINT.
70	520	THROAT SWAB CULTURE & SENSITIVITY	MANUAL	THROAT SWAB	STERILE SWAB	3 Days	A THROAT SWAB CULTURE, OR THROAT CULTURE, IS A TEST COMMONLY USED TO DIAGNOSE BACTERIAL INFECTIONS IN THE THROAT BY USING CONVENTIONAL CULTURE AND SENSITIVITY METHOD. THESE INFECTIONS CAN INCLUDE STREP THROAT, PNEUMONIA, TONSILLITIS, WHOOPING COUGH, AND MENINGITIS. A NEGATIVE THROAT SWAB CULTURE MEANS THAT NO INFECTIOUS BACTERIA ARE PRESENT IN YOUR THROAT. A POSITIVE TEST INDICATES THE PRESENCE OF STREPTOCOCCI (THE BACTERIA THAT CAUSES STREP THROAT) OR OTHER BACTERIA AND DEPENDING ON THE TYPE OF ORGANISM DETECTED, ANOTHER TEST CALLED SENSITIVITY OR SUSCEPTIBILITY TEST IS DONE IN ORDER TO DETERMINE WHICH MEDICATION WILL WORK BEST AGAINST THAT ORGANISM.
71	551	TIP CULTURE & SENSITIVITY	MANUAL	CATHETER TIP	STERILE CONTAINER	3 Days	THE CULTURE AND SENSITIVITY CATHETER TEST INVOLVES A CATHETER TUBE WITH A POINTED TIP WHICH IS INSERTED INTO THE INFECTED AREA FOR COLLECTING THE SAMPLE OF THE WOUND. ONCE THE SPECIMEN FOR THE STUDY HAS BEEN COLLECTED, THE CATHETER IS REMOVED. A 5 CM PORTION OF IT IS CUT OFF AND SENT FOR TESTING. THIS TEST IS VERY USEFUL IN IDENTIFYING DISEASES AND CRITICAL HEALTH CONDITIONS CAUSED DUE TO BACTERIA, FUNGI, AND OTHER MICROORGANISMS. THE TEST IDENTIFIES THE PRESENCE OF SUCH ORGANISMS.
72	571	URETHRAL SMEAR FOR GONOCOCCI	GRAM STAIN/MICROSCOPY	URETHRAL DISCHARGE	SMEAR STERILE CONTAINER	4 Hours	GONORRHEA IS A SEXUALLY TRANSMITTED DISEASE CAUSED BY THE BACTERIUM-NEISSERIA GONORRHOEAE, WHICH INVOLVES INFLAMMATORY DISCHARGE FROM THE URETHRA OR VAGINA. IT CAN CAUSE SERIOUS COMPLICATIONS IF NOT DETECTED AND TREATED. A POSITIVE TEST INDICATES AN ACTIVE GONORRHEA INFECTION THAT REQUIRES TREATMENT WITH A COURSE OF ANTIBIOTICS.
73	849	URETHRAL SWAB CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	URETHRAL SWAB	STERILE SWAB	3 Days	THIS TEST IS OFTEN CARRIED OUT TO DETERMINE THE CAUSATIVE FACTORS BEHIND THE URETHRAL INFECTION AND TO DETERMINE THE RIGHT COURSE OF TREATMENT.
74	521	URETHRAL SWAB CULTURE & SENSITIVITY	MANUAL	URETHRAL SWAB	STERILE SWAB	3 Days	THIS TEST IS OFTEN CARRIED OUT TO DETERMINE THE CAUSATIVE FACTORS BEHIND THE URETHRAL INFECTION AND TO DETERMINE THE RIGHT COURSE OF TREATMENT. THIS TEST IS QUITE EFFECTIVE IN DETERMINING ALL FORMS OF URETHRA INFECTIONS INCLUDING-STDs, GONORRHEA, ETC.

DEPT NAME: MICROBIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
75	512	URINE CULTURE & SENSITIVITY	MANUAL	URINE	STERILE CONTAINER	3 Days	URINE CULTURE IS THE GOLD STANDARD FOR DETECTING ASYMPTOMATIC BACTERIURIA. WHEN A PATIENT IS SUSPECTED OF HAVING A URINARY TRACT INFECTION (UTI), A URINE SAMPLE FOR CULTURE AND SENSITIVITY IS ORDERED. COMMON ORGANISMS ISOLATED ARE E.COLI, KLEBSIELLA, S.SAPROPHYTICUS, S.AUREUS, ENTEROCOCCUS, PROTEUS AND PSUDOMONAS. WHEN AN ORGANISM IS ISOLATED, ANTIBIOTIC SENSITIVITIES ARE PERFORMED TO GUIDE THE ANTIBIOTIC SELECTION.
76	530	URINE DEPOSIT FOR GONOCOCCI	MICROSCOPY	URINE	URINE CONTAINER	4 Hours	GONORRHEA IS A SEXUALLY TRANSMITTED DISEASE CAUSED BY THE BACTERIUM-NEISSERIA GONORRHOEAE, WHICH INVOLVES INFLAMMATORY DISCHARGE FROM THE URETHRA OR VAGINA. IT CAN CAUSE SERIOUS COMPLICATIONS IF NOT DETECTED AND TREATED. A POSITIVE TEST INDICATES AN ACTIVE GONORRHEA INFECTION THAT REQUIRES TREATMENT WITH A COURSE OF ANTIBIOTICS.
77	563	VAGINAL SMEAR-CANDIDA/TV	WET FILM M/S	VAGINAL/CERVICAL DISCHARGE	SMEAR	4 Hours	TRICHOMONAS VAGINALIS (TV) IS A SEXUALLY TRANSMITTED INFECTION (STI) CAUSED BY A TINY ORGANISM THAT LIVES INSIDE THE VAGINA AND URETHRA. CANDIDIASIS IS AN INFECTION CAUSED BY A YEAST (A TYPE OF FUNGUS) CALLED CANDIDA. IT CAN CAUSE VAGINAL CANDIDIASIS. VAGINAL SMEAR TEST HELPS IN DIAGNOSING THESE INFECTIONS.
78	846	VAGINAL SWAB CULTURE & AUTOMATED SUSCEPTIBILITY	AUTOMATED	VAGINAL SWAB	STERILE SWAB	3 Days	THE CULTURE AND SENSITIVITY - VAGINAL SWAB IS OFTEN PERFORMED TO DETECT THE PATHOGEN THAT CAUSES VAGINAL IRRITATION AND DISCHARGE. THIS TEST HELPS TO IDENTIFY AND TREAT VAGINAL INFECTIONS. AUTOMATED ANTIMICROBIAL SUSEPTIBILITY TESTING SYSTEMS PROVIDE AUTOMATED INOCULATION, READING AND INTERPRETATION. THESE SYSTEMS HAVE THE ADVANTAGE OF BEING RAPID AND CONVENIENT. THE CULTURE IS THEN TESTED TO CONFIRM THE CAUSATIVE FACTORS BEHIND THE INFECTION, AS WELL AS TO DETERMINE THE SENSITIVITY OF THE BACTERIA TO ANTIBIOTICS. THIS SENSITIVITY TEST IS USED TO DETERMINE THE BEST COURSE OF TREATMENT AGAINST THE INFECTION.
79	522	VAGINAL SWAB CULTURE & SENSITIVITY	MANUAL	VAGINAL SWAB	STERILE SWAB	3 Days	THE CULTURE AND SENSITIVITY VAGINAL SWAB IS OFTEN PERFORMED TO DETECT THE PATHOGEN THAT CAUSES VAGINAL IRRITATION AND DISCHARGE. THIS TEST HELPS TO IDENTIFY AND TREAT VAGINAL INFECTIONS. THE CULTURE IS THEN USED TO CHECK AND CONFIRM THE CAUSATIVE FACTORS BEHIND THE INFECTION AS WELL AS TO DETERMINE THE SENSITIVITY OF THE BACTERIA TO ANTIBIOTICS. THIS SENSITIVITY TEST IS USED TO DETERMINE THE BEST COURSE OF TREATMENT AGAINST THE INFECTION.

MOLECULAR BIOLOGY

DEPT NAME: MOLECULAR BIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
1	001221	TRUENAT H1N1 PCR	TRUENAT RT PCR	NASOPHARYNGEAL /THROAT SWAB	IN VTM	4 Hours	H1N1 INFLUENZA, A SUBTYPE OF INFLUENZA A VIRUS, IS AN INFECTIOUS VIRAL ILLNESS THAT CAUSES BOTH UPPER AND, IN SOME CASES, LOWER RESPIRATORY TRACT INFECTIONS IN ITS HOST. H1N1 INFLUENZA INFECTIONS CAN CAUSE SYMPTOMS SUCH AS RHINORRHEA, RIGORS, MYALGIA, HEADACHE, FEVER, CHILLS, LOSS OF APPETITE, AND, POSSIBLY, LOWER RESPIRATORY TRACT DISEASE AND GASTROINTESTINAL DISEASE. RARELY, H1N1 INFLUENZA CAN LEAD TO ACUTE RESPIRATORY DISTRESS SYNDROME AND SECONDARY BACTERIAL SEPSIS.
2	001770	TRUENAT INFLUENZA A&B RTPCR	TRUENAT RT PCR	NASOPHARYNGEAL /THROAT SWAB	IN VTM	4 Hours	INFLUENZA IS A COMMUNICABLE VIRAL DISEASE THAT AFFECTS THE UPPER AND LOWER RESPIRATORY TRACT. A WIDE SPECTRUM OF INFLUENZA VIRUSES (TYPES A,B,C,D) CAUSES IT. INFLUENZA A AND B ARE THE TWO MAIN TYPES THAT ROUTINELY SPREAD IN HUMANS AND CAUSE SEASONAL FLU EPIDEMICS.SOME OF THESE VIRUSES CAN INFECT HUMANS, AND SOME ARE SPECIFIC TO DIFFERENT SPECIES. THESE VIRUSES ARE TRANSMISSIBLE THROUGH RESPIRATORY DROPLETS EXPELLED FROM THE MOUTH AND RESPIRATORY SYSTEM DURING COUGHING, TALKING, AND SNEEZING. THE INFLUENZA VIRUSES CAN BE TRANSMITTED BY TOUCHING INANIMATE OBJECTS SOILED WITH THE VIRUS AND TOUCHING THE NOSE OR EYE.
3	001546	TRUENAT HIV-1/HIV-2 VIRAL LOAD ASSAY	TRUENAT RT PCR	WHOLE BLOOD/ EDTA	EDTA CONTAINER	4 Hours	HIV-1 AND HIV-2 ARE TWO TYPES OF HUMAN IMMUNODEFICIENCY VIRUS, WITH HIV-1 BEING THE MORE COMMON AND AGGRESSIVE FORM GLOBALLY. HIV-2 IS PRIMARILY FOUND IN WEST AFRICA AND PROGRESSES MORE SLOWLY, WITH A LOWER RISK OF TRANSMISSION COMPARED TO HIV-1. BOTH VIRUSES CAN LEAD TO ACQUIRED IMMUNODEFICIENCY SYNDROME (AIDS)
4	001692	TRUENAT CYTOMEGALOVIRUS (CMV) DNA, QUANTITATIVE PCR	TRUENAT RT PCR	WHOLEBLOOD/ EDTA ,TISSUE	EDTA CONTAINER, TISSUE	4 Hours 24 Hours(Tissue)	HUMAN CYTOMEGALOVIRUS (CMV), ALSO KNOWN AS HERPESVIRUS-5, IS A UBIQUITOUS VIRUS THAT BELONGS TO THE HERPESVIRIDAE, A LARGE FAMILY OF DNA VIRUSES. PRIMARY INFECTION WITH CMV CAN MANIFEST AS EITHER ASYMPTOMATIC INFECTION, A FLU-LIKE SYNDROME, MONONUCLEOSIS, OR OTHER SYNDROMES. WHEN PATIENTS ARE IMMUNOCOMPROMISED, EG, AFTER SOLID ORGAN AND HEMATOPOIETIC TRANSPLANTATION OR ON CHEMOTHERAPY CMV CAN REACTIVATE, CAUSING VIREMIA WHICH CAN CAUSE SEVERE ORGAN DAMAGE, INCLUDING CMV ESOPHAGITIS, COLITIS ETC.
5	001601	TRUENAT LEPTOSPIRA-REAL TIME RTPCR	TRUENAT RT PCR	WHOLE BLOOD/ EDTA	EDTA CONTAINER	4 Hours	LEPTOSPIROSIS IS A DISEASE CAUSED BY THE BACTERIUM LEPTOSPIRA THAT CAN BE FOUND IN CONTAMINATED WATER OR SOIL.WITHOUT TREATMENT, LEPTOSPIROSIS IN PEOPLE CAN LEAD TO KIDNEY DAMAGE, MENINGITIS, LIVER FAILURE, TROUBLE BREATHING, AND EVEN DEATH.HUMAN LEPTOSPIRAL INFECTIONS RESULT PRIMARILY FROM DIRECT OR INDIRECT EXPOSURE TO THE URINE OF INFECTED ANIMALS.
6	001338	*HPV-HR REAL TIME DUPLEX PCR	REAL TIME-PCR	CERVICAL SWAB	VTM	4 Hours	THE HUMAN PAPILLOMAVIRUS (HPV) IS A NON-ENVELOPED, DOUBLE-STRANDED, CIRCULAR DNA VIRUS THAT IS RESPONSIBLE FOR CAUSING MULTIPLE EPITHELIAL LESIONS AND CANCERS. IT CAN MANIFEST AS CUTANEOUS AND ANOGENITAL WARTS, WHICH DEPENDING ON THE SUBTYPE, MAY PROGRESS TO CARCINOMA.

DEPT NAME: MOLECULAR BIOLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
7	001219	TRUENAT HLA-B27 PCR	TRUENAT RT PCR	WHOLE BLOOD/ EDTA	EDTA CONTAINER	4 Hours	THE HLA B27 STANDS FOR HUMAN LEUKOCYTE ANTIGEN B27 TEST ISUSED TO DETERMINE THE PRESENCE OF THE HLA B27 ANTIGEN ON THE SURFACE OF WHITE BLOOD CELLS. THE HLA-B27 ANTIGEN IS A CLASS I MAJOR HISTOCOMPATIBILITY COMPLEX MOLECULE WHOSE PRESENCE IS ASSOCIATED WITH DISEASES SUCH AS UVEITIS, PSORIASIS, REITER'S SYNDROME, INFLAMMATORY BOWEL DISEASES (CROHN'S DISEASE, ULCERATIVE COLITIS), SERONEGATIVE ARTHROPATHIES (PSORIATIC ARTHROPATHIES, ANKYLOSING SPONDYLITIS) AND WHIPPLE'S DISEASE.
8	001636	HBV DNA QUANTITATIVE REAL-TIME PCR	REAL TIME-PCR	WHOLE BLOOD/ EDTA	EDTA CONTAINER	48 Hours	HEPATITIS B IS CAUSED BY INFECTION WITH HBV. HBV IS A 40-42-NANOMETER ENVELOPED VIRUS CLASSIFIED IN THEHEPADNAVIRIDAEFAMILY.HBV IS HIGHLY INFECTIOUS AND CAN BE TRANSMITTED BY PERCUTANEOUS, MUCOSAL, OR NONINTACT SKIN EXPOSURE TO INFECTIOUS BLOOD, SEMEN, AND/OR OTHER BODY FLUIDS. HBV CAN CAUSE BOTH ACUTE/CHRONIC HEPATITIS, SYMPTOMS CAN RANGE FROM ASYMPTOMATIC OR MILD DISEASE TO, RARELY,CIRRHOSIS, FULMINANT HEPATITIS TO EXTRACELLULAR CARCINOMA
9	001314	SARS CoV-2 RTPCR	RT-PCR	NASOPHARYNGEAL /THROAT SWAB	IN VTM	48 Hours	SEVERE ACUTE RESPIRATORY SYNDROME CORONAVIRUS 2 (SARS-COV-2) IS A HIGHLY TRANSMISSIBLE AND PATHOGENIC CORONAVIRUS THAT EMERGED IN LATE 2019 AND HAS CAUSED A PANDEMIC OF ACUTE RESPIRATORY DISEASE, NAMED 'CORONAVIRUS DISEASE 2019' (COVID-19), TRANSMISSION OF SARS-COV-2 CAN OCCUR THROUGH DIRECT, INDIRECT, OR CLOSE CONTACT WITH INFECTED PEOPLE THROUGH INFECTED SECRETIONS SUCH AS SALIVA AND RESPIRATORY SECRETIONS OR THEIR RESPIRATORY DROPLETS. INFECTION WITH SARS-COV-2 PRIMARILY CAUSES RESPIRATORY ILLNESS RANGING FROM MILD DISEASE TO SEVERE DISEASE AND DEATH, AND SOME PEOPLE INFECTED WITH THE VIRUS NEVER DEVELOP SYMPTOMS.
10	001044	GENE X PERT HCV VIRAL LOAD ASSAY (Quantitative PCR)	REAL TIME-PCR	WHOLE BLOOD/EDTA PLASMA (3ML)	EDTA CONTAINER	4 Hours	HEPATITIS C IS A LIVER DISEASE CAUSED BY HCV, THE HEPATITIS C VIRUS BELONGS TO THE GENUS <u>HEPACIVIRUS</u> , A MEMBER OF THE FAMILY <u>FLAVIVIRIDAE</u> . HEPATITIS C CAN RANGE FROM A MILD ILLNESS LASTING A FEW WEEKS TO A SERIOUS, LONG-TERM ILLNESS.HCV IS HIGHLY INFECTIOUS AND CAN BE TRANSMITTED BY PERCUTANEOUS, MUCOSAL, OR NONINTACT SKIN EXPOSURE TO INFECTIOUS BLOOD, SEMEN, AND/OR OTHER BODY FLUIDS. HCV CAN CAUSE BOTH ACUTE/CHRONIC HEPATITIS , SYMPTOMS CAN RANGE FROM ASYMPTOMATIC OR MILD DISEASE TO, RARELY,CIRRHOSIS,FULMINANT HEPATITIS TO EXTRACELLULAR CARCINOMA.
11	001790	JAK2 V617F MUTATION DETECTION	REAL TIME-PCR	WHOLE BLOOD	EDTA CONTAINER	3 Days	THIS TEST IS DESIGNED FOR THE QUALITATIVE DETECTION OF THE V617F MUTATION IN THE JANUS KINASE 2 GENE (JAK2) BY REAL TIME PCR ON THE GENOMIC DNA OF PATIENTS SUSPECTED OF MYELOPROLIFERATIVE DISORDERS LIKE POLYCYTHEMIA VERA, PRIMARY MYELOFIBROSIS AND ESSENTIAL THROMBOCYTHEMIA.
11	001790	JAK2 EXON 12 MUTATION DETECTION	REAL TIME-PCR	WHOLE BLOOD	EDTA CONTAINER	3 Days	IS A BLOOD TEST THAT IDENTIFIES SPECIFIC MUTATIONS WITHIN JAK-2 GENE, SPECIFICALLY IN EXON-12, WHICH IS ASSOCIATED WITH CERTAIN BLOOD DISORDERS, PARTICULARLY POLYCYTHEMIA VERA (PV). THIS ISSUED WHEN PATIENT TEST NEGATIVE FOR MORE COMMON JAK2 V617F MUTATION BUT STILL PRESENTS WITH SYMPTOMS OF PV.



SEROLOGY



DEPT NAME: SEROLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
1	585	A.S.O TITRE	IMMUNO-TURBIDIMETRY	SERUM	SERUM CONTAINER	2 Hours	ANTI ANTISTREPTOLYSIN O TITER (ASO) IS A BLOOD TEST USED TO DETERMINE IF YOU'VE HAD A RECENT INFECTION CAUSED BY GROUP A STREPTOCOCCUS BACTERIA.
2	166	ANA SCREENING	CLIA	SERUM	SERUM CONTAINER	5 Hours	ANTI ANA TEST IS A BLOOD TEST THAT LOOKS FOR ANTINUCLEAR ANTIBODIES IN YOUR BLOOD. ANA TEST HELP IN THE DIAGNOSIS OF AUTOIMMUNE DISORDERS, SUCH AS: SYSTEMIC LUPUS ERYTHEMATOSUS (SLE RHEUMATOID ARTHRITIS, SCLERODERMA, SJÖGREN'S SYNDROME, ETC.
3	574	ANTI-HCV	CMIA	SERUM	SERUM CONTAINER	2 Hours	A BLOOD TEST, CALLED AN HCV ANTIBODY TEST, IS USED TO FIND OUT IF SOMEONE HAS EVER BEEN INFECTED WITH THE HEPATITIS C VIRUS. THE HCV ANTIBODY TEST, SOMETIMES CALLED THE ANTI-HCV TEST, LOOKS FOR ANTIBODIES TO THE HEPATITIS C VIRUS IN BLOOD.
4	168	ANTI ds DNA.	CLIA	SERUM	SERUM CONTAINER	5 Hours	THE ANTI-DOUBLE STRANDED DNA (ANTI-DSDNA) TESTS ARE USED TO HELP DIAGNOSE AND MONITOR LUPUS, ALSO CALLED SYSTEMIC LUPUS ERYTHEMATOSUS OR SLE, A CHRONIC INFLAMMATORY AUTOIMMUNE DISORDER IN WHICH THE IMMUNE SYSTEM MISTAKENLY TARGETS THE BODY'S OWN CELLS AND TISSUES.
5	887	ANTI HBS	CMIA	SERUM	SERUM CONTAINER	2 Hours	THE PRESENCE OF ANTI-HBS IS GENERALLY INTERPRETED AS RECOVERY AND IMMUNITY FROM HEPATITIS B VIRUS INFECTION. ANTI-HBS ALSO DEVELOPS IN A PERSON WHO HAS BEEN SUCCESSFULLY VACCINATED AGAINST HEPATITIS B. AMONG VACCINE RESPONDERS WHO COMPLETED A VACCINE SERIES, ANTI-HBS LEVELS CAN DECLINE OVER TIME, HOWEVER THE MAJORITY ARE STILL IMMUNE AND WILL MOUNT A RESPONSE WHEN EXPOSED TO HBV.
6	1178	CMV IgG ANTIBODY	CLIA	SERUM	SERUM CONTAINER	5 Hours	CMV INFECTION IS A DISEASE CAUSED BY A TYPE OF HERPES VIRUS. THE CMV BLOOD TEST IS PERFORMED TO DETECT ACTIVE CMV INFECTION, OR PAST CMV INFECTION IN PEOPLE WHO ARE AT RISK FOR REACTIVATION OF INFECTION. THOSE PEOPLE INCLUDES, ORGAN TRANSPLANT RECIPIENT AND THOSE WITH A SUPPRESSED IMMUNE SYSTEM.
7	588	C.M.V IgM ANTIBODY	CLIA	SERUM	SERUM CONTAINER	5 Hours	CMV INFECTION IS A DISEASE CAUSED BY A TYPE OF HERPES VIRUS. THE CMV BLOOD TEST IS PERFORMED TO DETECT ACTIVE CMV INFECTION, OR PAST CMV INFECTION IN PEOPLE WHO ARE AT RISK FOR REACTIVATION OF INFECTION. THESE PEOPLE INDUDES, ORGAN TRANSPLANT RECIPIENTS AND THOSE WITH A SUPPRESSED IMMUNE SYSTEM .

DEPT NAME: SEROLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
8	169	C-ANCA (PR3-PROTEINASE)	ELISA	SERUM	SERUM CONTAINER	30 Hours	THIS TEST LOOKS FOR ANTINEUTROPHIL CYTOPLASMIC ANTIBODIES (ANCA IN A SAMPLE OF YOUR BLOOD. IT IS USED PRIMARILY IN THE DIAGNOSIS AND MONITORING OF PATIENTS WITH WEGENER'S GRANULOMATOSIS (WG), IT IT ALSO AN IMPORTANT MARKER FOR THE DIFFERENTIAL DIAGNOSIS OF AUTE IMMUNE VASCULITIS.
9	155	CARDIOLIPIN ANTIBODY IgM	CLIA	SERUM	SERUM CONTAINER	5 Hours	ANTICARDIOLIPIN IGM (ACLA IGM) ANTIBODIES ARE AUTOANTIBODIES THAT MISTAKENLY ATTACK CARDIOLIPIN, A TYPE OF LIPID THAT IS A CRUCIAL COMPONENT OF CELL MEMBRANES, PARTICULARLY THE INNER MITOCHONDRIAL MEMBRANES AND PLATELETS. THE PRESENCE OF THESE ANTIBODIES IS ASSOCIATED WITH AN INCREASED RISK OF BLOOD CLOTS E.G DEEP VEIN THROMBOSIS (DVT), PULMONARY EMBOLISM (PE) AND OTHER COMPLICATIONS. ELEVATED LEVELS OF ANTICARDIOLIPIN IGM ANTIBODIES ARE LINKED TO AN INCREASED RISK OF RECURRENT MISCARRIAGES, STILLBIRTHS, PREECLAMPSIA, AND OTHER PREGNANCY-RELATED PROBLEMS.
10	156	CARDIOLIPIN ANTIBODY IgG	CLIA	SERUM	SERUM CONTAINER	5 Hours	ANTI-CARDIOLIPIN IGG (ACLA IGG) ARE ANTIBODIES THAT TARGET CARDIOLIPIN, A PHOSPHOLIPID. THESE ANTIBODIES ARE ASSOCIATED WITH ANTIPHOSPHOLIPID SYNDROME (APS), AN AUTOIMMUNE DISORDER LINKED TO AN INCREASED RISK OF BLOOD CLOTS AND RECURRENT MISCARRIAGES. THE TEST MEASURES IGG ANTIBODIES AGAINST CARDIOLIPIN IN THE BLOOD TO HELP DIAGNOSE CONDITIONS LIKE APS OR INVESTIGATE UNEXPLAINED BLOOD CLOTS AND RECURRENT MISCARRIAGES.
11	172	CHIKUNGUNYA CARD IgM	IMMUNO CHROMATOGRAPHY	SERUM	SERUM CONTAINER	2 Hours	THIS TEST IS A RAPID LATERAL FLOW CHROMATOGRAPHIC ASSAY & MAY BE USED AS AN AID TO DIAGNOSE RECENT INFECTION WITH CHIKUNGUNYA VIRUS.
12	589	C-REACTIVE PROTEIN	IMMUNO TURBIDIMETRY	SERUM	SERUM CONTAINER	2 Hours	CRP IS A BLOOD TEST THAT MEASURES THE LEVEL OF C-REACTIVE PROTEIN IN BLOOD. CRP IN PRIMARILY PRODUCED BY THE LIVER IN RESPONSE TO INFLAMMATION IN THE BODY IT IS ALSO USED TO MONITOR CHRONIC INFLAMMATORY DISEASE, EVALUATE TREATMENT EFFECTIVENESS & CHECK POST SURGICAL COMPLICATION
13	182	DENGUE IgM	ELISA	SERUM	SERUM CONTAINER	12 Hours	A POSITIVE IGM TEST MEANS YOU HAVE LIKELY BEEN INFECTED WITH DENGUE RECENTLY. A NEGATIVE IGM TEST COULD MEAN YOU ARE NOT INFECTED OR YOU TESTED TOO SOON. IGM ANTIBODY LEVELS ALSO INDICATE WHETHER THIS IS A PRIMARY (FIRST) OR SECONDARY (PAST) INFECTION.
14	786	EBV ANTIBODY TO VIRAL CAPSID ANTIGEN-IGM	ELFA	SERUM	SERUM CONTAINER	5 Hours	EPSTEIN-BARR VIRUS (EBV) SPREADS THROUGH SALIVA OR SHARING UTENSILS AND IS THE LEADING CAUSE OF INFECTIOUS MONONUCLEOSIS ("MONO"), COMMON IN TEENS AND ADULTS. SYMPTOMS INCLUDE FATIGUE, FEVER, SORE THROAT, SWOLLEN GLANDS, TONSILS, RASH, AND SOMETIMES AN ENLARGED SPLEEN. THE INFECTION USUALLY RESOLVES ON ITS OWN. THIS TEST DETECTS EBV-SPECIFIC ANTIBODIES; ANTI-VCA IGM APPEARS EARLY IN INFECTION AND TYPICALLY DISAPPEARS WITHIN 4-6 WEEKS, INDICATING RECENT INFECTION.

DEPT NAME: SEROLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
15	558	H.S.V 1 IgG Antibody	ELISA	SERUM	SERUM CONTAINER	30 Hours	AN IGG BLOOD TEST IS USED TO DIAGNOSE HERPES SIMPLEX VIRUS INFECTION. IT DETECTS THE ANTIBODIES YOUR BODY PRODUCES TO FIGHT AGAINST THE VIRUS. A TEST RESULT SHOWING IGG ANTIBODIES WILL TELL A PERSON THAT THEY HAVE EXPERIENCED AN INFECTION IN THE PAST AND THAT THEY HAVE ANTIBODIES AGAINST THE VIRUS.
16	559	H.S.V 1 IgM Antibody	ELISA	SERUM	SERUM CONTAINER	30 Hours	A HERPES IGM TEST IS A BLOOD TEST THAT CAN DETECT EARLY HERPES SIMPLEX VIRUS (HSV 1) INFECTION.
17	581	H.S.V 2 IgG Antibody	ELISA	SERUM	SERUM CONTAINER	30 Hours	AN IGG BLOOD TEST IS USED TO DIAGNOSE HERPES SIMPLEX VIRUS INFECTION. IT DETECTS THE ANTIBODIES YOUR BODY PRODUCES TO FIGHT AGAINST THE VIRUS. A TEST RESULT SHOWING IGG ANTIBODIES WILL TELL A PERSON THAT THEY HAVE EXPERIENCED AN INFECTION IN THE PAST AND THAT THEY HAVE ANTIBODIES AGAINST THE VIRUS.
18	562	H.S.V 2 IgM Antibody	ELISA	SERUM	SERUM CONTAINER	30 Hours	A HERPES IGM TEST IS A BLOOD TEST THAT CAN DETECT EARLY HERPES SIMPLEX VIRUS (HSV2) INFECTION.
19	572	HAVAb-IgM	CMIA	SERUM	SERUM CONTAINER	2 Hours	THIS TEST LOOKS FOR ANTIBODIES IN YOUR BLOOD AGAINST HEPATITIS A VIRUS, IN ORDER TO FIND OUT IF YOU ARE INFECTED WITH THE HEPATITIS A VIRUS (HAV).
20	1033	HBE ANTIBODY	CLIA	SERUM	SERUM CONTAINER	5 Hours	IN HEPATITIS B VIRUS (HBV) CARRIERS AND IN PATIENTS WITH CHRONIC HEPATITIS B, POSITIVE ANTI-HBE RESULTS USUALLY INDICATE INACTIVITY OF THE VIRUS AND LOW INFECTIVITY OF THE PATIENTS.
21	591	HBE ANTIGEN	CLIA	SERUM	SERUM CONTAINER	5 Hours	DIAGNOSIS AND MONITORING OF HEPATITIS B VIRUS INFECTIVITY. HEPATITIS BE ANTIGEN (HBEAG) INDICATES ONGOING VIRUS REPLICATION AND INCREASED INFECTIVITY.
22	613	HBsAg	CMIA	SERUM	SERUM CONTAINER	2 Hours	HBSAG (HEPATITIS B SURFACE ANTIGEN) A "POSITIVE" OR "REACTIVE" HBSAG TEST RESULT MEANS THAT THE PERSON IS INFECTED WITH HEPATITIS B VIRUS.
23	549	HBsAg CARD	IMMUNO-CHROMATOGRAPHY	SERUM	SERUM CONTAINER	2 Hours	AN HBSAG CARD TEST IS A RAPID DIAGNOSTIC TEST TO DETECT THE PRESENCE OF HBSAG IN A BLOOD SAMPLE, IT IS A COMMON SCREENING TOOL USED IN VARIOUS ROUTINE CHECKUPS, PRENATAL SCREENINGS & BEFORE BLOOD DONATION.

DEPT NAME: SEROLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
24	1666	HCV-CARD	IMMUNO-CHROMATOGRAPHY	SERUM	SERUM CONTAINER	2 Hours	A BLOOD TEST, CALLED AN HCV ANTIBODY TEST, IS USED TO FIND OUT IF SOMEONE HAS EVER BEEN INFECTED WITH THE HEPATITIS C VIRUS. THE HCV ANTIBODY TEST, SOMETIMES CALLED THE ANTI-HCV TEST, LOOKS FOR ANTIBODIES TO THE HEPATITIS C VIRUS IN BLOOD.
25	1693	HELICOBACTER PYLORI RAPID STOOL TEST	IMMUNO-CHROMATOGRAPHY	STOOL	AIR TIGHT CONTAINER	4 Hours	HELICOBACTER PYLORI (H PYLORI) BACTERIA IS A COMMON CAUSE OF PEPTIC ULCERS (SORES IN THE LINING OF THE STOMACH, SMALL INTESTINE, OR ESOPHAGUS). IN THIS TEST, A STOOL SAMPLE IS TESTED TO CHECK FOR H PYLORI ANTIGENS.
26	1620	HEPATITIS A ANTIBODY (ANTI HAV) IGM CARD	IMMUNO-CHROMATOGRAPHY	SERUM	SERUM CONTAINER	2 Hours	THE PRESENCE OF ANTI-HAV (IGM) IN HUMAN SERUM OR PLASMA IS INDICATIVE OF PAST OR PRESENT INFECTION WITH HEPATITIS A VIRUS (HAV) OR VACCINATION AGAINST HAV. THE TEST FOR ANTI-HAV IS PRIMARILY USED TO DETERMINE EXPOSURE TO HAV EITHER NATURALLY OR DUE TO VACCINATION.
27	1175	HSV1 & 2 IGG ANTIBODY	CLIA	SERUM	SERUM CONTAINER	5 Hours	AN IGG BLOOD TEST IS USED TO DIAGNOSE HERPES SIMPLEX VIRUS INFECTION. IT DETECTS THE ANTIBODIES YOUR BODY PRODUCES TO FIGHT AGAINST THE VIRUS. A TEST RESULT SHOWING IGG ANTIBODIES WILL TELL A PERSON THAT THEY HAVE EXPERIENCED AN INFECTION IN THE PAST AND THAT THEY HAVE ANTIBODIES AGAINST THE VIRUS (HSV1 AND HSV2).
28	1176	HSV1 & 2 IGM ANTIBODY	CLIA	SERUM	SERUM CONTAINER	5 Hours	A HERPES IGM TEST IS A BLOOD TEST THAT CAN DETECT EARLY HERPES SIMPLEX VIRUS (HSV1 AND HSV2) INFECTION.
29	181	LEPTOSPIRA IGM	ELISA	SERUM	SERUM CONTAINER	8 Hours	LEPTOSPIROSIS IS A ZOO NOTIC DISEASE TRANSMITTED THROUGH SOIL, FOOD OR WATER CONTAMINATED BY URINE OF INFECTED ANIMALS. THIS TEST IS INTENDED TO BE USED AS AN AID FOR THE DIAGNOSIS OF ACUTE OR RECENT LEPTOSPIROSIS DUE TO INFECTION WITH LEPTOSPIRA SPECIES.
30	966	MEASLES IgG	CLIA	SERUM	SERUM CONTAINER	5 Hours	MEASLES IS A CONTAGIOUS DISEASE MAINLY INFECTING CHILDREN AND IMMUNOCOMPROMISED PATIENTS. IGG DETECTS PREVIOUS EXPOSURE OR IMMUNITY AGAINST MEASLES VIRUS.
31	174	MUMPS IgG	CLIA	SERUM	SERUM CONTAINER	30 Hours	IGG DETECTS PREVIOUS EXPOSURE OR IMMUNITY AGAINST MUMPS VIRUS.

DEPT NAME: SEROLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
32	165	P-ANCA (MPO-MYELOPEROXIDASE)	ELISA	SERUM	SERUM CONTAINER	30 Hours	ANTIBODIES AGAINST MPO ARE LINKED TO VASCULITIS, PARTICULARLY THREE FORMS: GRANULOMATOSIS WITH POLYANGIITIS (GPA), MICROSCOPIC POLYANGIITIS (MPA), AND EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS (EGPA).
33	164	PHOSPHOLIPID ANTIBODY-IGM	ELISA	SERUM	SERUM CONTAINER	30 Hours	ANTIPHOSPHOLIPID ANTIBODIES (APLA) IGM ARE THE ANTIBODIES DIRECTED AGAINST PHOSPHOLIPIDS RESPONSIBLE FOR BLOOD CLOTTING. THIS TEST IS TO CHECK FOR THE PRESENCE OF APL THAT CAUSE ABNORMAL BLOOD CLOTTING AND THEIR PRESENCE INDICATES AN AUTOIMMUNE DISORDER CALLED ANTIPHOSPHOLIPID SYNDROME (APS).
34	163	PHOSPHOLIPID Ab-IgG	ELISA	SERUM	SERUM CONTAINER	30 Hours	ANTIPHOSPHOLIPID ANTIBODIES (APLA) IGM ARE THE ANTIBODIES DIRECTED AGAINST PHOSPHOLIPIDS RESPONSIBLE FOR BLOOD CLOTTING. THIS TEST IS TO CHECK FOR THE PRESENCE OF APL THAT CAUSE ABNORMAL BLOOD CLOTTING AND THEIR PRESENCE INDICATES AN AUTOIMMUNE DISORDER CALLED ANTIPHOSPHOLIPID SYNDROME (APS).
35	617	RAPID DENGUE TEST (NS1 Ag & IgM/IgG Ab)	IMMUNO-CHROMATOGRAPHY	SERUM	SERUM CONTAINER	2 Hours	DETECTION OF NS1 ANTIGEN, IGG AND IGM ANTIBODIES OF DENGUE VIRUS IN THE GIVEN SAMPLE.
36	66	RAPID MALARIAL TEST	IMMUNO CHROMATOGRAPHY	EDTA WHOLE BLOOD	SERUM CONTAINER	2 Hours	A RAPID MALARIAL TEST IS AN ALTERNATE WAY OF QUICKLY ESTABLISHING THE DIAGNOSIS OF MALARIA INFECTION BY DETECTING SPECIFIC MALARIA ANTIGENS IN A PERSON'S BLOOD.
37	548	RETRO VIRAL SCREENING CARD	FLOW THROUGH CASSETTE	SERUM	SERUM CONTAINER	2 Hours	SCREENING TEST FOR HIV 1&2 INFECTION.
38	615	RETROVIRAL EARLY SCREEN	CMIA	SERUM	SERUM CONTAINER	2 Hours	HIV TESTING SHOWS WHETHER A PERSON IS INFECTED WITH HIV. HIV STANDS FOR HUMAN IMMUNODEFICIENCY VIRUS. HIV IS THE VIRUS THAT CAUSES AIDS (ACQUIRED IMMUNODEFICIENCY SYNDROME). AIDS IS THE MOST ADVANCED STAGE OF HIV INFECTION.
39	592	RHEUMATOID FACTOR	IMMUNO-TURBIDIMETRY	SERUM	SERUM CONTAINER	2 Hours	RHEUMATOID FACTORS ARE PROTEINS PRODUCED BY THE IMMUNE SYSTEM THAT CAN ATTACK HEALTHY TISSUES IN THE BODY. HIGH LEVELS OF RHEUMATOID FACTOR ARE MOST OFTEN ASSOCIATED WITH AUTOIMMUNE DISEASES, SUCH AS RHEUMATOID ARTHRITIS AND SJOGREN'S SYNDROME.

DEPT NAME: SEROLOGY


Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
40	600	RHEUMATOID FACTOR (SYNOVIAL FLUID)	CMIA	FLUID	AIR TIGHT CONTAINER	3 Hours	RHEUMATOID FACTORS ARE PROTEINS PRODUCED BY YOUR IMMUNE SYSTEM THAT CAN ATTACK HEALTHY TISSUES IN YOUR BODY. HIGH LEVELS OF RHEUMATOID FACTOR ARE MOST OFTEN ASSOCIATED WITH AUTOIMMUNE DISEASES, SUCH AS RHEUMATOID ARTHRITIS AND SJOGREN'S SYNDROME.
41	1180	RUBELLA IgG Ab	CLIA	SERUM	SERUM CONTAINER	5 Hours	DETECTS POST EXPOSURE TO RUBELLA INFECTION(GERMAN MEASLES) OR THE STATUS OF VACCINATION.
42	594	RUBELLA IgM Ab	CLIA	SERUM	SERUM CONTAINER	5 Hours	DETECTS RECENT OR ACUTE PHASE INFECTION WITH RUBELLA VIRUS.
43	1369	SCRUB TYPHUS IGG & IGM RAPID ASSAY	IMMUNO-CHROMATOGRAPHY	SERUM	SERUM CONTAINER	2 Hours	SCRUB TYPHUS, ALSO KNOWN AS BUSH TYPHUS, IS A DISEASE CAUSED BY A BACTERIA CALLED ORIENTIA TSUTSUGAMUSHI (OT). THIS HELPS DETECT IGG IGM ANTIBODIES IN HUMAN SERUM TO OT-DERIVED RECOMBINANT ANTIGEN, WHICH HELP DIAGNOSE HUMAN EXPOSURE TO OT SPECIES.
44	511	TREPONEMA PALLIDUM HEMAGGLUTINATION	CMIA	SERUM	SERUM CONTAINER	4 Hours	THE TREPONEMA PALLIDUM HEMAGGLUTINATION ASSAY IS AN INDIRECT AGGLUTINATION ASSAY USED FOR THE DETECTION AND TITRATION OF ANTIBODIES AGAINST THE CAUSATIVE AGENT OF SYPHILIS.
45	1363	TB GOLD ASSAY (IGRA)	ELISA	BLOOD	LITHIUM HEPARIN VACUTAINER	24 Hours	TB GOLD TEST OR INTERFERON-GAMMA RELEASE ASSAYS (IGRAS) ARE WHOLE-BLOOD TESTS THAT CAN AID IN DIAGNOSING MYCOBACTERIUM TUBERCULOSIS INFECTION. IGRAS MEASURE A PERSON'S IMMUNE REACTIVITY TO M. TUBERCULOSIS. WHITE BLOOD CELLS OF MOST PERSONS THAT HAVE BEEN INFECTED WITH M. TUBERCULOSIS WILL RELEASE HIGH LEVELS OF INTERFERON-GAMMA WHEN MIXED WITH ANTIGENS DERIVED FROM M.TUBERCULOSIS.
46	596	TOXOPLASMA IgG Ab	CLIA	SERUM	SERUM CONTAINER	5 Hours	TOXOPLASMOSIS IS CAUSED BY THE PARASITE TOXOPLASMA GONDI. IT DETECTS PAST INFECTION WITH TOXOPLASMA GONDII.
47	597	TOXOPLASMA IgM Ab	CLIA	SERUM	SERUM CONTAINER	5 Hours	TOXOPLASMOSIS IS CAUSED BY THE PARASITE TOXOPLASMA GONDI. IT DETECTS ACUTE PHASE INFECTION WITH TOXOPLASMA GONDII.


DEPT NAME: SEROLOGY

Sl. No.	Test Code	Test Name	Method	Specimen	Container	TAT	Test Description
48	806	RAPID PLASMA REAGIN (RPR) TEST	SLIDE AGGLUTINATION	SERUM	SERUM CONTAINER	5 Hours	THE RAPID PLASMA REAGIN (RPR) TEST IS A MODIFIED VERSION OF THE VDRL TEST USED TO SCREEN FOR SYPHILIS, CAUSED BY TREPONEMA PALLIDUM. IT USES CARBON PARTICLES FOR EASIER VISUAL INTERPRETATION AND DETECTS NON-SPECIFIC REAGIN ANTIBODIES USING A CARDIOLIPIN ANTIGEN FROM BEEF HEART. SINCE THE TEST IS NON-SPECIFIC, REACTIVE RESULTS SHOULD BE CONFIRMED WITH MORE SPECIFIC TREPONEMAL TESTS LIKE TPHA AND CORRELATED WITH CLINICAL FINDINGS.
49	968	VARICELLA-ZOSTER IgG	CLIA	SERUM	SERUM CONTAINER	5 Hours	VARICELLA ZOSTER IS THE VIRUS THAT CAUSE CHICKEN POX. IGG INDICATES PREVIOUS EXPOSURE OR STATUS OF VACCINATION.
50	535	WIDAL TUBE TEST	TUBE AGGLUTINATION	SERUM	SERUM CONTAINER	24 Hours	THE WIDAL TEST IS AN ADVANCED WAY TO CHECK FOR ANTIBODIES THAT YOUR BODY MAKES AGAINST THE SALMONELLA BACTERIA THAT CAUSES TYPHOID FEVER. IT LOOKS FOR O AND H ANTIBODIES IN A PATIENT'S SERUM.
51	579	WEIL FELIX TEST	TUBE AGGLUTINATION	SERUM	SERUM CONTAINER	48 Hours	THE WEIL-FELIX TEST IS AN AGGLUTINATION TEST FOR THE DIAGNOSIS OF RICKETTSIAL INFECTIONS. THE BASIS OF THE TEST IS TO DETECT THE ANTI-RICKETTSIAL ANTIBODIES PRESENT IN THE PATIENT'S SERUM SAMPLE.



Scan the QR code with your smartphone's camera, tap the link, and install our mobile collection app from play store





PACKAGES

1. Aswini Care Basic Health Checkup
2. Aswini Wellness Health Checkup - Male
3. Aswini Wellness Health Checkup - Female
4. Aswini Wellness Premium Health Checkup



aswini
DIAGNOSTIC SERVICES
A UNIT OF ADS RESEARCH CENTRE LLP
DIRECTORY OF SERVICES



An ISO 15189: 2022 Laboratory Accredited to National Accreditation Board for Testing & Calibration Laboratories (NABL)

Jail Road, Calicut-673004.

www.aswinicalicut.net, mail@aswinicalicut.net