

Minimizing the central circle in a ring of circles of radii $1, 2, \dots, n$: Supnick orders, floating circles, and a breakdown cascade

Maurizio Falconi*

June 2026 — v4.2 (submission)

Abstract

Non-overlapping circles of radii $1, 2, \dots, n$ must all be externally tangent to a central circle. In what cyclic order should they be arranged so that the radius R of the central circle is minimized? The question was posed by Dan (Mathematics Stack Exchange, January 2023) together with a conjectured optimal arrangement, and is listed as problem 295 of the UQ benchmark of unsolved questions. We resolve the structure of the problem. First, we show that the angular-separation matrix $\theta_R(a, b) = 2 \arcsin \sqrt{ab/((R+a)(R+b))}$ is, for every R , a symmetric anti-Monge (inverse Supnick) matrix; by a classical theorem of Supnick (1957) the cyclic order minimizing the total angle is therefore a single fixed tour, independent of R , namely $\sigma^* = \langle 1, n-1, 3, n-3, 5, \dots, n-2, 2, n \rangle$. This proves the conjectured “pyramid” arrangement optimal whenever the resulting necklace is geometrically realizable, and yields the unconditional lower bound $R^*(n) \geq R_{\text{chain}}(\sigma^*)$. Second, realizability fails in a structured cascade: from $n = 8$ the smallest circle can no longer separate its two large neighbours and becomes a *floating* circle; from $n = 13$ the same mechanism reaches circle 2 inside the chain. We formalize the problem as a circular system of pairwise angular constraints (a simple temporal network), give an exact certified branch-and-bound, and determine the optimal arrangements for all $n \leq 14$, with optima verified to 50 decimal digits. The optimal radii interpolate four structural regimes which we characterize exactly via Descartes’ Circle Theorem. The cascade continues self-similarly: circle 2 leaves the chain at $n = 14$ (certified) and, heuristically, circle 3 by $n = 18$. We further conjecture, with supporting analysis, that $R^*(n) = \frac{n^2}{8} (1 + o(1))$, and we determine the radius-*maximizing* arrangement as the complementary Supnick tour $\langle 1, 3, 5, \dots, n, \dots, 4, 2 \rangle$.

1 Introduction

The following question was asked by the user Dan on Mathematics Stack Exchange in January 2023 [1] and, despite substantial attention (27 votes), has remained unanswered; it appears as question 295 in the UQ benchmark of unsolved Stack Exchange questions [2].

Non-overlapping circles of radius $1, 2, 3, \dots, n$ are all externally tangent to a middle circle. How should we arrange the surrounding circles in order to minimize the middle circle’s radius R ?

*Independent researcher, Parma, Italy. maurizio.falconi47@gmail.com

Dan conjectured a specific “pyramid-fashion” arrangement and observed two phenomena that make the problem subtle: (i) the natural closure equation

$$\sum_{k=1}^n \arccos\left(1 - \frac{2r_k r_{k+1}}{(R + r_k)(R + r_{k+1})}\right) = 2\pi \quad (r_{n+1} = r_1) \quad (1)$$

treats consecutive circles as mutually tangent, yet (ii) for $n \geq 8$ the smallest circles can be “placed last anywhere they fit”, i.e. the optimal configuration contains circles tangent to the central circle only. A comment by Rei Henigman [1] verified the conjecture for $n = 10$ by enumerating permutations of $\{2, \dots, 10\}$, manually excluding circle 1 from the chain.

This paper resolves the structure of the problem on three levels.

1. The ordering problem is a Supnick TSP. Writing $\theta_R(a, b)$ for the minimum angular separation, at central radius R , between the centers of tangent circles of radii a and b (Section 2), we show that the matrix $(\theta_R(r_i, r_j))_{i,j}$ is symmetric *anti-Monge* for every $R > 0$ (Lemma 5). Minimizing total angle over cyclic orders is therefore a *maximum* traveling salesman problem on a Supnick matrix, which by a classical theorem of Supnick [3] (see also the survey [4]) is solved by one fixed tour independent of the matrix entries:

$$\sigma^* = \langle 1, n-1, 3, n-3, 5, n-5, \dots, n-2, 2, n \rangle.$$

A short self-consistency argument (the same order is optimal at *every* R) transfers optimality from fixed R to the variable- R problem (Theorem 6). The order σ^* coincides with Dan’s conjectured pyramid arrangement. In particular $R_{\text{chain}}(\sigma^*)$, defined by (1), is an unconditional lower bound for $R^*(n)$, and it is the exact optimum whenever the σ^* -necklace is geometrically realizable.

2. Realizability breaks in a cascade, which we determine exactly. The closure equation (1) ignores non-adjacent pairs. For $n \geq 8$ the configuration it describes for σ^* is *not realizable*: circle 1, seated between circles n and $n-1$, no longer keeps them apart, i.e. $\theta_R(n, 1) + \theta_R(1, n-1) < \theta_R(n, n-1)$. Circle 1 then leaves the chain and becomes a *floating* circle, tangent to the central circle only. We prove that the correct model is a circular system of $\binom{n}{2}$ pairwise angular constraints — equivalently a simple temporal network (STN) — and we give an exact, certified algorithm for it (Section 4). Exhaustive certified optimization for $3 \leq n \leq 13$ (Section 5) reveals four regimes, governed by two closed-form quantities of the Supnick tour on $\{2, \dots, n\}$: its geometric validity and the largest Descartes pocket it opens.

regime	n	floating set	optimal chain
I (full necklace)	$3 \leq n \leq 7$	$\emptyset$	σ^* on $\{1, \dots, n\}$
II (paid floater)	$n = 8, 9$	$\{1\}$	distorted tour on $\{2, \dots, n\}$
III (free floater)	$n = 10, 11, 12$	$\{1\}$	σ^* on $\{2, \dots, n\}$
IV (second breakdown)	$n = 13$	$\{1\}$	constrained tour on $\{2, \dots, n\}$
II' (cascade, level 2)	$n = 14$	$\{1, 2\}$	constrained tour on $\{3, \dots, n\}$

3. Quantitative results. We report certified optima, binding structures and floating sets for all $n \leq 13$, verified to 50 decimal digits (Table 1); the radius-maximizing (worst) arrangement, which is the *minimum*-tour counterpart $\langle 1, 3, 5, \dots, n, \dots, 4, 2 \rangle$ of Supnick’s theorem (Proposition 8); and the asymptotic law $R^*(n) \sim n^2/8$ (Section 6).

Related work. Supnick’s fixed-tour theorem [3] and the Monge/Supnick taxonomy of polynomially solvable TSPs are surveyed in [4]. Circle packing literature concerns containers (packing circles *inside* regions, see e.g. [6]); the present problem — ordering unequal circles *around* a central one — appears not to have been treated. The tangency mechanics are governed by Descartes’ Circle Theorem [5]. To our knowledge the link between Supnick matrices and rings of tangent circles is new.

2 The model

Let the central circle have radius $R > 0$ and center O . A surrounding circle of radius r tangent externally to it has its center at distance $R + r$ from O ; its position is a single angle φ . The whole configuration is a vector of angles.

Lemma 1 (Exact angular reformulation). *Circles of radii a, b , both externally tangent to the central circle, with angular positions separated by $\Delta \in [0, \pi]$, are non-overlapping if and only if*

$$\Delta \geq \theta_R(a, b) := 2 \arcsin \sqrt{\frac{ab}{(R+a)(R+b)}}.$$

Moreover $\theta_R(a, b) \in (0, \pi)$ always, θ_R is symmetric, strictly decreasing in R , and strictly increasing in each radius.

Proof. By the law of cosines the squared distance between the centers is $(R+a)^2 + (R+b)^2 - 2(R+a)(R+b)\cos\Delta$, and non-overlap means this is at least $(a+b)^2$. Rearranging gives $\cos\Delta \leq 1 - \frac{2ab}{(R+a)(R+b)}$, i.e. $\Delta \geq \theta_R(a, b)$ with θ_R as stated, using $\arccos(1-2s^2) = 2 \arcsin s$ for $s \in [0, 1]$. The argument of the square root is strictly less than 1 since $(R+a)(R+b) - ab = R(R+a+b) > 0$, whence $\theta_R < \pi$; the monotonicity claims are immediate from the formula. $\square$

Thus the problem is: minimize R such that there exist angles $\varphi_1, \dots, \varphi_n$ with circular separations $\geq \theta_R(r_i, r_j)$ for *all* pairs. For a fixed cyclic order this is a circular system of difference constraints — a simple temporal network — and feasibility at fixed R is decidable by shortest paths (Section 4). Two derived quantities recur:

Definition 2. For a cyclic order σ of a multiset of radii, $R_{\text{chain}}(\sigma)$ is the unique solution of $\sum_i \theta_R(\sigma_i, \sigma_{i+1}) = 2\pi$. (Existence and uniqueness: the left side tends to $|\sigma|\pi \geq 2\pi$ as $R \rightarrow 0^+$, to 0 as $R \rightarrow \infty$, and is strictly decreasing.) For a feasible pair (order, R), summing the adjacent gaps shows $\sum_i \theta_R(\sigma_i, \sigma_{i+1}) \leq 2\pi$ and hence $R \geq R_{\text{chain}}(\sigma)$: the chain value is an unconditional per-order lower bound.

Definition 3 (Floating circle). A circle c is *floating* at a global optimum if there exists an optimal configuration in which c is tangent only to the central circle (every pairwise constraint involving c is strictly slack).

Definition 4 (Descartes pocket). For adjacent, mutually tangent circles a, b on the necklace at central radius R , the *pocket* is the largest circle externally tangent to the central circle and to both: by Descartes’ Circle Theorem its curvature is $k_{\text{pocket}} = \frac{1}{R} + \frac{1}{a} + \frac{1}{b} + 2\sqrt{\frac{1}{Ra} + \frac{1}{ab} + \frac{1}{Rb}}$. A circle of radius ρ floats freely between a and b iff $\rho \leq 1/k_{\text{pocket}}$, equivalently $\theta_R(a, \rho) + \theta_R(\rho, b) \leq \theta_R(a, b)$.

3 The optimal cyclic order is a fixed Supnick tour

Lemma 5 (Supermodularity). *For every $R > 0$ the matrix $(\theta_R(r_i, r_j))$, rows and columns indexed by the radii in increasing order, is symmetric strictly anti-Monge: for $a < a'$ and $b < b'$,*

$$\theta_R(a, b) + \theta_R(a', b') > \theta_R(a, b') + \theta_R(a', b).$$

Proof. Write $\theta_R(a, b) = 2 \arcsin(t_a t_b)$ with $t_x = \sqrt{x/(R+x)} \in (0, 1)$, which is strictly increasing in x . For $f(x, y) = \arcsin(xy)$ on $(0, 1)^2$,

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial y} \frac{y}{\sqrt{1-x^2 y^2}} = \frac{(1-x^2 y^2) + x^2 y^2}{(1-x^2 y^2)^{3/2}} = (1-x^2 y^2)^{-3/2} > 0,$$

so f is strictly supermodular; supermodularity is preserved by strictly increasing reparametrizations of each coordinate, and the claim follows. $\square$

Equivalently, $-\theta_R$ is a Supnick matrix (symmetric Monge). Supnick's theorem [3], in the form stated in [4, Sec. 2], says that on Supnick matrices the minimum TSP tour is $\langle 1, 3, 5, \dots, n, \dots, 6, 4, 2 \rangle$ and the *maximum* TSP tour is

$$\sigma^* = \langle 1, n-1, 3, n-3, 5, n-5, \dots, n-2, 2, n \rangle,$$

independently of the matrix entries. Maximizing $\sum(-\theta_R)$ is minimizing $\sum \theta_R$, hence:

Theorem 6 (Optimal order). *Let $r_1 < \dots < r_n$ be arbitrary positive radii.*

- (i) *For every $R > 0$, among all cyclic orders the tour σ^* minimizes $\sum_i \theta_R(r_{\sigma(i)}, r_{\sigma(i+1)})$.*
- (ii) *σ^* minimizes R_{chain} over all cyclic orders; hence $R^* \geq R_{\text{chain}}(\sigma^*)$ unconditionally.*
- (iii) *If the σ^* -necklace at $R_{\text{chain}}(\sigma^*)$ is geometrically realizable (no non-adjacent pair violated), then it is a global optimum: $R^* = R_{\text{chain}}(\sigma^*)$.*

Proof. (i) is Lemma 5 plus Supnick's theorem. (ii): fix any order σ and let $R = R_{\text{chain}}(\sigma)$. By (i), $\sum \theta_R(\sigma^*) \leq \sum \theta_R(\sigma) = 2\pi$; since $\sum \theta_R(\sigma^*)$ is strictly decreasing in R and equals 2π at $R_{\text{chain}}(\sigma^*)$, we get $R_{\text{chain}}(\sigma^*) \leq R$. The unconditional bound follows since every feasible (σ, R) has $R \geq R_{\text{chain}}(\sigma)$. (iii): realizability makes the lower bound attained. $\square$

Remark 7. Theorem 6 holds for *arbitrary* distinct radii, not only $1, \dots, n$; the arithmetic sequence enters only in the realizability analysis below. For $n = 10$ the lower bound of (ii) is ≈ 9.90 , the value observed in [1] when circle 1 is (incorrectly) kept in the chain; the true optimum is $9.9799\dots$

Proposition 8 (Worst arrangement). *For every R the cyclic order maximizing $\sum_i \theta_R$ is the complementary Supnick tour $\sigma_{\dagger} = \langle 1, 3, 5, \dots, n, \dots, 6, 4, 2 \rangle$; consequently $\sigma_{\dagger}$ maximizes R_{chain} . For all $3 \leq n \leq 13$ the $\sigma_{\dagger}$ -necklace is realizable, so among necklace arrangements it maximizes the central radius exactly; certified values in Table 3.*

4 Exact algorithm and certification

Feasibility oracle. Fix a cyclic order and R . Unrolling the circle at one circle ($\varphi_0 = 0$) gives, for every pair $i < j$ of positions, the two constraints $\varphi_j - \varphi_i \geq \theta_R(r_i, r_j)$ and $\varphi_j - \varphi_i \leq 2\pi - \theta_R(r_i, r_j)$. This is a simple temporal network; feasibility is the absence of a negative cycle in the associated distance graph and is decided by Floyd–Warshall in $O(n^3)$. The minimum feasible R for the order, $R_{\text{full}}(\sigma)$, is found by bisection (the constraint set relaxes monotonically in R), bracketed below by $R_{\text{chain}}(\sigma)$. A witness configuration is recovered from shortest-path potentials and verified independently in Cartesian coordinates.

Lower bounds and search. For the search over orders we use

$$\text{LB}(\sigma) = \max \{ R_{\text{chain}}(\sigma), R_{\text{chain}}(\sigma \setminus \{1\}), R_{\text{chain}}(\sigma \setminus \{1, 2\}) \},$$

where $\sigma \setminus S$ denotes the induced cyclic order (valid since induced adjacencies are a subset of the full pairwise constraints). Orders are enumerated canonically (largest radius pinned, reflections halved), LB is computed vectorized for all $(n-1)!/2$ orders, and R_{full} is evaluated in ascending LB until the next bound meets the incumbent: this certifies global optimality up to an absolute guard of 10^{-10} in R . Optimal binding structures were then re-verified in 50-digit arithmetic (mpmath); the values in Table 1 are exact to all displayed digits, with maximal residual slack below 10^{-40} . The $n = 13$ certification enumerates $12!/2 \approx 2.4 \times 10^8$ orders (≈ 1.8 h on a consumer 8-core laptop); $n = 14$ enumerates $13!/2 \approx 3.1 \times 10^9$ orders (≈ 22 h). Code and artifacts: <https://github.com/falker47/ringmin>.

Essential binding structure. A pairwise constraint is *essential* if perturbing it alone by $+\varepsilon$ makes the STN infeasible at R^* ; a circle is floating iff it carries no essential constraint. This distinguishes true tangencies from placement artifacts of a particular witness.

5 Certified optima for $n \leq 14$ and the regime cascade

For $n \leq 7$ the optimal cycle is exactly σ^* on $\{1, \dots, n\}$ (regime I), as guaranteed by Theorem 6(iii) once realizability is checked. The transitions are governed by two computable properties of the Supnick tour on the reduced set $\{2, \dots, n\}$:

Proposition 9 (First breakdown). *In the σ^* -necklace on $\{1, \dots, n\}$, circle 1 sits between n and $n-1$. The realizability inequality $\theta_R(n, 1) + \theta_R(1, n-1) \geq \theta_R(n, n-1)$ at $R = R_{\text{chain}}(\sigma^*)$ holds for $n \leq 7$ and fails for $8 \leq n \leq 13$ (violated pair $(n, n-1)$; at $n = 13$ also $(13, 11)$ across circle 2). Hence regime I ends at $n = 7$. We conjecture failure for all $n \geq 8$; cf. open problem (1).*

Proposition 10 (Free vs. paid floater). *Let $\rho_{\text{max}}(n)$ be the largest Descartes pocket of the σ^* -necklace on $\{2, \dots, n\}$ at its chain radius. Then $\rho_{\text{max}} = 0.858, 0.988, 1.139, 1.262, 1.409$ for $n = 8, \dots, 12$. For $n = 10, 11, 12$ one has $\rho_{\text{max}} > 1$: circle 1 floats freely and $R^*(n) = R_{\text{chain}}(\sigma^* \text{ on } \{2, \dots, n\})$ exactly (regime III). For $n = 8, 9$, $\rho_{\text{max}} < 1$: the optimum distorts the reduced tour to open a unit pocket, at certified extra cost $\delta(8) = 0.04080$, $\delta(9) = 0.00532$ over the (then unattainable) reduced chain bound (regime II).*

Proposition 11 (Second breakdown). *At $n = 13$ the Supnick tour on $\{2, \dots, 13\}$ is itself unrealizable: circle 2, seated between 13 and 12, violates $\theta_R(13, 2) + \theta_R(2, 12) = 0.80448 <$*

Table 1: Certified optima, rounded to 12 decimals from the 50-digit verified values (30-digit table in the Appendix). Cycles list radii in circular order; the floating circle is shown in its (non-unique) hosting gap. “Essential pair” is the tangent non-adjacent pair bridged by the floater.

n	$R^*(n)$	optimal cycle	floating	essential pair
3	0.260869565217 (= 6/23)	(3,1,2)	—	—
4	0.844453589561	(4,1,3,2)	—	—
5	1.695494081203	(5,1,4,3,2)	—	—
6	2.794919518897	(6,1,5,3,4,2)	—	—
7	4.153189553744	(7,1,6,3,4,5,2)	—	—
8	5.767794284590	(8,1,6,4,5,3,7,2)	{1}	(8, 6)
9	7.726726552611	(9,2,8,1,5,6,4,7,3)	{1}	(8, 5)
10	9.979907385863	(10,2,9,4,7,1,6,5,8,3)	{1}	(7, 6)
11	12.488720487188	(11,2,10,4,8,6,7,1,5,9,3)	{1}	(7, 5)
12	15.258870430448	(12,2,11,4,9,6,7,8,5,1,10,3)	{1}	(5, 10)
13	18.317563047217	(13,3,1,12,2,10,6,8,7,9,5,11,4)	{1}	(3, 12)
14	21.665395182215	(14,3,13,2,9,8,7,10,6,11,5,1,12,4)	{1, 2}	(13, 9), (5, 12)

$0.83488 = \theta_R(13, 12)$ at $R = 18.3176$. The certified optimum demotes circle 2 to the pair (12, 10), where the separation constraint holds with slack 6×10^{-3} rad; circle 1 floats in the (3, 12) pocket (radius 1.086). Thus the mechanism that expelled circle 1 at $n = 8$ reaches circle 2 at $n = 13$ (regime IV).

Remark 12 (Structural summary). For all $n \leq 13$ the optimum has the form: a single tangent necklace carrying a Supnick-like order *constrained to geometric realizability*, plus floating circles in Descartes pockets. The cascade continues self-similarly: circle 2 first floats at $n = 14$ (now *certified*, Table 1) via a *paid* distortion that opens a pocket of radius 2.013 between circles 13 and 9 — the regime-II mechanism one level up, at extra cost 0.0565 over the (then unattainable) chain bound of the Supnick tour on $\{3, \dots, 14\}$. At $n = 16$ the best known value coincides to 10^{-10} with the chain value of the Supnick tour on $\{3, \dots, 16\}$, the level-2 analogue of regime III. Heuristically, circle 3 joins by $n = 18$ (Table 2). Moreover the Supnick tour on $\{3, \dots, n\}$ itself becomes unrealizable at $n = 17$ — circle 3 failing at the (17, 16) seam with slack -0.0104 — so the *seam-failure* onsets for circles 1, 2, 3 are $n = 8, 13, 17$: each cascade level reproduces the full regime pattern of the previous one. The onsets $n = 8, 14, 18$ for floaters 1, 2, 3 are consistent with Dan’s $n/4$ Descartes-based upper bound on the number of ignorable circles [1].

Table 2: Best known arrangements for $15 \leq n \leq 18$ (local search, 200–600 restarts per n ; *non-exhaustive*). The last column counts restarts reaching the best value.

n	best R	floating	cycle	hits
15	25.290151636714	{1, 2}	(15,3,14,5,12,2,9,1,8,10,7,11,6,13,4)	59
16	29.194988498141	{1, 2}	(16,1,4,14,6,12,8,10,9,2,11,7,13,5,15,3)	18
17	33.394425022062	{1, 2}	(17,4,16,3,15,6,1,13,8,11,10,9,2,12,7,14,5)	19
18	37.870755667393	{1, 2, 3}	(18,4,17,3,14,8,12,10,11,9,2,13,7,15,6,16,1,5)	9

6 Asymptotics

For $R \gg n$ one has $\theta_R(a, b) = 2\sqrt{ab}/R(1 + O(n/R))$, so the closure equation gives $R \approx \frac{1}{\pi} \sum_i \sqrt{r_i r_{i+1}}$. Since $\sqrt{ab}$ is supermodular, Theorem 6(i) applies verbatim: the minimizing order of $\sum \sqrt{r_i r_{i+1}}$ is again σ^* , whose value on $1, \dots, n$ is

$$\min_{\sigma} \sum_i \sqrt{r_{\sigma(i)} r_{\sigma(i+1)}} = (1 + o(1)) \int_0^n \sqrt{x(n-x)} dx = \frac{\pi n^2}{8} (1 + o(1)),$$

because adjacent pairs in σ^* are $\{k, n - k + O(1)\}$ with each odd k contributing twice. Self-consistency ($R \sim n^2/8 \gg n$) further requires the floating circles to be asymptotically negligible, in the sense $\sum_{k \in F(n)} \sqrt{kn} = o(n^2)$ — guaranteed if $|F(n)| = o(n)$, consistent with $|F(18)| = 3$, but open (problem (2) below). Under this assumption we are led to:

Conjecture 13. $R^*(n) = \frac{n^2}{8}(1 + o(1))$. Empirically the deficit $n^2/8 - R(n)$ equals 2.52, 2.64, 2.74, 2.81, 2.83 for the certified $n = 10, \dots, 14$, and 2.83, 2.81, 2.73, 2.63 for the heuristic $n = 15, \dots, 18$ (the latter are lower bounds on the true deficit, heuristic radii being upper bounds for R^*): bounded so far, and in any case growing strictly slower than n . We tentatively conjecture the stronger form $n^2/8 - R^*(n) = O(\sqrt{n})$.

Two-sided bounds making the leading order rigorous (via $\arcsin x \geq x$ and the explicit σ^* configuration as an upper bound) appear within reach but are not pursued here.

7 Open problems

(1) Prove Propositions 9–11 for all n analytically (the inequalities are explicit; monotonicity in n is the missing step). (2) Characterize the floating set $F(n)$ asymptotically (float onsets: circle 1 at $n = 8$ and circle 2 at $n = 14$, both certified, circle 3 at $n = 18$ heuristically; certified seam-failure onsets $n = 8, 13, 17$); is $|F(n)| \sim cn^{1/2}$, or does Dan’s $n/4$ bound give the right order? (3) The same machinery applies to radii $r_k = k^\alpha$ or arbitrary sequences: Theorem 6 is general, the cascade analysis is not. (4) Three dimensions: spheres of radii $1, \dots, n$ tangent to a central sphere — the ordering structure disappears; what replaces Supnick?

Acknowledgments

The problem and the pyramid conjecture are due to Dan [1]; Rei Henigman’s $n = 10$ computation provided an independent check. This work was produced in an AI-assisted workflow: problem selection, research design, the mathematical arguments, and supervision of all computations were carried out with substantial assistance from Anthropic’s Claude; the software implementation was written by OpenAI’s Codex under that supervision, with the author acting as orchestrator. All mathematical claims were reviewed as stated, and all certified values were independently re-verified in 50-digit arithmetic as described in Section 4.

References

- [1] Dan, *Circles of radius $1, 2, 3, \dots, n$ all touch a middle circle. How to make the middle circle as small as possible?*, Mathematics Stack Exchange, question 4619480, January 2023. <https://math.stackexchange.com/q/4619480>
- [2] The UQ Project, *Unsolved Questions* benchmark, question 295. <https://uq.stanford.edu/question/295> (accessed June 2026).
- [3] F. Supnick, *Extreme Hamiltonian lines*, Annals of Mathematics (2) **66** (1957), 179–201.
- [4] R.E. Burkard, V.G. Deineko, R. van Dal, J.A.A. van der Veen, G.J. Woeginger, *Well-solvable special cases of the TSP: a survey*, SIAM Review **40**(3) (1998), 496–546.
- [5] J.C. Lagarias, C.L. Mallows, A.R. Wilks, *Beyond the Descartes circle theorem*, Amer. Math. Monthly **109** (2002), 338–361.
- [6] M. Hifi, R. M’Hallah, *A literature review on circle and sphere packing problems: models and methodologies*, Advances in Operations Research (2009).

A High-precision values and figures

Table 3: The radius-maximizing necklace arrangement $\sigma_{\dagger}$ (Proposition 8): certified chain values, all geometrically realizable.

n	3	4	5	6	7	8	9	10	11	12	13
$R_{\dagger}$	0.2609	0.8673	1.7786	3.0001	4.5375	6.3934	8.5690	11.0647	13.8806	17.0164	20.4720

Per- n configuration figures for all $3 \leq n \leq 18$ are available in the repository.

n	R^*
3	0.260869565217391304347826086957
4	0.844453589560855604347528524674
5	1.69549408120271081351328017371
6	2.79491951889692485617024406797
7	4.15318955374381246513863858202
8	5.7677942845896143026361805725
9	7.72672655261128921886246177604
10	9.97990738586347760966552641468
11	12.4887204871876588517468786264
12	15.2588704304484933617250043503
13	18.3175630472173206282821941532
14	21.6653951822145150956462891793

Table 4: Certified optima, 30 significant digits.

n	Supnick min tour	R_{chain}	R_{full}
3	[1, 3, 2]	0.260869565217	0.260869565217
4	[1, 3, 4, 2]	0.867281640255	0.867281640255
5	[1, 3, 5, 4, 2]	1.778550949884	1.778550949884
6	[1, 3, 5, 6, 4, 2]	3.000138442323	3.000138442323
7	[1, 3, 5, 7, 6, 4, 2]	4.537467833562	4.537467833562
8	[1, 3, 5, 7, 8, 6, 4, 2]	6.393358320114	6.393358320114
9	[1, 3, 5, 7, 9, 8, 6, 4, 2]	8.568988935955	8.568988935955
10	[1, 3, 5, 7, 9, 10, 8, 6, 4, 2]	11.064726494858	11.064726494858
11	[1, 3, 5, 7, 9, 11, 10, 8, 6, 4, 2]	13.880577932626	13.880577932626
12	[1, 3, 5, 7, 9, 11, 12, 10, 8, 6, 4, 2]	17.016408238128	17.016408238128
13	[1, 3, 5, 7, 9, 11, 13, 12, 10, 8, 6, 4, 2]	20.472039120681	20.472039120681
14	[1, 3, 5, 7, 9, 11, 13, 14, 12, 10, 8, 6, 4, 2]	24.247291190092	24.247291190092
15	[1, 3, 5, 7, 9, 11, 13, 15, 14, 12, 10, 8, 6, 4, 2]	28.342000288518	28.342000288518
16	[1, 3, 5, 7, 9, 11, 13, 15, 16, 14, 12, 10, 8, 6, 4, 2]	32.756022334220	32.756022334220
17	[1, 3, 5, 7, 9, 11, 13, 15, 17, 16, 14, 12, 10, 8, 6, 4, 2]	37.489233291550	37.489233291550
18	[1, 3, 5, 7, 9, 11, 13, 15, 17, 18, 16, 14, 12, 10, 8, 6, 4, 2]	42.541527273519	42.541527273519

Table 5: Worst-arrangement comparison; Supnick min tour values.




Figure 1: $R^*(n)$, the worst-arrangement curve, and the $n^2/8$ asymptote.




Figure 2: The certified optimum for $n = 14$; floating circles 1, 2 highlighted.