

Tomato yellow leaf curl virus disease on tomato - Ghana

Tomato yellow leaf curl virus TLCV

	Prevention	Monitoring	Direct Control	Direct Control	Restrictions
 Stunting/dwarfism of young leaves. Source: Scot Nelson, via Flickr	- Whiteflies carry this disease, so don't plant new crops close to infected ones. - Prevent build-up of the whitefly population whenever possible. - Tomato is the main host, but avoid growing crops like okra, peppers beans or cucumbers/melons close to tomato fields.	- TYLCV causes severe stunting of young leaves and shoots, resulting in bushy growth of infected seedlings. - Symptoms are clear at the seedling stage. - Use yellow sticky traps to monitor the presence of whiteflies. If a lot of whiteflies are seen or if TYLCV symptoms are seen take action.	- Remove infected seedlings and destroy by burning or burying. - If possible, use reflective mulches (Aluminium/silver coloured) between rows to discourage feeding of whiteflies. - Spray with 0.25% or 0.5% horticultural oil (2-4 teaspoons/ 4.5L of water) with horticultural soap.	- When using a pesticide, always wear protective clothing and follow the instructions on the product label. - Do not use chemicals with the same mode of action year after year as this can lead to resistance. - Always consult the most recent list of registered pesticides (PCPB).	WHO Class II (Moderately hazardous). IRAC 4A. REI 24hrs; PHI 1 day.
 Yellowing and curling of leaves. Source: Scot Nelson, via Flickr	- Remove Billygoat weed which is an alternative host of the virus. - Remove and destroy old crop residues on which the whitefly are feeding and all volunteer tomatoes in other crops. - Plant early to avoid peak of the whitefly population and protect the nursery beds from whitefly with nets. - Plant resistant varieties if available.	- Look out for stunting and dwarfism, upward and inward curling of leaves with yellow margins. - Yellowing of young leaves. - Flower drops, random distribution of affected plants in patches.		- Thiamethoxam at 30 g/15 l for 2 consecutive times at 14 day intervals. - Lambda cyhalothrin 5 EC (45ml/15L). Systemic pyrethroid with contact activity. Maximum 2 applications in the morning and later in the day.	- Don't use both imidacloprid & thiamethoxam. WHO Class II; IRAC 4A. PHI 3 days. Toxic to bees and other non-target arthropods. - WHO Class II. IRAC 3A. PHI 2 days. REI 24 hrs. Eye and skin irritant. Highly toxic to bees and other non-target arthropods. Toxic to aquatic organisms; avoid using near waterways.

Ghana

CREATED/UPDATED: February 2023

AUTHOR(S): Grace Vanderpujje, Emmanuella Lekete, Awudu Gariba, Joseph Honger, Andrew Sarkordie

EDITED BY: Plantwise

LOSE LESS, FEED MORE

PlantwisePlus is a CABI-led global initiative www.plantwise.org
©CAB International. Published under a CC BY-NC-SA 4.0 licence