

# Asteroseismology of GD358 with complex C/O core profiles

Agnès Bischoff-Kim<sup>1</sup>, J. L. Provencal<sup>2,3</sup>, M. H. Montgomery<sup>3,4</sup>, Samuel T. Harrold<sup>4</sup>, B. Howard<sup>3</sup>, W. Strickland<sup>5</sup>, D. Chandler<sup>5</sup>, D. Campbell<sup>5</sup>, A. Arredondo<sup>5</sup>, R. Linn<sup>5</sup>, D. P. Russell<sup>5</sup>, D. Doyle<sup>5</sup>, A. Brickhouse<sup>5</sup>, R. Linn<sup>5</sup>, D. Peters<sup>5</sup>, A. V. Kusakin<sup>6</sup>, A. V. Sergeev<sup>7</sup>, M. Andreev<sup>7</sup>, A. Maksim<sup>7</sup>, S. Velichko<sup>7</sup>, R. Janulis<sup>8</sup>, F. Aliçavuş<sup>9</sup>, N. Horoz<sup>9</sup>, S. Zola<sup>10,11</sup>, W. Ogłóza<sup>10,11</sup>, D. Koziel-Wierzbowska<sup>10,11</sup>, T. Kundera<sup>10,11</sup>, D. Jableka<sup>10,11</sup>, B. Debski<sup>10,11</sup>, S. Meingast<sup>12</sup>

## ABSTRACT

We report on .....

*Subject headings:* Stars: oscillations — Stars: variables: general — white dwarfs

## 1. Astrophysical Context

White dwarfs are the end product of evolution for around 98% of the stars in our Galaxy. Buried in their interiors are the records of physical processes that take place during earlier

---

<sup>1</sup>Penn State Worthington Scranton, Dunmore, PA 18512; axk55@psu.edu

<sup>2</sup>University of Delaware, Department of Physics and Astronomy Newark, DE 19716; jlp@udel.edu

<sup>3</sup>Delaware Asteroseismic Research Center, Mt. Cuba Observatory, Greenville, DE 19807

<sup>4</sup>Department of Astronomy, University of Texas, Austin, TX 78712; mikemon@rocky.as.utexas.edu

<sup>5</sup>Meyer Observatory and Central Texas Astronomical Society, 209 Paintbrush, Waco, TX 76705; chandler@vvm.com

<sup>6</sup>Fesenkov Astrophysical Institute, Almaty 050020, Kazakhstan

<sup>7</sup>Ukrainian National Academy of Sciences, Main Astronomical Observatory, Golosiiv, Kiev 022 252650; sergeev@terskol.com

<sup>8</sup>Institute of Theoretical Physics and Astronomy, Vilnius University, Vilnius, Lithuania; jr@itpa.lt

<sup>9</sup>Ulupinar Observatory, Çanakkale Onsekiz Mart University, Turkey

<sup>10</sup>Astronomical Observatory of the Jagiellonian University, ul. Orla 171, 30-244 Cracow, Poland; szola@oa.uj.edu.pl

<sup>11</sup>Mount Suhora Observatory, Cracow Pedagogical University, Ul. Podchorazych 2, 30-084 Krakow, Poland; zola@astro1.as.ap.krakow.pl

<sup>12</sup>Institut für Astrophysik, Universität Wien, Türkenschanzstrasse 17, 1180 Wien, Austria

stages in the life of the star. Nuclear reaction rates during the core helium burning phase set the core composition of white dwarfs, while the relative time spent burning hydrogen and helium during the asymptotic-giant-branch (AGB) phase and mass-loss episodes determine the thickness of the helium layer (Lawlor & MacDonald 2006; Althaus et al. 2005). Helium atmosphere white dwarfs (DBs) comprise roughly 20% of the population of field white dwarfs, with most of the remaining 80% consisting of their hydrogen atmosphere (DA) cousins. The bifurcation into two spectral classes is thought to occur during post-AGB evolution when, in some cases, a very late thermal pulse burns off the residual hydrogen in the envelope, producing a nearly pure helium atmosphere (Iben et al. 1983). Such objects are then supposed to return to the white dwarf cooling track as PG 1159 stars, which are widely believed to be the precursors of most DB white dwarfs. DBs are found to pulsate at effective temperatures ranging between 21,000 K and 28,000 K (Beauchamp et al. 1999; Castanheira et al. 2005).

The subject of this paper, GD358 (V777 Her) is the brightest ( $m_v = 13.7$ ) and best studied helium atmosphere white dwarf pulsator. It is located near the red edge of the instability strip, with  $T_{\text{eff}}$  of 24000 K and  $\log g$  of 7.8 (Koester 2013). GD358’s pulsation spectrum contains a series of independent modes, many with complex frequency structure. Models involving magnetic fields and oblique rotation have been proposed to explain that structure (Montgomery et al. 2010).

Pulsating white dwarfs do not pulsate in all their available modes at all times. We have maintained an active observing program of this complex star in an effort to identify additional frequencies. These new observations have identified additional periods in GD358’s pulsation spectrum, bringing the total known modes to 15. Thirteen of these modes belong to a consecutive  $\ell = 1$  sequence, the longest sequence observed in a DBV. This sequence is critical to constrain the asteroseismic fits and better define the limitations of high precision white dwarf asteroseismology.

Paradoxically, among the DBVs that have enough detected periods to be fitted asteroseismically, GD358 is the only one that has not been analyzed using the complex C/O profiles adapted and parameterized from stellar evolution calculations (e.g. Salaris et al. (1997); Althaus et al. (2005)). The most recent fits of GD358 (Metcalf et al. 2003) were performed using 11 observed modes and simple models where the oxygen abundance drops linearly from its central value to zero. We present a new detailed asteroseismic analysis of GD358, taking advantage of its long sequence of  $\ell = 1$  modes to better constrain the asteroseismic fits and define the limits of high precision white dwarf asteroseismology.

The present analysis allows us to place GD358 in the context of stellar evolution. According to the models, the precursors of DBs emerge from the born-again phase with envelopes containing a nearly uniform mixture of helium (He), carbon (C), and oxygen (O) out to the

photosphere (Dreizler & Heber 1998; Herwig et al. 1999). As the PG 1159 stars cool, the helium diffuses upward and gradually accumulates to form a chemically pure surface layer. Through the DBV instability strip, this process is still ongoing so that instead of a pure helium layer surrounding the carbon and oxygen core, one has a region where the carbon and helium are still mixed. This leads to a double layered structure, with the pure He surface layer overlying the remainder of the uniform He/C/O envelope, all above the degenerate C/O core. A key prediction of the diffusion models is that, for a given stellar mass, the pure He surface layer will steadily grow thicker as the DB star cools. GD358 is the fourth DBV we can use to check this theory. The other three DBVs (Bischoff-Kim et al. (2014); Sullivan et al. (2008); Metcalfe et al. (2003)) have painted a picture qualitatively consistent with the diffusion calculations, with the hotter best fit models having thinner pure helium layers. With GD358, we seek to confirm this trend.

In Section 2, we present our new observations and outline the data reduction process. In Section 3, we establish the framework for frequency identification, and present the list of frequencies used for the asteroseismic investigation. We perform further analysis of the observed frequencies in Section 4, and motivate the  $\ell$  and  $m$  identification of the modes. We present the asteroseismic fitting of GD358’s pulsation spectrum in section 5, present the results in section 6 and discuss our results in section 7.

## 2. New Observations and Data Reduction

GD358 was discovered in 1982 (Winget et al. 1982) and has been the target of the Whole Earth Telescope (WET) in 1990, 1994, 2000, and 2006 (Provencal et al. 2009; Kepler et al. 2003; Winget et al. 1994). New observations presented here include 254 individual observing runs (1105.1 hrs) spanning 2007-2015 (Table 1). Each season of observations was obtained as part of multisite WET campaigns.

Standard WET procedure calls for observers to transfer raw images and calibration files to WET headquarters for analysis at the end of each local night. CCD data reduction follows the steps described in Provencal et al. (2012). In brief, the new observations were obtained with different CCD photometers, each with distinct effective bandpasses. We minimize the bandpass issues by using CCDs with similar detectors whenever possible and employing a red cutoff filter (BG40 or S8612) to normalize wavelength response and reduce extinction effects. We corrected each image for bias and thermal noise, and normalized by the flat field. Aperture photometry using the Maestro photometry pipeline described by Dalessio et al. (2010) was performed on each image, utilizing a range of aperture sizes for the target and comparison stars. For each individual nightly run, we chose the combination of aperture size

and comparison star(s) producing the highest signal/noise light curve.

We used the WQED pipeline (Thompson & Mullally 2009) to examine each nightly light curve for photometric quality, remove outlying points, divide by suitable comparison stars, and correct for differential extinction. Our observational technique is therefore not sensitive to oscillations longer than a few hours. The result is a series of light curves with amplitude variations represented as fractional intensity (mmi). The unit is a linear representation of the fractional intensity of modulation ( $1 \text{ mmi} \approx 1 \text{ mmag}$ ). We present our Fourier transforms (FTs) in units of modulation amplitude ( $1 \text{ mma} = 1 \times 10^{-3} \text{ ma} = 0.1\% = 1 \text{ ppt}$ ).

The final reduction step is to combine the individual light curves (one example shown in Fig. 1) to produce complete light curves for GD358 for each observing season. We assume GD358 oscillates about a mean light level. This assumption allows us to carefully assess overlapping segments from different telescopes and identify and correct any vertical offsets. As discussed in detail in Provencal et al. (2009), we find no significant differences between the noise levels of amplitude spectra using: 1) the combination of all light curves including overlapping segments from different telescopes, 2) the combination of light curves where we retain only higher signal to noise observations in overlapping segments and 3) combining all light curves using data weighted by telescope aperture.

The result is a series of new light curves for each observing season between 2007 and 2015. 2007 contains 8.1 hrs of observation, 2008 26.1 hrs, 2009 45 hrs, 2010 201.3 hrs, 2011 401.1 hrs, 2012 150.5 hrs, 2013 87 hrs, 2014 184.6 hrs, and 2015 55 hrs. Our coverage is not complete, and this incompleteness produces spectral leakage in the amplitude spectra. To quantify this, we sampled single sinusoids using exactly the same times as the original data for each season. The resulting amplitude spectrum, or “spectral window”, is the pattern produced in the FT by a single frequency. The FTs for 2010, 2011, and 2014 are given in Fig. 2.

### 3. Frequency Identification

Our goal is to compile a complete list of GD358’s observed independent and combination frequencies to be used in a comprehensive asteroseismic analysis. GD358 is known for large scale changes in amplitudes and small but not insignificant frequency variations on a variety of timescales (Provencal et al. 2009). The amplitude and frequency variations evident in Fig. 2 will produce artifacts in FTs, greatly complicating any frequency analysis. We reanalyzed all available archival data (Provencal et al. 2009; Kepler et al. 2003; Winget et al. 1994, 1982) as well as our new observations to insure continuity in methodology. We use *Pe-*

*riod04* (Lenz & Breger 2005) for Fourier analysis and nonlinear least squares fitting to select statistically significant peaks in each FT. As discussed in Provencal et al. (2012), we adopt the criterion that a peak must have an amplitude at least four times greater than the average noise level in the given frequency range. We define “noise” as the frequency-dependent average amplitude after prewhitening by the dominant frequencies. This is a conservative estimate, as it is impossible to ensure that all of the “real” frequencies are removed when determining the noise level. This is certainly true for GD358, where the peaks above  $\approx 2500$   $\mu\text{Hz}$  are combination frequencies (Provencal et al. 2009). Fig. 3 displays the average noise as a function of frequency for the 2007-2015 observing seasons. A similar plot for the archival data is presented in Fig. 3 of Provencal et al. (2009). To confirm our uncertainty estimates, we calculated Monte Carlo simulations using the routine provided in *Period04*. This routine generates a set of light curves using the original times, the fitted frequencies and amplitudes, and added Gaussian noise. A least squares fit is performed on each simulated light curve, with the distribution of fit parameters giving the uncertainties. Our Monte Carlo results are consistent with our average noise estimates.

Our frequency selection procedure involves identifying the largest amplitude resolved peak in the FT, fitting a sinusoid at that frequency to the data set, subtracting the fit from the light curve, recomputing the FT, examining the residuals, and repeating the process until no significant power remains. This technique, known as prewhitening, must be employed with an over-abundance of caution, especially since we are aware of amplitude and/or frequency modulation in our data set. A detailed discussion of the prewhitening procedure and steps taken to minimize the effects of amplitude modulation is given in (Provencal et al. 2009). Our final identifications of independent frequencies detected in each observing season are given in Table 2 and Table 3.

## 4. Frequency Analysis

### 4.1. Frequencies Distribution

Perusal of Tables 2 and 3 shows that GD358’s observed frequencies vary in two important ways. First, frequencies detected in a given observing season are not found in all observing seasons. Asteroseismology is based on the assumption that the available pulsation frequencies are linked to stellar structure. Since we are fairly certain that GD358’s internal structure does not change on the timescales of the observations, we can assume that GD358 excites different subsets of its available pulsations at different times. While this is a common phenomenon seen in white dwarf pulsators, the selection mechanism remains unknown. The best way to determine GD358’s complete set of pulsation frequencies is to combine frequency

identifications from multiple observing seasons as outlined in Kleinman et al. (1998).

Fig. 4 presents a schematic representation of GD358’s pulsation frequencies for all the available data. The features of asteroseismic importance are the localized bands between 900 and 2400  $\mu\text{Hz}$  (1100 and 400 s). We interpret each band to represent a single radial overtone. The bands themselves illustrate the second way in which GD358’s frequencies vary. Each band consists of frequency detections from multiple observing seasons, but each band is significantly wider in frequency space than the frequency error of any single measurement, arguing that some process is acting on the frequencies. The simplest explanation is rotational splitting. However, with the exception of the two highest frequency (shortest period) bands ( $k = 8$  and  $k = 9$  from Provencal et al. (2009)), we find no clear evidence of multiplet structure in the frequency distributions of the lower frequency (longer period) bands. The lower frequency band widths are most likely the result of amplitude and/or frequency modulation that obscures any underlying signatures of rotational splitting.

Fig. 4 also shows that the widths of the bands is not constant, but changes from band to band. Bell et al. (2015) presents an interesting analysis of the hydrogen atmosphere pulsator (DAV) KIC4552982, in which they identify 17 bands of pulsation frequencies. This DAV follows the general pattern exhibited by GD358: the highest frequency (shortest period) mode shows evidence of rotational splitting, while the lower frequency (longer period) modes are complex bands. The authors note that this DAV’s bands have different widths in frequency space, and that there may be astronomical significance to this. Although the observational timebases and coverage are quite different (20 continuous months for KIC4552982 vs 30 incomplete years for GD358), our data presents the opportunity to investigate this for a DB pulsator. We measured the width of each of GD358’s band. We define ”width” as the difference between the lowest and highest frequency detected in each band. Fig. 6 presents the results. The widths of GD358’s bands are commensurate with those found in KIC4552982. We also find a general increase in width with decreasing frequency (increasing period), until we reach the mode at 1238  $\mu\text{Hz}$  (807 s). This band has a width at least twice as wide as any other. What is different about this mode???

## 4.2. Mode Identification

Our current work has produced a well defined sequence of modes, adding to previous studies (Montgomery et al. 2010; Provencal et al. 2009; Metcalfe et al. 2000; Winget et al. 1994). The previous identification of these modes as a series of  $l=1$  radial overtones is based mostly on the pulsation frequency distribution and limited spectroscopic analysis (Kotak et al. 2002; Castanheira et al. 2005). It is important to further investigate these

identifications as we initiate an indepth asteroseismic investigation.

GD358’s combination frequencies provide a tool by which we can bolster  $\ell$  identifications. Combination frequencies are typically observed in the FTs of moderate to large amplitude pulsators. They are identified by their exact numerical relationships with parent frequencies. The combinations themselves are not independent, but result from nonlinear effects associated with the surface convection zone (Brickhill 1992; Brassard et al. 1995; Wu 2001; Ising & Koester 2001). Wu (2001) lays the groundwork, showing that observed amplitudes of the combination frequencies depend on geometric factors such as the  $(\ell, m)$  indices of the parents and the inclination of the pulsation axis to the line of sight.

The methods outlined in Provencal et al. (2012) and Montgomery et al. (2010) work best when applied to larger amplitude frequencies detected in high signal to noise data sets such as provided by extensive WET runs. The primary reason for this is that combination frequencies are lower amplitude than their parents, and so are more difficult to detect in sparse data. We chose the 1990, 1994, 2006, 2010, 2011, and 2014 observing seasons, and looked at pulsation frequencies with amplitudes above 10 mma.

Fig. 7 shows the resulting probability distribution of  $\ell$  and  $m$  values for the 1735.96  $\mu\text{Hz}$  frequency as detected in 2014. The mode is clearly identified as an  $\ell = 1$ ,  $m = 1$  mode.

Add table with all l,m ids

Finally, given the  $l=1$  identification, but the lack of definite multiplet structure, the best way to determine the  $l=0$  component for each band is simply to average the frequencies in the bands. We experimented with numerous weighting techniques, and determined there is no significant difference in our solutions. Our final frequency identification for each band is given in Table 4. We use these periods in our asteroseismic fitting.

## 5. Asteroseismic fitting

The basic method in our asteroseismic fitting consists of calculating grids of white dwarf models and running a fitting subroutine to match the periods of the models ( $P_{\text{calc}}$ ) with the observed periods and calculated residuals using the usual formula

$$\sigma_{\text{RMS}} = \sqrt{\frac{\sum_1^{n_{\text{obs}}} (P_{\text{calc}} - P_{\text{obs}})^2}{n_{\text{obs}}}}, \quad (1)$$

where  $n_{\text{obs}}$  is the number of periods present in the pulsation spectrum.

### 5.1. The Models

To compute our models, we used the White Dwarf Evolution Code (WDEC). The WDEC evolves hot polytrope models from temperatures above 100,000 K down to the temperature of our choice. Models in the temperature range of interest for the present study are thermally relaxed solutions to the stellar structure equations. Each model we compute for our grids is the result of such an evolutionary sequence. The WDEC is described in detail in Lamb & van Horn (1975) and Wood (1990). We used smoother core composition profiles and implemented more complex profiles that result from stellar evolution calculations (Salaris et al. 1997). We updated the envelope equation of state tables from those calculated by Fontaine et al. (1977) to those given by Saumon et al. (1995). We use OPAL opacities (Iglesias & Rogers 1996) and plasmon neutrino rates published by Itoh et al. (1996).

DBVs are younger than their cooler cousins the DAVs. Time-dependent diffusion calculations (e.g. Dehner & Kawaler 1995; Althaus et al. 2005) show that at 24,000 K, a typical temperature for a DBV, the carbon has not fully settled into the core of the star yet. We expect the helium layer to be separated into a mixed He/C layer with a pure He layer on top, as shown in Fig. 8. Following Metcalfe et al. (2005), we adopted and parameterized this structure in our models.

### 5.2. Initial Grid Search

In our asteroseismic fits, we initially varied six parameters; the effective temperature, the mass and four structure parameters. There are two parameters associated with the shapes of the oxygen (and carbon) core composition profiles: the central oxygen abundance ( $X_o$ ) and the edge of the homogeneous carbon and oxygen core ( $q_{\text{fm}}$ , as a fraction of stellar mass. For envelope structure,  $M_{\text{env}}$  marks the location of the base of the helium layer and  $M_{\text{He}}$  the location where the helium abundance rises to 1 (see Fig. 8).  $M_{\text{env}}$  and  $M_{\text{He}}$  are mass coordinates, defined as e.g.  $M_{\text{env}} = -\log(1 - M(r)/M_*)$ , where  $M(r)$  is the mass enclosed in radius  $r$  and  $M_*$  is the stellar mass.

We started with a master grid (table 5) chosen so that it covered all relevant area of parameter space and had sufficient resolution to find any region of local minimum in the fitness parameter (Bischoff-Kim & Metcalfe 2011; Bischoff-Kim et al. 2014). We used the minimum resolution that was computationally manageable. The master grid involved the computation of 10,483,200 models. We fit simultaneously all 15 periods, requiring all of them to be  $\ell = 1$  modes. A fitness map of this initial fit are shown in the left panel of Fig. 9, and the parameters of the best fit model are listed in table 5.



### 5.3. Asymptotic Period Spacing

Before we refine the period-by-period fitting optimization, it is worth stepping back and consider what we can learn from the average period spacing of GD358. The average period spacing provides an asteroseismic measure of the mass and temperature of the star, independent of the details of internal chemical composition profiles. Higher  $k$  modes are not strongly trapped in the core and according to asymptotic theory, they should be nearly evenly spaced in period. This spacing is given by (Unno et al. 1989)

$$\Delta P = \frac{\pi}{\ell(\ell+1)^{1/2}} \left[ \int_{r_1}^{r_2} \frac{N}{r} dr \right]^{-1}, \quad (2)$$

where  $r_1$  and  $r_2$  are turning points of the mode and  $N$  is the Brunt-Väisälä frequency. The asymptotic period spacing is  $\ell$  dependent, with higher  $\ell$  modes having smaller spacing. In the case of GD 358, we have a single  $\ell = 1$  sequence so we only need to worry about the dependence of the asymptotic period spacing on the Brunt-Väisälä frequency. Much if not all of GD 358’s pulsation spectrum is close to the asymptotic limit, because the shortest period observed is a  $k = 8$  mode.

The dependence of  $\Delta P$  on the Brunt-Väisälä frequency leads to higher mass and lower temperature models having a smaller period spacing (their interior is less compressible). This effect appears in asteroseismic fitting of white dwarfs and also sdB stars as a ubiquitous diagonal trend in contour maps of the quality of the fits in the mass-effective temperature plane (e.g Bischoff-Kim et al. 2014; Castanheira & Kepler 2009; Charpinet et al. 2008). One requirement for the periods of the model to match the observed periods is that the average period spacing in the models matches the average period spacing in the observed pulsation spectrum. If a good match occurs for a given mass and effective temperature, then models with lower mass but higher effective temperature will also match well.

We can use the sequence of 13 consecutive  $\ell = 1$  modes found in GD358’s pulsation spectrum to calibrate our models. Using the  $\ell = 1$  sequence of modes, we compute an average period spacing of 39.08 seconds. We call this  $\Delta P_{\text{obs}}$ . For each model in the master grid, we compute an asymptotic period spacing ( $\Delta P_{\text{calc}}$ ). This asymptotic period spacing is calculated by first discarding the 10 lowest  $k$  modes. The exact value of 10 is somewhat arbitrary, but it is chosen so that the modes we use in our computation are indeed in the asymptotic limit. The higher  $k$  modes show weaker trapping than the lower  $k$  modes. We then fit a line through the set  $(k_i, P_i)$ . The slope gives us the asymptotic period spacing in the model. We also calculate the residuals of the fits and discard the models that have residuals above a certain limit. The limit is chosen by checking the procedure by eye on a

few models.

We show a contour map of the location of the models that best match the average period spacing of 39.08 seconds in Fig. 9, right panel. We place it side by side with a contour map showing the location of the best fit model in the same region of parameter space, based on the master grid fitting described in section 5.2. Note how the best fit model falls right within the valley where the period spacings between GD358 and the models match. This should come as no surprise, as in order for 15 periods to fit reasonably well, the model period list should have a spacing similar to that of the observed period spectrum. It is also somewhat comforting to find that the spectroscopic data point also falls within the swath where the period spacings match (though the best fit model based on the master grid search is further down the valley).

One can fit simultaneously the average period spacing and the individual periods formally while performing the fits by using some prescription to calculate the goodness of fit. This leads to a more complex relation than defined in equation 1. Note that the period spacing is a much weaker constraint than the individual period fit. If one takes 5.0 seconds as an upper limit for goodness of fit, that includes 9 points in the period-by-period fit plot (left panel in Fig 9), but the entire parameter plane for the average period spacing fit plot (right panel). We bypass the difficulty of assigning proper weights to each constraint by taking the average period spacing fit into account as an initial step to refine our fits. We refine our fits by calculating models on a higher resolution grid that fall within the region defined by the two diagonal lines shown in the right panel of Fig. 9 (covering the effective temperature range 25,000 K-27,500 K), then performing a period-by-period fit using equation 1.

#### 5.4. Optimal Grid Resolution

Having determined a more restricted region of parameter space to search for the best fit models, we now turn to the question of how fine we need to make our refined grid. We want to have a high enough resolution grid that we can be sure we captured a true minimum, but on the other hand, there are computational limitations to how many models we can afford to calculate, save, and process.

One way to gain a sense of how fine the grid needs to be is to make single parameter cuts through parameter space. Fig. 10 shows such a cut for master grid models. The plot was made by fixing 5 of the 6 parameters to the best fit values of the best initial fit (see table 5). We allowed the effective temperature to vary through the entire range considered for the master grid. The fits seem to settle to a minimum in a smooth way, except for the

striking anomaly at 26,600 K. Upon further investigation, we discovered that anomaly to be due to a single period in the pulsation spectrum: the 968 s mode. In the models, that mode appears highly sensitive to some parameters, and in a non-continuous way.

This is unsettling, as it is easy to see that one might miss the best fit models if the grid is not fine enough. In order to quantify "fine enough", we ran systematic scans, computing models with 5 parameter fixed and allowing the 6th parameter to vary in very fine steps. We went down in step sizes to  $\Delta T_{\text{eff}} = 1$  K,  $\Delta M_* = 0.0001 M_{\odot}$ ,  $\Delta M_{\text{env}} = \Delta M_{\text{He}} = 0.01$  dex,  $\Delta X_o = 0.01$ , and  $\Delta q_{\text{fm}} = 0.001$ . Results for 4 different parameters are shown in Fig 11. The figure includes zooms to regions of interest. Other modes also exhibit a behavior similar to the 968 s mode, in different regions of parameter space. For instance, the jumps in  $\sigma_{\text{RMS}}$  for  $q_{\text{fm}}$  between 0.3 and 0.4 are due to sudden changes in the 854.5 and 902.5 s modes.

Regardless of the behavior of individual modes in the models as the parameters vary, we can assert from the test above what step sizes will allow us to minimize our risk of missing a minimum. We settled on optimal step sizes of  $\Delta T_{\text{eff}} = 20$ ,  $\Delta M_* = 0.0002$ ,  $\Delta M_{\text{He}} = 0.2$ ,  $\Delta X_o = 0.2$ , and  $\Delta q_{\text{fm}} = 0.01$  for our refined grid. To offset the computational cost, we fixed  $M_{\text{env}}$  to a canonical value of 2.80 (Dehner & Kawaler 1995). It is clear from the second panel in Fig. 11 that there is only a weak dependence of the fits on that parameter.

## 6. Results of the Period Fitting

The parameters for our best fit model based on the refined grid are listed in table 5 and the periods of that model in table 4. The goodness of fit of the model is  $\sigma_{\text{RMS}} = 4.22$  s. In Fig. 8, we show the interior structure of the best fit model, with the corresponding Brunt-Väisälä frequency profile.

### 6.1. Validation of the Fitting Method

With what we have discovered in section 5.4, it is only natural to be concerned about whether we have truly found a best fit. We performed a simple test to validate our fitting method, which consisted of using the exact same procedure to find a best fit to the periods of a model that was not on any of the grids we calculated, but that did have parameters that were very close to the best fit model. We used the period list for a model with parameters  $T_{\text{eff}} = 25570$  K,  $M_*/M_{\odot} = 0.5437$ ,  $M_{\text{He}} = -5.30$ ,  $X_o = 0.42$ , and  $q_{\text{fm}} = 0.175$ , fixing  $M_{\text{env}}$  to -2.80, and including only the subset of 15  $\ell = 1$  periods that match GD358's pulsation spectrum.

We first performed a fitting of the periods using the master (coarse) grid described in section 5.2. This placed the best fit model in the appropriate region of parameter space. Then we refined our fits, using the grids described at the end of section 5.3. We are able to recover the best fit parameters adequately, with the top 10 best fit models having parameters  $T_{\text{eff}} = 25620 \pm 67$  K,  $M_*/M_{\odot} = 0.5431 \pm 0.0011$ ,  $M_{\text{He}} = -5.32 \pm 0.098$ ,  $X_o = 0.40$  for all top 10 fits, and  $q_{\text{fm}} = 0.175 \pm 0.5$ . The top 10 best fits have  $\sigma_{\text{RMS}} \sim 0.5$  s. We remind the reader that the step sizes in the second phase of the fitting are 20 K for  $T_{\text{eff}}$ , 0.2 for  $M_{\text{He}}$ , 0.2 for  $X_o$ , and 0.01 for  $q_{\text{fm}}$ .

## 7. Discussion and Conclusions

We find a best fit effective temperature of 25,560 K for GD358, hotter than the recent spectroscopic determination of 24000 K (Koester 2013) by more than one sigma (the mass matches the spectroscopy). For reasons discussed in section 5.3, while it is possible to determine whether a star is closer to the red edge or the blue edge of the instability strip, effective temperature is poorly constrained in asteroseismic fitting. As we can see from Fig. 10, there are local minima that are almost as good as the global minimum we found, some at lower effective temperature. Structure parameters, especially the location of chemical transitions (with the exception of  $M_{\text{env}}$  in the case of GD358) are more tightly constrained.

In fitting GD358, we discovered that the best fit parameter space was semi-chaotic, with two very close sets of parameters able to yield very different fitness number. This is due to the fact that some periods can change values drastically for a small change in a given parameter. The phenomenon is only semi-chaotic, as the period will oscillate between two values. This is good news, because there remains a well defined minimum in parameter space, even if it is a double valued minimum. Any search algorithm that uses a small enough mesh should be able to find the minimum. This phenomenon likely became apparent while fitting GD358 because it is the first white dwarf for which we fit a long consecutive sequence of modes of the same  $\ell$  with high resolution grids.

In this study, we varied the parameters that have traditionally been varied in this type of asteroseismic fitting, because we wanted to be able to place our results side by side with other studies of DBVs, including KIC 8626021 (Bischoff-Kim et al. 2014), EC200058 (Bischoff-Kim & Metcalfe 2011), and CBS114 (Metcalfe et al. 2005). We now have 4 DBVs that were the object of asteroseismic fitting that used a consistent set of models. Unfortunately, GD358 does not fit neatly into the trend (neither does EC20058). It is true, however that the two blue edge objects (EC20058 and KIC 8626021) have thinner helium layers ( $M_{\text{He}} = -6.10$  and  $-7.90$  respectively) than the red edge stars ( $M_{\text{He}} = -5.40$  for GD358 and  $-5.94$  for

CBS114. Pinning down the proper effective temperature of each star is challenging. To be fully consistent, it is best to use the asteroseismic values, even if effective temperature is not strongly constrained asteroseismically. At best we can contrast blue edge versus red edge objects.

Numerical experiments of the type presented in section 5.4 (Bischoff-Kim 2015) have shown that, consistent with the theory of non-radial oscillations, the shape of the transition zones matter as much as where the transition zones occur. Modern asteroseismic fitting vary parameters attached to the shape of transition zones (Giammichele et al. 2015). We refrained from doing this in the present study because we wanted to compare our results with previous studies of DBVs. It is unclear how much of an effect on structure parameters a change in parameterization will have.

*Facilities:* MCAO:0.6m (), Struve(), KPNO:2.1m (), , BOAO:1.8m (), Lulin:1.8m (), Beijing:2.16m (), Maidanek:1.5m (), Peak Terskol

## REFERENCES

- Althaus, L. G., Serenelli, A. M., Panei, J. A., Córscico, A. H., García-Berro, E., & Scóccola, C. G. 2005, *A&A*, 435, 631
- Beauchamp, A., Wesemael, F., Bergeron, P., Fontaine, G., Saffer, R. A., Liebert, J., & Brassard, P. 1999, *ApJ*, 516, 887
- Bell, K. J., Hermes, J. J., Bischoff-Kim, A., Moorhead, S., Montgomery, M. H., Østensen, R., Castanheira, B. G., & Winget, D. E. 2015, *ApJ*, 809, 14
- Bischoff-Kim, A. 2015, in *Astronomical Society of the Pacific Conference Series*, Vol. 493, 19th European Workshop on White Dwarfs, ed. P. Dufour, P. Bergeron, & G. Fontaine, 175
- Bischoff-Kim, A., & Metcalfe, T. S. 2011, *MNRAS*, 414, 404
- Bischoff-Kim, A., Østensen, R. H., Hermes, J. J., & Provencal, J. L. 2014, *ApJ*, 794, 39
- Brassard, P., Fontaine, G., & Wesemael, F. 1995, *ApJS*, 96, 545
- Brickhill, A. J. 1992, *MNRAS*, 259, 529
- Castanheira, B. G., & Kepler, S. O. 2009, *MNRAS*, 396, 1709

- Castanheira, B. G., Nitta, A., Kepler, S. O., Winget, D. E., & Koester, D. 2005, *A&A*, 432, 175
- Charpinet, S., Van Grootel, V., Reese, D., Fontaine, G., Green, E. M., Brassard, P., & Chayer, P. 2008, *A&A*, 489, 377
- Dalessio, J., Provencal, J. L., Sullivan, D., & Shipman, H. L. 2010, in 17th European Workshop on White Dwarfs, ed. K. Werner & T. Rauch, AIP Conference Proceedings
- Dehner, B. T., & Kawaler, S. D. 1995, *ApJ*, 445, L141
- Dreizler, S., & Heber, U. 1998, *A&A*, 334, 618
- Fontaine, G., Graboske, H. C., J., & Van Horn, H. M. 1977, *ApJS*, 35, 293
- Giammichele, n., Charpinet, S., Fontaine, G., Brassard, P., & Zong, W. 2015, IAU General Assembly, 22, 2256942
- Herwig, F., Blöcker, T., Langer, N., & Driebe, T. 1999, *A&A*, 349, L5
- Iben, Jr., I., Kaler, J. B., Truran, J. W., & Renzini, A. 1983, *ApJ*, 264, 605
- Iglesias, C. A., & Rogers, F. J. 1996, *ApJ*, 464, 943
- Ising, J., & Koester, D. 2001, *A&A*, 374, 116
- Itoh, N., Nishikawa, A., & Kohyama, Y. 1996, *ApJ*, 470, 1015
- Kepler, S. O., et al. 2003, *A&A*, 401, 639
- Kleinman, S. J., et al. 1998, *ApJ*, 495, 424
- Koester, D. 2013, private communication
- Kotak, R., van Kerkwijk, M. H., & Clemens, J. C. 2002, *A&A*, 388, 219
- Lamb, D. Q., & van Horn, H. M. 1975, *ApJ*, 200, 306
- Lawlor, T. M., & MacDonald, J. 2006, *MNRAS*, 371, 263
- Lenz, P., & Breger, M. 2005, *Communications in Asteroseismology*, 146, 53
- Metcalfe, T. S., Montgomery, M. H., & Kanaan, A. 2003, American Astronomical Society Meeting, 203

- Metcalfe, T. S., Montgomery, M. H., & Kanaan, A. 2005, in *Astronomical Society of the Pacific Conference Series*, Vol. 334, 14th European Workshop on White Dwarfs, ed. D. Koester & S. Moehler, 465
- Metcalfe, T. S., Nather, R. E., & Winget, D. E. 2000, *ApJ*, 545, 974
- Montgomery, M. H., Provencal, J. L., Kanaan, A., Thompson, S., Dalessio, J., Shipman, H., & Winget, D. E. 2010, *ApJ*, 716, 84
- Provencal, J. L., et al. 2009, *ApJ*, 693, 564
- . 2012, *ApJ*, 751, 91
- Salaris, M., Dominguez, I., Garcia-Berro, E., Hernanz, M., Isern, J., & Mochkovitch, R. 1997, *ApJ*, 486, 413
- Saumon, D., Chabrier, G., & van Horn, H. M. 1995, *ApJS*, 99, 713
- Sullivan, D. J., et al. 2008, *MNRAS*, 387, 137
- Thompson, S. E., & Mullally, F. 2009, *Journal of Physics Conference Series*, 172, 012081
- Winget, D. E., van Horn, H. M., Tassoul, M., Fontaine, G., Hansen, C. J., & Carroll, B. W. 1982, *ApJ*, 252, L65
- Winget, D. E., et al. 1994, *ApJ*, 430, 839
- Wood, M. A. 1990, PhD thesis, The University of Texas at Austin
- Wu, Y. 2001, *MNRAS*, 323, 248




Fig. 1.— Light curve of GD358 obtained with the Peak Terskol 2.0m. Each point corresponds to a 10 s exposure. The nonlinear, multiperiodic nature of this star is clearly evident.






Fig. 2.— Fourier Transforms of GD358 for the 2010, 2011, and 2014 observing seasons. The frequency range of the observed series of  $\ell = 1$  modes discussed in section 4 is indicated by the left arrow (red). Peaks below  $\approx 2400 \mu\text{Hz}$  are combination frequencies discussed in section 4 (right arrow, blue). (A color version of this figure is available in the online journal.)




Fig. 3.— A comparison of the average noise as a function of frequency for the 2007-2015 observing seasons. Each data set was prewhitened by its dominant frequencies. The noise levels for each season are somewhat different. This must be taken into account during frequency analysis.




Fig. 4.— A schematic representation of GD358’s pulsation modes for all available data between 1982 and 2015. Systematic patterns of distribution are evident. The bands between 2400 and 1000  $\mu\text{Hz}$  (400 and 1000 s) are of particular importance for this work. (A color version of this figure is available in the online journal.)




Fig. 5.— The detailed schematic distribution of frequencies for the bands near 1300(*right* and 2360  $\mu\text{Hz}$  (*left*). The 1300  $\mu\text{Hz}$  band shows no multiplet structure, unlike the 2360  $\mu\text{Hz}$  band. Here we see clear evidence of multiplet structure. Assuming the triplet structure represents  $\ell = 1$ , this implies a rotation period of  $\approx 1.5$  days. Note changes in the scale for both axes. (A color version of this figure is available in the online journal.)




Fig. 6.— Width of each bank of power in Fig. 4. We find an overall increase in band thickness with decreasing frequency (increasing period). The band at 807 s has a width at least twice as large as other bands.




Fig. 7.— Probability distribution (from 0 to 1) of  $\ell$  and  $m$  values for the 1735.96 variation detected in 2014. The solutions are from individual simulations with  $Res_{rms} < 9.5^{-06}$ . The amplitudes of the observed combination frequencies argue that this is  $\ell = 1$ ,  $m=1$ .




Fig. 8.— *Upper panel:* Chemical composition profiles of the best fit model. The center of the model is to the left and the surface to the right.  $q = 2$  corresponds to  $M_r = 0.99 M_*$ . The vertical axis shows fractional abundances. *Lower panel:* The corresponding Brunt-Väisälä frequency ( $\log N^2$  on the vertical axis).




Fig. 9.— *Left panel:* Contour map showing the location of the best fit models. The quantity plotted in the mass-effective temperature plane is the fitness parameter defined in equation 1. The scale is in seconds. *Right panel:* Contour map of the difference between the average periods spacing determined from GD 358’s periods and the asymptotic period spacing of the models, in the mass-effective temperature plane. The scale is in 10s of seconds, so the worse matches pictured on the plot have  $|\Delta P_{\text{obs}} - \Delta P_{\text{calc}}| \sim 2$  s.




Fig. 10.— Dependence of the fitness parameter on effective temperature alone based on the master grid of models.




Fig. 11.— Dependence of the fitness parameter on different parameters. The 5 parameters other than the one shown on the horizontal axis in each plot are held to fixed values corresponding to the best fit model (table 8).



Table 1.

| Run Name      | Telescope    | Detector | Date       | Length<br>(hrs) |
|---------------|--------------|----------|------------|-----------------|
| mcao070524-01 | MCAO 0.6     | CCD      | 2007-05-24 | 4.0             |
| mcao070531-02 | MCAO 0.6     | CCD      | 2007-05-31 | 4.1             |
| mcao080613-01 | MCAO 0.6     | CCD      | 2008-06-13 | 2.0             |
| mcao080617-02 | MCAO 0.6     | CCD      | 2008-07-25 | 1.2             |
| mcao080726-01 | MCAO 0.6     | CCD      | 2008-07-26 | 1.7             |
| mcao080730-01 | MCAO 0.6     | CCD      | 2008-07-30 | 3.5             |
| mcao080802-01 | MCAO 0.6     | CCD      | 2008-08-02 | 3.7             |
| pjmo080706-03 | PJMO 0.6     | CCD      | 2008-07-06 | 5.1             |
| suho080810-19 | Suhora 0.6   | CCD      | 2008-08-10 | 4.7             |
| suho080811-19 | Suhora 0.6   | CCD      | 2008-08-11 | 2.5             |
| boao090528-17 | BOAO 1.8     | CCD      | 2009-05-28 | 2.0             |
| kore090529-16 | BOAO 1.8     | CCD      | 2009-05-29 | 2.4             |
| mcao090513-01 | MCAO 0.6     | CCD      | 2009-05-13 | 2.4             |
| mcao090519-03 | MCAO 0.6     | CCD      | 2009-05-19 | 3.2             |
| mcao090520-01 | MCAO 0.6     | CCD      | 2009-05-20 | 4.8             |
| mcao090521-01 | MCAO 0.6     | CCD      | 2009-05-21 | 4.9             |
| mcao090522-01 | MCAO 0.6     | CCD      | 2009-05-22 | 3.1             |
| mcao090531-02 | MCAO 0.6     | CCD      | 2009-05-31 | 2.1             |
| mcao090601-01 | MCAO 0.6     | CCD      | 2009-06-01 | 3.3             |
| mole090526-20 | Moletai 1.65 | CCD      | 2009-05-26 | 3.7             |
| suho090520-19 | Suhora 0.6   | CCD      | 2009-05-20 | 2.2             |
| suho090521-19 | Suhora 0.6   | CCD      | 2009-05-21 | 5.1             |
| vien090525-19 | Vienna 0.6   | CCD      | 2009-05-25 | 6.2             |
| krak100523-20 | Krakow 0.4   | CCD      | 2010-05-23 | 3.4             |
| krak100526-20 | Krakow 0.4   | CCD      | 2010-05-26 | 2.3             |
| krak100616-20 | Krakow 0.4   | CCD      | 2010-06-16 | 4.3             |
| krak100617-20 | Krakow 0.4   | CCD      | 2010-06-17 | 4.5             |
| mcao100516-01 | MCAO 0.6     | CCD      | 2010-05-16 | 2.0             |
| mcao100521-01 | MCAO 0.6     | CCD      | 2010-05-21 | 0.9             |
| mcao100521-05 | MCAO 0.6     | CCD      | 2010-05-21 | 0.8             |
| mcao100526-01 | MCAO 0.6     | CCD      | 2010-05-26 | 3.3             |
| mcd0100517-06 | McDonald 2.1 | CCD      | 2010-05-17 | 4.8             |

Table 1—Continued

| Run Name      | Telescope     | Detector | Date       | Length<br>(hrs) |
|---------------|---------------|----------|------------|-----------------|
| mole100516-22 | Moletai 1.65  | CCD      | 2010-05-26 | 1.8             |
| mole100517-22 | Moletai 1.65  | CCD      | 2010-05-17 | 2.0             |
| mole100520-21 | Moletai 1.65  | CCD      | 2010-05-20 | 3.3             |
| mole100521-21 | Moletai 1.65  | CCD      | 2010-05-21 | 3.5             |
| pjmo100519-04 | PJMO 0.6      | CCD      | 2010-05-19 | 4.0             |
| pjmo100520-02 | PJMO 0.6      | CCD      | 2010-05-20 | 3.8             |
| pjmo100521-05 | PJMO 0.6      | CCD      | 2010-05-21 | 3.3             |
| pjmo100522-02 | PJMO 0.6      | CCD      | 2010-05-22 | 7.0             |
| pjmo100523-04 | PJMO 0.6      | CCD      | 2010-05-23 | 6.0             |
| pjmo100524-02 | PJMO 0.6      | CCD      | 2010-05-24 | 7.5             |
| pjmo100527-02 | PJMO 0.6      | CCD      | 2010-05-27 | 8.3             |
| pjmo100528-02 | PJMO 0.6      | CCD      | 2010-05-28 | 8.3             |
| pjmo100619-03 | PJMO 0.6      | CCD      | 2010-06-19 | 4.0             |
| pjmo100622-02 | PJMO 0.6      | CCD      | 2010-06-22 | 0.7             |
| suho100617-21 | Suhora 0.6    | CCD      | 2010-06-17 | 4.3             |
| terb100516-17 | Terskol 2.0   | CCD      | 2010-05-16 | 2.4             |
| terb100520-17 | Terskol 2.0   | CCD      | 2010-05-20 | 8.9             |
| terb100521-20 | Terskol 2.0   | CCD      | 2010-05-21 | 4.0             |
| terb100524-19 | Terskol 2.0   | CCD      | 2010-05-24 | 5.6             |
| terb100525-17 | Terskol 2.0   | CCD      | 2010-05-25 | 7.3             |
| tueb100522-20 | Tuebingen 0.8 | SBig     | 2010-05-22 | 5.9             |
| tueb100523-20 | Tuebingen 0.8 | SBig     | 2010-05-23 | 5.9             |
| tueb100524-20 | Tuebingen 0.8 | SBig     | 2010-05-24 | 6.0             |
| tueb100604-20 | Tuebingen 0.8 | SBig     | 2010-06-04 | 5.8             |
| tueb100605-20 | Tuebingen 0.8 | SBig     | 2010-06-05 | 5.3             |
| turk100512-18 | Canakkale 1.2 | CCD      | 2010-05-12 | 6.6             |
| turk100513-22 | Canakkale 1.2 | CCD      | 2010-05-13 | 2.8             |
| turk100516-19 | Canakkale 1.2 | CCD      | 2010-05-16 | 6.4             |
| turk100520-19 | Canakkale 1.2 | CCD      | 2010-05-20 | 1.1             |
| turk100522-21 | Canakkale 1.2 | CCD      | 2010-05-22 | 2.6             |
| turk100523-19 | Canakkale 1.2 | CCD      | 2010-05-23 | 6.0             |
| turk100611-19 | Canakkale 1.2 | CCD      | 2010-06-11 | 5.2             |

Table 1—Continued

| Run Name      | Telescope     | Detector | Date       | Length<br>(hrs) |
|---------------|---------------|----------|------------|-----------------|
| turk100613-19 | Canakkale 1.2 | CCD      | 2010-06-13 | 5.3             |
| turk100615-20 | Canakkale 1.2 | CCD      | 2010-06-15 | 4.8             |
| turk100618-19 | Canakkale 1.2 | CCD      | 2010-06-18 | 4.5             |
| turk100620-20 | Canakkale 1.2 | CCD      | 2010-06-20 | 4.8             |
| hvar110521-22 | Hvar 1.0      | CCD      | 2011-05-21 | 2.6             |
| hvar110526-19 | Hvar 1.0      | CCD      | 2011-05-26 | 6.6             |
| hvar110527-19 | Hvar 1.0      | CCD      | 2011-05-27 | 3.5             |
| hvar110529-19 | Hvar 1.0      | CCD      | 2011-05-29 | 6.7             |
| hvar110531-19 | Hvar 1.0      | CCD      | 2011-05-31 | 4.3             |
| hvar110602-19 | Hvar 1.0      | CCD      | 2011-06-02 | 3.8             |
| hvar110604-19 | Hvar 1.0      | CCD      | 2011-06-04 | 6.6             |
| hvar110606-20 | Hvar 1.0      | CCD      | 2011-06-06 | 4.4             |
| krak110506-20 | Krakow 0.4    | CCD      | 2011-05-06 | 5.4             |
| krak110509-19 | Krakow 0.4    | CCD      | 2011-05-09 | 6.7             |
| krak110510-19 | Krakow 0.4    | CCD      | 2011-05-10 | 6.2             |
| krak110511-19 | Krakow 0.4    | CCD      | 2011-05-11 | 6.7             |
| krak110512-19 | Krakow 0.4    | CCD      | 2011-05-12 | 3.1             |
| krak110516-19 | Krakow 0.4    | CCD      | 2011-05-16 | 6.1             |
| krak110517-19 | Krakow 0.4    | CCD      | 2011-05-17 | 5.6             |
| krak110518-19 | Krakow 0.4    | CCD      | 2011-05-18 | 5.7             |
| krak110519-20 | Krakow 0.4    | CCD      | 2011-05-19 | 5.4             |
| krak110520-20 | Krakow 0.4    | CCD      | 2011-05-20 | 5.2             |
| krak110522-21 | Krakow 0.4    | CCD      | 2011-05-22 | 4.3             |
| krak110523-23 | Krakow 0.4    | CCD      | 2011-05-23 | 1.4             |
| krak110524-19 | Krakow 0.4    | CCD      | 2011-05-24 | 4.0             |
| krak110525-20 | Krakow 0.4    | CCD      | 2011-05-25 | 5.4             |
| krak110526-20 | Krakow 0.4    | CCD      | 2011-05-26 | 5.3             |
| krak110529-20 | Krakow 0.4    | CCD      | 2011-05-29 | 5.0             |
| krak110530-20 | Krakow 0.4    | CCD      | 2011-05-30 | 4.8             |
| krak110531-20 | Krakow 0.4    | CCD      | 2011-05-31 | 0.8             |
| krak110604-20 | Krakow 0.4    | CCD      | 2011-06-04 | 5.3             |
| mole110426-22 | Moletai 1.65  | CCD      | 2011-04-26 | 1.7             |

Table 1—Continued

| Run Name      | Telescope      | Detector | Date       | Length<br>(hrs) |
|---------------|----------------|----------|------------|-----------------|
| mole110427-21 | Moletai 1.65   | CCD      | 2011-04-27 | 3.7             |
| mole110430-20 | Moletai 1.65   | CCD      | 2011-04-30 | 0.8             |
| mole110502-20 | Moletai 1.65   | CCD      | 2011-05-02 | 0.4             |
| mole110502-21 | Moletai 1.65   | CCD      | 2011-05-02 | 2.6             |
| mole110505-20 | Moletai 1.65   | CCD      | 2011-05-05 | 1.1             |
| mole110510-21 | Moletai 1.65   | CCD      | 2011-05-10 | 3.5             |
| mtlm110426-05 | Mt. Lemmon 1.0 | CCD      | 2011-04-26 | 6.3             |
| mtlm110427-06 | Mt. Lemmon 1.0 | CCD      | 2011-04-27 | 5.5             |
| mtlm110428-04 | Mt. Lemmon 1.0 | CCD      | 2011-04-28 | 7.1             |
| mtlm110429-05 | Mt. Lemmon 1.0 | CCD      | 2011-04-29 | 6.8             |
| mtlm110430-05 | Mt. Lemmon 1.0 | CCD      | 2011-04-30 | 6.4             |
| mtlm110501-05 | Mt. Lemmon 1.0 | CCD      | 2011-05-01 | 6.6             |
| mtlm110502-05 | Mt. Lemmon 1.0 | CCD      | 2011-05-02 | 6.5             |
| naoc110426-13 | NAOC 0.5       | CCD      | 2011-04-26 | 6.9             |
| naoc110427-11 | NAOC 0.5       | CCD      | 2011-04-27 | 3.3             |
| naos110426-13 | NAOC 0.85      | CCD      | 2011-04-26 | 6.9             |
| naos110427-11 | NAOC 0.85      | CCD      | 2011-04-27 | 3.3             |
| naos110428-12 | NAOC 0.85      | CCD      | 2011-04-28 | 2.6             |
| naos110501-13 | NAOC 0.85      | CCD      | 2011-05-01 | 7.1             |
| naos110502-12 | NAOC 0.85      | CCD      | 2011-05-02 | 8.3             |
| naos110505-13 | NAOC 0.85      | CCD      | 2011-05-05 | 7.0             |
| pjmo110426-07 | PJMO 0.6       | CCD      | 2011-04-26 | 3.0             |
| pjmo110428-04 | PJMO 0.6       | CCD      | 2011-04-28 | 2.0             |
| pjmo110429-03 | PJMO 0.6       | CCD      | 2011-04-29 | 2.5             |
| pjmo110430-04 | PJMO 0.6       | CCD      | 2011-04-30 | 3.0             |
| pjmo110503-04 | PJMO 0.6       | CCD      | 2011-05-03 | 2.6             |
| pjmo110518-03 | PJMO 0.6       | CCD      | 2011-05-18 | 6.7             |
| pjmo110519-03 | PJMO 0.6       | CCD      | 2011-05-19 | 2.7             |
| pjmo110521-03 | PJMO 0.6       | CCD      | 2011-05-21 | 2.7             |
| pjmo110524-02 | PJMO 0.6       | CCD      | 2011-05-24 | 5.2             |
| pjmo110525-07 | PJMO 0.6       | CCD      | 2011-05-25 | 3.2             |
| pjmo110526-03 | PJMO 0.6       | CCD      | 2011-05-26 | 7.0             |

Table 1—Continued

| Run Name      | Telescope     | Detector | Date       | Length<br>(hrs) |
|---------------|---------------|----------|------------|-----------------|
| pjmo110527-02 | PJMO 0.6      | CCD      | 2011-05-27 | 0.8             |
| pjmo110527-03 | PJMO 0.6      | CCD      | 2011-05-27 | 6.7             |
| suho110521-20 | Suhora 0.6    | CCD      | 2011-05-21 | 5.2             |
| suho110522-19 | Suhora 0.6    | CCD      | 2011-05-22 | 5.9             |
| suho110523-20 | Suhora 0.6    | CCD      | 2011-05-23 | 5.6             |
| suho110527-20 | Suhora 0.6    | CCD      | 2011-05-27 | 2.0             |
| suho110529-20 | Suhora 0.6    | CCD      | 2011-05-29 | 4.6             |
| suho110530-20 | Suhora 0.6    | CCD      | 2011-05-30 | 4.9             |
| terb110527-20 | Terskol 2.0   | CCD      | 2011-05-27 | 2.4             |
| terb110529-17 | Terskol 2.0   | CCD      | 2011-05-27 | 4.4             |
| terb110530-18 | Terskol 2.0   | CCD      | 2011-05-30 | 2.6             |
| terb110531-17 | Terskol 2.0   | CCD      | 2011-05-31 | 1.8             |
| terb110601-19 | Terskol 2.0   | CCD      | 2011-06-01 | 3.3             |
| tueb110504-19 | Tuebingen 0.8 | SBig     | 2011-05-04 | 6.8             |
| tueb110505-19 | Tuebingen 0.8 | SBig     | 2011-05-05 | 6.8             |
| tueb110506-19 | Tuebingen 0.8 | SBig     | 2011-05-06 | 6.7             |
| tueb110508-19 | Tuebingen 0.8 | SBig     | 2011-05-08 | 6.6             |
| tueb110509-19 | Tuebingen 0.8 | SBig     | 2011-05-09 | 6.7             |
| tueb110513-19 | Tuebingen 0.8 | SBig     | 2011-05-13 | 2.2             |
| tueb110518-22 | Tuebingen 0.8 | SBig     | 2011-05-18 | 4.1             |
| tueb110523-20 | Tuebingen 0.8 | SBig     | 2011-05-23 | 5.9             |
| tueb110524-20 | Tuebingen 0.8 | SBig     | 2011-05-24 | 5.9             |
| tueb110525-20 | Tuebingen 0.8 | SBig     | 2011-05-24 | 6.0             |
| tueb110529-20 | Tuebingen 0.8 | SBig     | 2011-05-29 | 5.9             |
| tueb110530-20 | Tuebingen 0.8 | SBig     | 2011-05-30 | 3.4             |
| tubil10602-00 | Tubitak 1.0   | CCD      | 2011-06-02 | 1.1             |
| tubil10602-23 | Tubitak 1.0   | CCD      | 2011-06-02 | 2.0             |
| tubil10603-20 | Tubitak 1.0   | CCD      | 2011-06-03 | 5.2             |
| boao120418-18 | BOAO 1.8      | CCD      | 2012-04-18 | 1.7             |
| krak120418-00 | Krakow 0.4    | CCD      | 2012-04-18 | 1.8             |
| krak120421-00 | Krakow 0.4    | CCD      | 2012-04-21 | 4.6             |
| krak120423-00 | Krakow 0.4    | CCD      | 2012-04-23 | 1.7             |

Table 1—Continued

| Run Name      | Telescope      | Detector | Date       | Length<br>(hrs) |
|---------------|----------------|----------|------------|-----------------|
| krak120428-00 | Krakow 0.4     | CCD      | 2012-04-28 | 6.6             |
| krak120429-00 | Krakow 0.4     | CCD      | 2012-04-29 | 6.1             |
| krak120430-00 | Krakow 0.4     | CCD      | 2012-04-30 | 6.7             |
| mtlm120419-10 | Mt. Lemmon 1.0 | CCD      | 2012-04-19 | 1.7             |
| mtlm120420-10 | Mt. Lemmon 1.0 | CCD      | 2012-04-20 | 1.4             |
| mtlm120421-05 | Mt. Lemmon 1.0 | CCD      | 2012-04-21 | 6.2             |
| mtlm120422-07 | Mt. Lemmon 1.0 | CCD      | 2012-04-22 | 4.8             |
| na50120524-12 | NAOC 0.5       | CCD      | 2012-05-24 | 4.5             |
| na50120525-12 | NAOC 0.5       | CCD      | 2012-05-25 | 4.4             |
| na50120526-14 | NAOC 0.5       | CCD      | 2012-05-26 | 3.3             |
| na50120527-12 | NAOC 0.5       | CCD      | 2012-05-27 | 4.9             |
| na50120530-13 | NAOC 0.5       | CCD      | 2012-05-30 | 6.0             |
| pjmo120425-03 | PJMO 0.6       | CCD      | 2012-04-25 | 7.2             |
| pjmo120426-03 | PJMO 0.6       | CCD      | 2012-04-26 | 4.3             |
| prom120430-04 | PROMPT 0.4     | CCD      | 2012-04-30 | 2.2             |
| prom120430-07 | PROMPT 0.4     | CCD      | 2012-04-30 | 1.1             |
| prom120501-04 | PROMPT 0.4     | CCD      | 2012-05-01 | 4.5             |
| prom120502-04 | PROMPT 0.4     | CCD      | 2012-05-02 | 4.4             |
| prom120503-04 | PROMPT 0.4     | CCD      | 2012-05-03 | 0.9             |
| prom120504-04 | PROMPT 0.4     | CCD      | 2012-05-04 | 4.5             |
| prom120509-04 | PROMPT 0.4     | CCD      | 2012-05-09 | 4.3             |
| prom120510-04 | PROMPT 0.4     | CCD      | 2012-05-10 | 4.1             |
| prom120511-03 | PROMPT 0.4     | CCD      | 2012-05-11 | 4.1             |
| prom120512-03 | PROMPT 0.4     | CCD      | 2012-05-12 | 4.0             |
| prom120513-03 | PROMPT 0.4     | CCD      | 2012-05-13 | 2.1             |
| prom120514-03 | PROMPT 0.4     | CCD      | 2012-05-14 | 4.0             |
| prom120515-03 | PROMPT 0.4     | CCD      | 2012-05-15 | 4.0             |
| suho120501-20 | Suhora 0.6     | CCD      | 2012-05-01 | 6.0             |
| suho120502-21 | Suhora 0.6     | CCD      | 2012-05-02 | 5.1             |
| suho120503-23 | Suhora 0.6     | CCD      | 2012-05-03 | 2.4             |
| suho120505-19 | Suhora 0.6     | CCD      | 2012-05-05 | 2.1             |
| tubi120420-01 | Tubitak 1.0    | CCD      | 2012-04-20 | 1.0             |

Table 1—Continued

| Run Name      | Telescope      | Detector | Date       | Length<br>(hrs) |
|---------------|----------------|----------|------------|-----------------|
| tubi120423-19 | Tubitak 1.0    | CCD      | 2012-04-23 | 6.6             |
| tubi120512-20 | Tubitak 1.0    | CCD      | 2012-04-23 | 5.2             |
| caam130412-21 | Cannakkale 1.2 | CCD      | 2013-04-12 | 4.0             |
| caam130417-21 | Cannakkale 1.2 | CCD      | 2013-04-17 | 3.2             |
| caam130418-21 | Cannakkale 1.2 | CCD      | 2013-04-18 | 4.5             |
| krak130418-00 | Krakow 0.4     | CCD      | 2013-04-18 | 6.7             |
| krak130421-00 | Krakow 0.4     | CCD      | 2013-04-18 | 7.4             |
| krak130425-00 | Krakow 0.4     | CCD      | 2013-04-25 | 6.7             |
| krak130426-00 | Krakow 0.4     | CCD      | 2013-04-26 | 2.1             |
| naos130503-13 | NAOC 0.85      | CCD      | 2013-05-03 | 6.0             |
| pjmo130429-06 | PJMO 0.6       | CCD      | 2013-04-29 | 2.7             |
| pjmo130430-07 | PJMO 0.6       | CCD      | 2013-04-30 | 2.1             |
| pjmo130503-07 | PJMO 0.6       | CCD      | 2013-05-03 | 3.6             |
| pjmo130505-04 | PJMO 0.6       | CCD      | 2013-05-05 | 6.8             |
| suho130422-21 | Suhora 0.6     | CCD      | 2013-04-22 | 4.6             |
| suho130425-20 | Suhora 0.6     | CCD      | 2013-04-25 | 5.9             |
| suho130426-20 | Suhora 0.6     | CCD      | 2013-04-26 | 5.4             |
| suho130501-19 | Suhora 0.6     | CCD      | 2013-05-01 | 6.1             |
| suho130504-21 | Suhora 0.6     | CCD      | 2013-05-04 | 4.4             |
| suho130505-19 | Suhora 0.6     | CCD      | 2013-05-05 | 5.5             |
| caam140610-19 | Cannakkale 1.2 | CCD      | 2014-06-10 | 5.5             |
| krak140530-21 | Krakow 0.4     | CCD      | 2014-05-30 | 2.4             |
| krak140604-20 | Krakow 0.4     | CCD      | 2014-06-04 | 5.5             |
| krak140606-20 | Krakow 0.4     | CCD      | 2014-06-06 | 4.7             |
| krak140607-20 | Krakow 0.4     | CCD      | 2014-06-07 | 5.1             |
| krak140608-20 | Krakow 0.4     | CCD      | 2014-06-08 | 5.2             |
| krak140609-19 | Krakow 0.4     | CCD      | 2014-06-09 | 4.7             |
| krak140610-20 | Krakow 0.4     | CCD      | 2014-06-10 | 2.4             |
| mcao140602-05 | MCAO 0.6       | CCD      | 2014-06-02 | 3.0             |
| mcao140603-01 | MCAO 0.6       | CCD      | 2014-06-03 | 4.1             |
| mcao140607-01 | MCAO 0.6       | CCD      | 2014-06-07 | 4.0             |
| mcao140608-01 | MCAO 0.6       | CCD      | 2014-06-08 | 3.9             |

Table 1—Continued

| Run Name      | Telescope     | Detector | Date       | Length<br>(hrs) |
|---------------|---------------|----------|------------|-----------------|
| mole140526-20 | Moletai 1.65  | CCD      | 2014-05-26 | 2.5             |
| mole140527-20 | Moletai 1.65  | CCD      | 2014-05-27 | 3.7             |
| mole140605-20 | Moletai 1.65  | CCD      | 2014-06-05 | 3.3             |
| mole140607-20 | Moletai 1.65  | CCD      | 2014-06-07 | 2.9             |
| naos140605-15 | NAOC 0.85     | CCD      | 2014-06-05 | 5.7             |
| naos140606-17 | NAOC 0.85     | CCD      | 2014-06-06 | 3.9             |
| pjmo140519-04 | PJMO 0.6      | CCD      | 2014-05-19 | 4.0             |
| pjmo140529-03 | PJMO 0.6      | CCD      | 2014-05-29 | 5.4             |
| pjmo140530-03 | PJMO 0.6      | CCD      | 2014-05-30 | 4.3             |
| pjmo140531-02 | PJMO 0.6      | CCD      | 2014-05-31 | 5.0             |
| pjmo140601-03 | PJMO 0.6      | CCD      | 2014-06-01 | 3.3             |
| pjmo140602-02 | PJMO 0.6      | CCD      | 2014-06-02 | 4.5             |
| pjmo140603-02 | PJMO 0.6      | CCD      | 2014-06-03 | 7.1             |
| pjmo140604-04 | PJMO 0.6      | CCD      | 2014-06-04 | 5.8             |
| pjmo140611-03 | PJMO 0.6      | CCD      | 2014-06-11 | 5.9             |
| suho140604-19 | Suhora 0.6    | CCD      | 2014-06-04 | 4.4             |
| suho140606-20 | Suhora 0.6    | CCD      | 2014-06-06 | 4.7             |
| ters140604-20 | Terskol 0.6   | CCD      | 2014-06-04 | 2.0             |
| ters140619-20 | Terskol 0.6   | CCD      | 2014-06-19 | 6.0             |
| tsao140524-17 | Tien Shan 1.0 | CCD      | 2014-05-24 | 4.7             |
| tsao140525-16 | Tien Shan 1.0 | CCD      | 2014-05-25 | 6.0             |
| tsao140610-15 | Tien Shan 1.0 | CCD      | 2014-06-10 | 1.0             |
| tsao140611-16 | Tien Shan 1.0 | CCD      | 2014-06-11 | 6.0             |
| tsao140613-17 | Tien Shan 1.0 | CCD      | 2014-06-13 | 4.6             |
| tsao140616-19 | Tien Shan 1.0 | CCD      | 2014-06-16 | 1.4             |
| tsao140619-17 | Tien Shan 1.0 | CCD      | 2014-06-19 | 4.8             |
| tsao140620-18 | Tien Shan 1.0 | CCD      | 2014-06-20 | 3.1             |
| tsao140627-16 | Tien Shan 1.0 | CCD      | 2014-06-27 | 5.9             |
| tsao140628-16 | Tien Shan 1.0 | CCD      | 2014-06-28 | 5.9             |
| krak150420-20 | Krakow 0.4    | CCD      | 2015-05-20 | 2.1             |
| krak150421-19 | Krakow 0.4    | CCD      | 2015-05-21 | 7.5             |
| pjmo150420-03 | PJMO 0.6      | CCD      | 2015-04-20 | 7.3             |



Table 1—Continued

| Run Name      | Telescope  | Detector | Date       | Length<br>(hrs) |
|---------------|------------|----------|------------|-----------------|
| pjmo150421-05 | PJMO 0.6   | CCD      | 2015-04-21 | 5.4             |
| pjmo150422-03 | PJMO 0.6   | CCD      | 2015-04-22 | 4.7             |
| pjmo150426-03 | PJMO 0.6   | CCD      | 2015-04-26 | 7.4             |
| prom150428-08 | PROMPT 0.4 | CCD      | 2015-04-28 | 6.5             |
| prom150429-03 | PROMPT 0.6 | CCD      | 2015-04-29 | 6.6             |
| prom150501-03 | PROMPT 0.6 | CCD      | 2015-05-01 | 6.2             |

Table 2. 1982-2006 Detected Independent Frequencies

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| 1982                        |             |                  |              |
| $1236.483 \pm 0.07$         | 808.75      | $16.70 \pm 0.06$ | 9.4          |
| $1431.112 \pm 0.04$         | 698.76      | $30.86 \pm 0.06$ | 11.4         |
| $1613.842 \pm 0.05$         | 619.65      | $26.12 \pm 0.06$ | 9.7          |
| $1618.845 \pm 0.06$         | 617.72      | $29.34 \pm 0.06$ | 10.9         |
| $2368.563 \pm 0.10$         | 422.20      | $12.52 \pm 0.06$ | 5.8          |
| 1984                        |             |                  |              |
| $1124.701 \pm 3.5$          | 889.13      | $24.7 \pm 1.1$   | 7.2          |
| $1626.474 \pm 3.5$          | 614.83      | $25.9 \pm 1.1$   | 6.8          |
| 1985                        |             |                  |              |
| $1166.944 \pm 0.05$         | 856.94      | $6.43 \pm 0.06$  | 11           |
| $1176.489 \pm 0.03$         | 849.99      | $35.70 \pm 0.06$ | 16.0         |
| $1614.554 \pm 0.04$         | 619.37      | $9.15 \pm 0.06$  | 4.6          |
| $2351.762 \pm 0.04$         | 425.22      | $14.34 \pm 0.06$ | 12           |
| 1986                        |             |                  |              |
| $1081.025 \pm 0.03$         | 925.05      | $5.1 \pm 0.4$    | 4.6          |
| $1160.678 \pm 0.03$         | 861.57      | $4.9 \pm 0.4$    | 4.5          |
| $1223.803 \pm 0.02$         | 817.12      | $9.3 \pm 0.4$    | 9.1          |
| $1233.585 \pm 0.01$         | 810.65      | $14.9 \pm 0.4$   | 14.5         |
| $1385.240 \pm 0.03$         | 721.90      | $6.7 \pm 0.4$    | 6.4          |
| $1426.181 \pm 0.01$         | 701.17      | $14.2 \pm 0.4$   | 14.3         |
| $1525.030 \pm 0.01$         | 655.72      | $21.3 \pm 0.4$   | 21.4         |
| $1611.691 \pm 0.01$         | 620.47      | $20.9 \pm 0.4$   | 20.9         |

Table 2—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $2157.783 \pm 0.04$         | 463.44      | $2.4 \pm 0.4$    | 4            |
| $2165.564 \pm 0.03$         | 461.77      | $4.8 \pm 0.4$    | 4            |
| $2368.938 \pm 0.03$         | 422.13      | $5.8 \pm 0.4$    | 5.6          |
| 1990                        |             |                  |              |
| $1112.933 \pm 0.03$         | 898.53      | $2.31 \pm 0.32$  | 4.0          |
| $1114.180 \pm 0.03$         | 897.52      | $2.44 \pm 0.32$  | 4.1          |
| $1118.324 \pm 0.02$         | 894.20      | $5.24 \pm 0.32$  | 8.9          |
| $1119.089 \pm 0.03$         | 893.58      | $2.83 \pm 0.32$  | 4.8          |
| $1224.216 \pm 0.03$         | 816.83      | $22.17 \pm 0.32$ | 4.0          |
| $1233.413 \pm 0.03$         | 810.76      | $4.80 \pm 0.32$  | 8.4          |
| $1245.395 \pm 0.03$         | 802.96      | $2.17 \pm 0.32$  | 4.0          |
| $1288.995 \pm 0.03$         | 775.80      | $3.50 \pm 0.32$  | 6.3          |
| $1291.229 \pm 0.03$         | 774.46      | $4.31 \pm 0.32$  | 7.6          |
| $1295.400 \pm 0.03$         | 771.96      | $3.27 \pm 0.32$  | 5.8          |
| $1297.540 \pm 0.01$         | 770.69      | $14.76 \pm 0.32$ | 26.1         |
| $1304.075 \pm 0.03$         | 766.83      | $4.67 \pm 0.32$  | 8.3          |
| $1355.447 \pm 0.03$         | 737.76      | $2.20 \pm 0.32$  | 4.0          |
| $1361.728 \pm 0.03$         | 734.36      | $2.90 \pm 0.32$  | 5.2          |
| $1368.588 \pm 0.03$         | 730.68      | $3.32 \pm 0.32$  | 5.9          |
| $1375.434 \pm 0.03$         | 727.04      | $3.18 \pm 0.32$  | 5.7          |
| $1421.041 \pm 0.02$         | 703.71      | $8.22 \pm 0.32$  | 15.1         |
| $1423.704 \pm 0.03$         | 702.39      | $3.11 \pm 0.32$  | 5.8          |
| $1427.365 \pm 0.01$         | 700.59      | $19.39 \pm 0.32$ | 35.8         |
| $1428.663 \pm 0.03$         | 699.96      | $2.61 \pm 0.32$  | 4.8          |
| $1433.729 \pm 0.02$         | 697.48      | $7.29 \pm 0.32$  | 13.5         |
| $1435.209 \pm 0.03$         | 696.76      | $3.50 \pm 0.32$  | 6.5          |
| $1512.798 \pm 0.02$         | 661.03      | $5.57 \pm 0.32$  | 10.7         |
| $1518.661 \pm 0.02$         | 658.47      | $8.34 \pm 0.32$  | 15.9         |

Table 2—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $1519.372 \pm 0.02$         | 658.17      | $5.88 \pm 0.32$  | 11.2         |
| $1524.924 \pm 0.02$         | 655.77      | $5.98 \pm 0.32$  | 11.2         |
| $1525.498 \pm 0.02$         | 655.52      | $6.95 \pm 0.32$  | 13.0         |
| $1611.741 \pm 0.02$         | 620.45      | $6.09 \pm 0.32$  | 12.3         |
| $1617.380 \pm 0.03$         | 618.28      | $4.69 \pm 0.32$  | 9.7          |
| $1623.709 \pm 0.02$         | 615.87      | $5.04 \pm 0.32$  | 10.2         |
| $2154.009 \pm 0.03$         | 464.25      | $4.40 \pm 0.32$  | 11.2         |
| $2157.765 \pm 0.03$         | 463.44      | $2.27 \pm 0.32$  | 5.8          |
| $2358.946 \pm 0.02$         | 423.92      | $5.63 \pm 0.32$  | 14.1         |
| $2362.507 \pm 0.02$         | 423.28      | $5.71 \pm 0.32$  | 14.3         |
| $2366.408 \pm 0.03$         | 422.58      | $4.59 \pm 0.32$  | 11.5         |
| 1991                        |             |                  |              |
| $1296.087 \pm 0.13$         | 771.55      | $10.11 \pm 0.41$ | 17.7         |
| $1296.577 \pm 0.13$         | 771.26      | $18.25 \pm 0.41$ | 29.8         |
| $1308.777 \pm 0.05$         | 764.07      | $5.02 \pm 0.41$  | 9.9          |
| $1396.945 \pm 0.05$         | 715.85      | $3.95 \pm 0.41$  | 8.4          |
| $1419.934 \pm 0.01$         | 704.26      | $29.3 \pm 0.41$  | 45.6         |
| $1423.333 \pm 0.04$         | 702.58      | $5.23 \pm 0.41$  | 6.3          |
| $1427.062 \pm 0.04$         | 700.74      | $5.02 \pm 0.41$  | 7.4          |
| $1443.381 \pm 0.05$         | 692.82      | $4.81 \pm 0.41$  | 9.7          |
| $2150.307 \pm 0.05$         | 465.05      | $2.36 \pm 0.41$  | 4.1          |
| $2154.389 \pm 0.05$         | 464.17      | $2.79 \pm 0.41$  | 4.6          |
| $2157.939 \pm 0.05$         | 463.41      | $2.62 \pm 0.41$  | 4.4          |
| $2370.121 \pm 0.04$         | 421.92      | $5.01 \pm 0.41$  | 9.2          |
| $2374.211 \pm 0.05$         | 421.19      | $3.26 \pm 0.41$  | 5.7          |
| $2378.195 \pm 0.04$         | 420.49      | $5.58 \pm 0.41$  | 10.7         |
| 1992                        |             |                  |              |

Table 2—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $1035.357 \pm 0.05$         | 965.85      | $7.35 \pm 0.31$  | 4.1          |
| $1101.952 \pm 0.05$         | 907.48      | $6.40 \pm 0.31$  | 3.7          |
| $1195.349 \pm 0.003$        | 836.58      | $6.90 \pm 0.31$  | 4.1          |
| $1233.263 \pm 0.001$        | 810.86      | $20.79 \pm 0.31$ | 13.5         |
| $1242.854 \pm 0.003$        | 804.60      | $14.20 \pm 0.31$ | 8.8          |
| $1265.279 \pm 0.003$        | 790.34      | $12.25 \pm 0.31$ | 7.1          |
| $1420.845 \pm 0.001$        | 703.81      | $20.59 \pm 0.31$ | 13.1         |
| $1438.411 \pm 0.003$        | 695.21      | $17.71 \pm 0.31$ | 11.9         |
| $1622.795 \pm 0.003$        | 616.22      | $14.26 \pm 0.31$ | 9.2          |
| $1628.812 \pm 0.004$        | 613.94      | $10.9 \pm 0.317$ | 6.5          |
| $2162.104 \pm 0.007$        | 462.51      | $4.26 \pm 0.31$  | 4.2          |
| $2166.094 \pm 0.007$        | 461.66      | $5.66 \pm 0.31$  | 4.8          |
| $2351.041 \pm 0.007$        | 425.34      | $6.75 \pm 0.31$  | 5.2          |
| $2359.162 \pm 0.007$        | 423.88      | $6.55 \pm 0.31$  | 9.2          |
| $2366.443 \pm 0.007$        | 422.58      | $7.17 \pm 0.31$  | 6.3          |
| 1994                        |             |                  |              |
| $939.264 \pm 0.007$         | 1064.66     | $2.36 \pm 0.10$  | 5.7          |
| $1024.773 \pm 0.011$        | 975.83      | $3.46 \pm 0.10$  | 8.8          |
| $1064.931 \pm 0.012$        | 939.03      | $2.82 \pm 0.10$  | 7.4          |
| $1106.833 \pm 0.013$        | 903.48      | $2.56 \pm 0.10$  | 7.0          |
| $1113.548 \pm 0.012$        | 898.03      | $3.95 \pm 0.10$  | 9.4          |
| $1164.637 \pm 0.012$        | 858.64      | $3.12 \pm 0.10$  | 8.6          |
| $1176.684 \pm 0.013$        | 849.85      | $2.79 \pm 0.10$  | 7.2          |
| $1224.306 \pm 0.012$        | 816.79      | $3.28 \pm 0.10$  | 9.1          |
| $1234.488 \pm 0.013$        | 810.05      | $2.70 \pm 0.10$  | 7.7          |
| $1235.491 \pm 0.005$        | 809.39      | $13.13 \pm 0.10$ | 37.3         |
| $1242.357 \pm 0.013$        | 804.92      | $3.33 \pm 0.10$  | 9.1          |

Table 2—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $1246.494 \pm 0.013$        | 802.25      | $2.69 \pm 0.10$  | 7.8          |
| $1286.550 \pm 0.003$        | 777.27      | $9.45 \pm 0.10$  | 27.6         |
| $1291.023 \pm 0.009$        | 774.58      | $6.07 \pm 0.10$  | 17.8         |
| $1293.244 \pm 0.009$        | 773.25      | $5.85 \pm 0.10$  | 16.1         |
| $1297.737 \pm 0.005$        | 770.57      | $21.5 \pm 0.10$  | 61.9         |
| $1298.710 \pm 0.010$        | 769.99      | $4.45 \pm 0.10$  | 12.5         |
| $1304.464 \pm 0.009$        | 766.64      | $6.84 \pm 0.10$  | 19.9         |
| $1305.352 \pm 0.013$        | 766.08      | $2.54 \pm 0.10$  | 7.0          |
| $1309.003 \pm 0.013$        | 763.94      | $2.76 \pm 0.10$  | 8.4          |
| $1312.045 \pm 0.013$        | 762.17      | $2.67 \pm 0.10$  | 7.9          |
| $1419.641 \pm 0.003$        | 704.40      | $18.70 \pm 0.10$ | 54.4         |
| $1422.947 \pm 0.013$        | 702.77      | $2.86 \pm 0.10$  | 8.2          |
| $1426.395 \pm 0.003$        | 701.07      | $16.05 \pm 0.10$ | 46.6         |
| $1430.851 \pm 0.005$        | 698.88      | $10.35 \pm 0.10$ | 30.1         |
| $1433.167 \pm 0.010$        | 697.75      | $4.06 \pm 0.10$  | 11.6         |
| $1437.607 \pm 0.006$        | 695.60      | $8.28 \pm 0.10$  | 24.1         |
| $1438.523 \pm 0.011$        | 695.16      | $3.68 \pm 0.10$  | 10.7         |
| $1440.997 \pm 0.012$        | 693.96      | $2.48 \pm 0.10$  | 7.2          |
| $1441.910 \pm 0.009$        | 693.52      | $4.09 \pm 0.10$  | 11.9         |
| $1611.351 \pm 0.009$        | 620.60      | $5.12 \pm 0.10$  | 14.2         |
| $1617.450 \pm 0.010$        | 618.26      | $3.61 \pm 0.10$  | 9.5          |
| $1618.545 \pm 0.009$        | 617.84      | $4.04 \pm 0.10$  | 11.2         |
| $1624.624 \pm 0.009$        | 615.53      | $5.83 \pm 0.10$  | 16.1         |
| $1625.634 \pm 0.009$        | 615.14      | $4.89 \pm 0.10$  | 13.9         |
| $2150.498 \pm 0.010$        | 465.01      | $3.22 \pm 0.10$  | 9.4          |
| $2154.130 \pm 0.010$        | 464.22      | $4.75 \pm 0.10$  | 13.7         |
| $2157.844 \pm 0.012$        | 463.43      | $2.70 \pm 0.10$  | 7.6          |
| $2358.880 \pm 0.010$        | 423.93      | $4.53 \pm 0.10$  | 13.5         |
| $2362.636 \pm 0.005$        | 423.26      | $9.29 \pm 0.10$  | 31.2         |
| $2366.505 \pm 0.011$        | 422.56      | $4.26 \pm 0.10$  | 13.2         |

Table 2—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| 1996                        |             |                  |              |
| $937.695 \pm 0.028$         | 1066.44     | $4.27 \pm 0.28$  | 3.5          |
| $1024.276 \pm 0.028$        | 976.30      | $6.78 \pm 0.28$  | 6.4          |
| $1104.483 \pm 0.025$        | 905.40      | $5.55 \pm 0.28$  | 5.1          |
| $1178.461 \pm 0.025$        | 848.56      | $5.34 \pm 0.28$  | 5.5          |
| $1227.936 \pm 0.022$        | 814.37      | $7.66 \pm 0.28$  | 7.5          |
| $1233.14 \pm 0.015$         | 810.94      | $13.50 \pm 0.28$ | 12.1         |
| $1234.506 \pm 0.015$        | 810.04      | $12.53 \pm 0.28$ | 11.2         |
| $1247.959 \pm 0.022$        | 801.31      | $7.84 \pm 0.28$  | 4.1          |
| $1261.275 \pm 0.015$        | 792.85      | $11.08 \pm 0.28$ | 11.6         |
| $1291.169 \pm 0.015$        | 774.49      | $10.71 \pm 0.28$ | 9.7          |
| $1294.899 \pm 0.025$        | 772.26      | $6.95 \pm 0.28$  | 7.9          |
| $1297.194 \pm 0.010$        | 770.89      | $22.05 \pm 0.28$ | 2.4          |
| $1420.057 \pm 0.010$        | 704.20      | $20.26 \pm 0.28$ | 19.7         |
| $1426.443 \pm 0.010$        | 701.04      | $18.41 \pm 0.28$ | 8.3          |
| $1430.253 \pm 0.012$        | 699.18      | $13.15 \pm 0.28$ | 12.8         |
| $1436.261 \pm 0.022$        | 696.25      | $8.20 \pm 0.28$  | 9.1          |
| $1628.296 \pm 0.028$        | 614.14      | $6.13 \pm 0.28$  | 6.3          |
| $2154.117 \pm 0.028$        | 464.23      | $7.40 \pm 0.28$  | 7.1          |
| $2358.806 \pm 0.025$        | 423.94      | $6.01 \pm 0.28$  | 4.7          |
| $2362.617 \pm 0.014$        | 423.26      | $9.87 \pm 0.28$  | 9.0          |
| $2367.288 \pm 0.025$        | 422.42      | $8.92 \pm 0.28$  | 11.6         |
| 2000                        |             |                  |              |
| $938.991 \pm 0.015$         | 1064.95     | $3.09 \pm 0.01$  | 9.4          |
| $946.238 \pm 0.015$         | 1056.82     | $1.20 \pm 0.01$  | 4.0          |
| $1110.999 \pm 0.015$        | 900.09      | $1.98 \pm 0.01$  | 7.0          |

Table 2—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $1171.564 \pm 0.015$        | 853.56      | $1.89 \pm 0.01$  | 7.0          |
| $1173.021 \pm 0.012$        | 852.50      | $2.32 \pm 0.01$  | 8.8          |
| $1251.851 \pm 0.004$        | 798.82      | $3.21 \pm 0.01$  | 12.5         |
| $1254.503 \pm 0.002$        | 797.13      | $8.60 \pm 0.01$  | 33.1         |
| $1255.583 \pm 0.002$        | 796.44      | $14.72 \pm 0.01$ | 67.1         |
| $1256.248 \pm 0.005$        | 796.02      | $8.04 \pm 0.01$  | 42.6         |
| $1257.268 \pm 0.010$        | 795.38      | $3.05 \pm 0.01$  | 15.2         |
| $1258.288 \pm 0.010$        | 794.73      | $2.06 \pm 0.01$  | 5.7          |
| $1296.603 \pm 0.001$        | 771.25      | $28.08 \pm 0.01$ | 109.5        |
| $1378.795 \pm 0.008$        | 725.27      | $4.33 \pm 0.01$  | 17.3         |
| $1379.737 \pm 0.010$        | 724.78      | $2.64 \pm 0.01$  | 10.4         |
| $1420.101 \pm 0.001$        | 704.18      | $29.78 \pm 0.01$ | 118.8        |
| $1423.597 \pm 0.010$        | 702.45      | $3.08 \pm 0.01$  | 12.3         |
| $1736.660 \pm 0.014$        | 575.82      | $1.02 \pm 0.01$  | 4.0          |
| $2150.515 \pm 0.013$        | 465.01      | $2.99 \pm 0.01$  | 11.2         |
| $2154.040 \pm 0.013$        | 464.24      | $5.38 \pm 0.01$  | 19.9         |
| $2157.736 \pm 0.012$        | 463.45      | $2.51 \pm 0.01$  | 9.5          |
| $2359.118 \pm 0.008$        | 423.89      | $5.51 \pm 0.01$  | 23.7         |
| $2366.271 \pm 0.010$        | 422.61      | $5.90 \pm 0.01$  | 25.1         |
| 2006                        |             |                  |              |
| $923.976 \pm 0.001$         | 1082.28     | $1.43 \pm 0.01$  | 5.2          |
| $938.216 \pm 0.002$         | 1065.85     | $1.23 \pm 0.01$  | 4.4          |
| $1024.496 \pm 0.002$        | 976.09      | $1.44 \pm 0.01$  | 4.8          |
| $1033.760 \pm 0.002$        | 967.34      | $1.85 \pm 0.01$  | 6.9          |
| $1039.075 \pm 0.001$        | 962.39      | $7.95 \pm 0.01$  | 27.2         |
| $1039.474 \pm 0.001$        | 962.02      | $2.84 \pm 0.01$  | 9.7          |
| $1041.535 \pm 0.001$        | 960.12      | $1.18 \pm 0.01$  | 4.3          |
| $1044.381 \pm 0.002$        | 957.50      | $1.63 \pm 0.01$  | 6.2          |



Table 2—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $1113.582 \pm 0.001$        | 898.02      | $2.71 \pm 0.01$  | 9.4          |
| $1120.404 \pm 0.001$        | 892.53      | $2.09 \pm 0.01$  | 7.3          |
| $1120.902 \pm 0.001$        | 892.14      | $2.98 \pm 0.01$  | 10.2         |
| $1121.704 \pm 0.002$        | 891.50      | $1.26 \pm 0.01$  | 4.4          |
| $1130.144 \pm 0.002$        | 884.84      | $1.91 \pm 0.01$  | 7.3          |
| $1161.552 \pm 0.001$        | 860.92      | $2.74 \pm 0.01$  | 9.6          |
| $1173.015 \pm 0.001$        | 852.50      | $7.26 \pm 0.01$  | 25.4         |
| $1178.096 \pm 0.002$        | 848.83      | $1.13 \pm 0.01$  | 4.3          |
| $1184.470 \pm 0.002$        | 844.26      | $1.64 \pm 0.01$  | 6.2          |
| $1222.199 \pm 0.002$        | 818.20      | $1.72 \pm 0.01$  | 6.3          |
| $1222.751 \pm 0.001$        | 817.83      | $5.04 \pm 0.01$  | 18.3         |
| $1222.945 \pm 0.001$        | 817.70      | $4.59 \pm 0.01$  | 5.1          |
| $1228.185 \pm 0.002$        | 814.21      | $2.71 \pm 0.01$  | 9.4          |
| $1228.791 \pm 0.001$        | 813.81      | $5.27 \pm 0.01$  | 19.0         |
| $1234.124 \pm 0.001$        | 810.29      | $24.87 \pm 0.01$ | 88.0         |
| $1239.510 \pm 0.001$        | 806.77      | $5.07 \pm 0.01$  | 18.3         |
| $1240.237 \pm 0.002$        | 806.30      | $2.85 \pm 0.01$  | 9.9          |
| $1244.790 \pm 0.002$        | 803.35      | $1.90 \pm 0.01$  | 6.9          |
| $1245.219 \pm 0.001$        | 803.07      | $4.75 \pm 0.01$  | 17.1         |
| $1246.032 \pm 0.001$        | 802.55      | $4.44 \pm 0.01$  | 15.2         |
| $1421.012 \pm 0.002$        | 703.72      | $2.81 \pm 0.01$  | 7.1          |
| $1429.209 \pm 0.001$        | 699.69      | $5.65 \pm 0.01$  | 22.5         |
| $1512.141 \pm 0.002$        | 661.31      | $1.79 \pm 0.01$  | 6.4          |
| $1736.301 \pm 0.001$        | 575.94      | $16.38 \pm 0.01$ | 75.2         |
| $1737.962 \pm 0.002$        | 575.39      | $1.80 \pm 0.01$  | 7.6          |
| $1741.666 \pm 0.001$        | 574.16      | $11.01 \pm 0.01$ | 49.7         |
| $1743.733 \pm 0.001$        | 573.48      | $5.59 \pm 0.01$  | 8.4          |
| $1749.083 \pm 0.001$        | 571.73      | $11.88 \pm 0.01$ | 50.2         |
| $1856.845 \pm 0.002$        | 538.55      | $1.44 \pm 0.01$  | 6.4          |
| $2150.393 \pm 0.001$        | 465.03      | $4.06 \pm 0.01$  | 17.3         |

Table 2—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $2154.223 \pm 0.001$        | 464.20      | $5.50 \pm 0.01$  | 21.8         |
| $2158.073 \pm 0.001$        | 463.38      | $7.24 \pm 0.01$  | 29.0         |
| $2359.052 \pm 0.001$        | 423.90      | $5.94 \pm 0.01$  | 22.1         |
| $2363.058 \pm 0.002$        | 423.18      | $1.7 \pm 0.011$  | 6.0          |
| $2366.524 \pm 0.001$        | 422.56      | $6.31 \pm 0.01$  | 23.0         |

Table 3. 2007-2015 Detected Independent Frequencies

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| 2007                        |             |                  |              |
| $1088.951 \pm 0.05$         | 918.31      | $6.9 \pm 0.7$    | 5            |
| $1121.169 \pm 0.04$         | 891.93      | $10.19 \pm 0.7$  | 8            |
| $1233.956 \pm 0.02$         | 810.40      | $23 \pm 0.7$     | 20           |
| $1251.1930.05$              | 799.24      | $9.56 \pm 0.7$   | 8            |
| $1735.4440.03$              | 576.22      | $13.47 \pm 0.7$  | 13           |
| $2156.2880.03$              | 463.76      | $7.26 \pm 0.7$   | 7            |
| 2008                        |             |                  |              |
| $1235.745 \pm 0.004$        | 809.23      | $25.14 \pm 0.33$ | 23.9         |
| $1735.716 \pm 0.004$        | 576.13      | $21.77 \pm 0.33$ | 22.6         |
| $1741.459 \pm 0.009$        | 574.23      | $10.14 \pm 0.33$ | 9.8          |
| $1750.185 \pm 0.012$        | 571.37      | $4.47 \pm 0.33$  | 5.6          |
| $2150.245 \pm 0.011$        | 465.06      | $5.66 \pm 0.33$  | 7.7          |
| $2158.416 \pm 0.01$         | 463.30      | $11.07 \pm 0.33$ | 11           |
| $2358.895 \pm 0.01$         | 423.93      | $7.95 \pm 0.33$  | 10.2         |
| $2366.872 \pm 0.01$         | 422.50      | $7.93 \pm 0.33$  | 10.2         |
| 2009                        |             |                  |              |
| $1088.445 \pm 0.018$        | 918.74      | $7.82 \pm 0.28$  | 9            |
| $1235.845 \pm 0.007$        | 809.16      | $9.64 \pm 0.28$  | 22.4         |
| $1236.849 \pm 0.018$        | 808.51      | $7.91 \pm 0.28$  | 9.1          |
| $1300.029 \pm 0.018$        | 576.29      | $4.10 \pm 0.28$  | 4.7          |
| $1735.699 \pm 0.006$        | 576.14      | $20.60 \pm 0.28$ | 22.8         |
| $1741.415 \pm 0.018$        | 574.25      | $7.15 \pm 0.28$  | 8.8          |
| $1750.525 \pm 0.012$        | 571.26      | $5.98 \pm 0.28$  | 6.8          |
| $2150.261 \pm 0.014$        | 465.06      | $5.22 \pm 0.28$  | 6.5          |

Table 3—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma  | Signal/Noise |
|-----------------------------|-------------|-------------------|--------------|
| $2158.474 \pm 0.003$        | 463.29      | $10.72 \pm 0.28$  | 13.4         |
| $2358.952 \pm 0.026$        | 423.92      | $5.89 \pm 0.28$   | 8.1          |
| $2366.893 \pm 0.018$        | 422.49      | $7.28 \pm 0.28$   | 10.1         |
| 2010                        |             |                   |              |
| $1087.790 \pm 0.024$        | 919.29      | $2.47 \pm 0.15$   | 5.1          |
| $1104.397 \pm 0.023$        | 891.34      | $2.57 \pm 0.15$   | 5.1          |
| $1112.868 \pm 0.009$        | 898.58      | $6.22 \pm 0.15$   | 13.2         |
| $1121.901 \pm 0.005$        | 891.34      | $12.6 \pm 0.15$   | 26.6         |
| $1122.674 \pm 0.021$        | 889.89      | $2.73 \pm 0.15$   | 5.2          |
| $1123.731 \pm 0.020$        | 891.33      | $3.31 \pm 0.15$   | 7.1          |
| $1200.999 \pm 0.017$        | 832.64      | $3.33 \pm 0.15$   | 7.1          |
| $1211.906 \pm 0.021$        | 825.15      | $2.99 \pm 0.15$   | 6.4          |
| $1236.278 \pm 0.003$        | 808.88      | $18.9 \pm 0.15$   | 39.7         |
| $1237.011 \pm 0.023$        | 808.41      | $4.10 \pm 0.15$   | 9.2          |
| $1241.721 \pm 0.014$        | 805.33      | $3.32 \pm 0.15$   | 7.0          |
| $1299.274 \pm 0.018$        | 769.66      | $3.89 \pm 0.15$   | 7.6          |
| $1735.786 \pm 0.003$        | 576.11      | $21.72 \pm 0.015$ | 46.9         |
| $1741.367 \pm 0.004$        | 574.26      | $8.34 \pm 0.15$   | 17.2         |
| $1745.084 \pm 0.025$        | 573.04      | $2.53 \pm 0.15$   | 5.2          |
| $1750.642 \pm 0.005$        | 571.19      | $10.93 \pm 0.15$  | 25.2         |
| $1859.431 \pm 0.023$        | 537.80      | $2.50 \pm 0.15$   | 5.2          |
| $2008.645 \pm 0.025$        | 497.85      | $2.66 \pm 0.15$   | 5.2          |
| $2150.138 \pm 0.025$        | 465.09      | $4.52 \pm 0.15$   | 9.7          |
| $2154.432 \pm 0.025$        | 464.16      | $2.17 \pm 0.15$   | 5.1          |
| $2158.524 \pm 0.006$        | 463.28      | $9.95 \pm 0.15$   | 23.4         |
| $2358.818 \pm 0.008$        | 423.94      | $9.05 \pm 0.15$   | 23.1         |
| $2362.603 \pm 0.017$        | 423.26      | $3.19 \pm 0.15$   | 6.7          |
| $2366.990 \pm 0.008$        | 422.48      | $7.40 \pm 0.15$   | 19           |

Table 3—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| 2011                        |             |                  |              |
| $954.564 \pm 0.008$         | 1047.59     | $5.2 \pm 0.17$   | 10.9         |
| $954.922 \pm 0.02$          | 1047.21     | $4.22 \pm 0.17$  | 8.7          |
| $1113.07 \pm 0.013$         | 898.45      | $5.72 \pm 0.17$  | 11.6         |
| $1113.641 \pm 0.02$         | 897.96      | $3.2 \pm 0.17$   | 6.4          |
| $1235.415 \pm 0.004$        | 809.44      | $9.76 \pm 0.17$  | 20.1         |
| $1362.917 \pm 0.03$         | 733.72      | $1.66 \pm 0.17$  | 4.0          |
| $1511.537 \pm 0.03$         | 661.58      | $1.85 \pm 0.17$  | 4.3          |
| $1614.827 \pm 0.016$        | 619.26      | $4.40 \pm 0.17$  | 10.5         |
| $1735.871 \pm 0.003$        | 576.08      | $20.3 \pm 0.17$  | 150.5        |
| $1736.14 \pm 0.037$         | 575.99      | $4.70 \pm 0.17$  | 11.1         |
| $1747.123 \pm 0.037$        | 572.37      | $3.30 \pm 0.17$  | 7.9          |
| $1750.833 \pm 0.013$        | 571.16      | $3.77 \pm 0.17$  | 8.9          |
| $1856.872 \pm 0.03$         | 538.54      | $1.62 \pm 0.17$  | 3.8          |
| $2008.742 \pm 0.02$         | 497.82      | $3.00 \pm 0.17$  | 7.4          |
| $2150.138 \pm 0.016$        | 465.09      | $4.98 \pm 0.17$  | 13.4         |
| $2154.294 \pm 0.028$        | 464.19      | $3.73 \pm 0.17$  | 8.9          |
| $2158.571 \pm 0.007$        | 463.27      | $9.10 \pm 0.17$  | 24.6         |
| $2358.707 \pm 0.009$        | 423.96      | $7.63 \pm 0.17$  | 27.4         |
| $2359.911 \pm 0.04$         | 423.74      | $1.58 \pm 0.17$  | 2.47         |
| $2366.986 \pm$              | 422.48      | $8.9 \pm 0.17$   | 26.4         |
| 2012                        |             |                  |              |
| $1113.254 \pm 0.014$        | 898.27      | $4.1 \pm 0.24$   | 6.2          |
| $1161.469 \pm 0.01$         | 860.98      | $5.21 \pm 0.24$  | 7.7          |
| $1211.371 \pm 0.006$        | 825.51      | $9.89 \pm 0.24$  | 14.7         |
| $1212.822 \pm 0.009$        | 824.52      | $6.46 \pm 0.24$  | 9.6          |

Table 3—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $1213.995 \pm 0.006$        | 823.73      | $8.66 \pm 0.24$  | 12.8         |
| $1215.915 \pm 0.01$         | 822.43      | $5.72 \pm 0.24$  | 8.5          |
| $1222.598 \pm 0.004$        | 817.930     | $11.53 \pm 0.24$ | 17.2         |
| $1226.836 \pm 0.008$        | 815.11      | $6.89 \pm 0.24$  | 10.3         |
| $1233.355 \pm 0.004$        | 810.80      | $12.53 \pm 0.24$ | 18.6         |
| $1235.431 \pm 0.009$        | 809.43      | $6.70 \pm 0.24$  | 9.9          |
| $1246.397 \pm 0.004$        | 802.31      | $13.35 \pm 0.24$ | 19.9         |
| $1258.482 \pm 0.009$        | 794.61      | $6.26 \pm 0.24$  | 9.3          |
| $1723.487 \pm 0.015$        | 580.22      | $4.30 \pm 0.24$  | 6.1          |
| $1734.393 \pm 0.006$        | 576.57      | $8.54 \pm 0.24$  | 12.1         |
| $1735.975 \pm 0.004$        | 576.05      | $12.4 \pm 0.24$  | 17.7         |
| $1745.543 \pm 0.012$        | 572.89      | $4.44 \pm 0.24$  | 6.3          |
| $1747.152 \pm 0.008$        | 572.36      | $6.74 \pm 0.24$  | 9.5          |
| $1748.895 \pm 0.009$        | 571.79      | $6.36 \pm 0.24$  | 9.0          |
| $1750.345 \pm 0.008$        | 571.32      | $6.19 \pm 0.24$  | 8.9          |
| $2150.072 \pm 0.02$         | 465.10      | $2.74 \pm 0.24$  | 4.0          |
| $2155.981 \pm 0.015$        | 463.83      | $3.68 \pm 0.24$  | 5.2          |
| $2158.563 \pm 0.006$        | 463.27      | $8.86 \pm 0.24$  | 12.2         |
| $2181.89 \pm 0.015$         | 458.32      | $3.60 \pm 0.24$  | 4.9          |
| $2355.788 \pm 0.01$         | 424.47      | $5.48 \pm 0.24$  | 8.7          |
| $2358.721 \pm 0.01$         | 423.96      | $5.17 \pm 0.24$  | 8.4          |
| $2366.807 \pm 0.009$        | 422.51      | $5.67 \pm 0.24$  | 9.4          |
| $2372.715 \pm 0.015$        | 421.46      | $3.81 \pm 0.24$  | 6.2          |
| 2013                        |             |                  |              |
| $1086.678 \pm 0.007$        | 920.24      | $5.28 \pm 0.18$  | 6.5          |
| $1097.639 \pm 0.014$        | 911.05      | $5.28 \pm 0.18$  | 5.4          |
| $1123.417 \pm 0.023$        | 890.14      | $4.46 \pm 0.18$  | 5.9          |
| $1235.981 \pm 0.005$        | 809.07      | $15.5 \pm 0.18$  | 15.0         |

Table 3—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $1241.938 \pm 0.013$        | 805.19      | $7.50 \pm 0.18$  | 6.4          |
| $1362.571 \pm 0.027$        | 733.91      | $4.74 \pm 0.18$  | 5.2          |
| $1614.872 \pm 0.007$        | 619.24      | $10.3 \pm 0.18$  | 10.2         |
| $1629.282 \pm 0.015$        | 613.77      | $4.80 \pm 0.18$  | 5.3          |
| $1736.468 \pm 0.008$        | 575.88      | $11.1 \pm 0.18$  | 11.4         |
| $1737.008 \pm 0.008$        | 575.70      | $11.2 \pm 0.18$  | 11.4         |
| $1746.007 \pm 0.024$        | 572.73      | $5.66 \pm 0.18$  | 5.7          |
| $2158.532 \pm 0.006$        | 463.28      | $8.32 \pm 0.18$  | 9.4          |
| $2162.432 \pm 0.014$        | 462.44      | $5.55 \pm 0.18$  | 6.4          |
| $2358.635 \pm 0.007$        | 423.97      | $7.50 \pm 0.18$  | 9.2          |
| $2364.019 \pm 0.026$        | 423.013     | $3.20 \pm 0.18$  | 6.4          |
| $2367.037 \pm 0.014$        | 422.47      | $6.82 \pm 0.18$  | 8.5          |
| 2014                        |             |                  |              |
| $1023.122 \pm 0.010$        | 977.40      | $2.38 \pm 0.17$  | 6.1          |
| $1113.15 \pm 0.008$         | 898.35      | $3.53 \pm 0.17$  | 8.7          |
| $1158.561 \pm 0.014$        | 863.14      | $2.33 \pm 0.17$  | 6.0          |
| $1161.856 \pm 0.008$        | 860.69      | $3.33 \pm 0.17$  | 8.6          |
| $1171.782 \pm 0.010$        | 853.40      | $2.96 \pm 0.17$  | 8.0          |
| $1172.914 \pm 0.011$        | 852.58      | $2.20 \pm 0.17$  | 5.7          |
| $1221.633 \pm 0.004$        | 818.58      | $5.19 \pm 0.17$  | 13.6         |
| $1230.125 \pm 0.008$        | 812.93      | $3.11 \pm 0.17$  | 8.1          |
| $1234.599 \pm 0.003$        | 809.98      | $10.08 \pm 0.17$ | 26.4         |
| $1235.145 \pm 0.001$        | 809.62      | $8.63 \pm 0.17$  | 48.7         |
| $1236.228 \pm 0.004$        | 808.96      | $5.96 \pm 0.17$  | 15.6         |
| $1237.719 \pm 0.001$        | 807.94      | $9.60 \pm 0.17$  | 25.0         |
| $1248.182 \pm 0.01$         | 801.17      | $2.79 \pm 0.17$  | 7.3          |
| $1299.081 \pm 0.003$        | 769.77      | $10.58 \pm 0.17$ | 27.6         |
| $1311.786 \pm 0.008$        | 762.32      | $3.29 \pm 0.17$  | 8.5          |

Table 3—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $1312.442 \pm 0.008$        | 761.94      | $3.10 \pm 0.17$  | 8.3          |
| $1361.767 \pm 0.018$        | 734.34      | $1.61 \pm 0.17$  | 4.1          |
| $1371.106 \pm 0.014$        | 729.34      | $1.93 \pm 0.17$  | 5.0          |
| $1429.540 \pm 0.009$        | 699.53      | $2.97 \pm 0.17$  | 7.5          |
| $1511.465 \pm 0.006$        | 661.61      | $4.25 \pm 0.17$  | 10.2         |
| $1520.357 \pm 0.018$        | 657.74      | $1.53 \pm 0.17$  | 3.7          |
| $1633.117 \pm 0.023$        | 612.33      | $1.19 \pm 0.17$  | 2.7          |
| $1735.955 \pm 0.001$        | 576.05      | $18.4 \pm 0.17$  | 43.1         |
| $1739.822 \pm 0.019$        | 574.77      | $1.47 \pm 0.17$  | 3.5          |
| $1740.357 \pm 0.012$        | 574.59      | $2.22 \pm 0.17$  | 5.3          |
| $1741.406 \pm 0.019$        | 574.25      | $1.3 \pm 0.17$   | 3.3          |
| $2003.559 \pm 0.009$        | 499.11      | $1.9 \pm 0.17$   | 4.4          |
| $2008.717 \pm 0.018$        | 497.83      | $1.53 \pm 0.17$  | 3.5          |
| $2150.280 \pm 0.010$        | 465.06      | $2.78 \pm 0.17$  | 6.7          |
| $2154.315 \pm 0.012$        | 464.18      | $2.55 \pm 0.17$  | 6.1          |
| $2158.634 \pm 0.002$        | 463.26      | $14.41 \pm 0.17$ | 34.8         |
| $2358.801 \pm 0.010$        | 423.94      | $6.97 \pm 0.17$  | 18.2         |
| $2362.782 \pm 0.005$        | 423.23      | $5.86 \pm 0.17$  | 15.3         |
| $2366.854 \pm 0.003$        | 422.50      | $8.24 \pm 0.17$  | 21.4         |
| 2015                        |             |                  |              |
| $1172.340 \pm 0.024$        | 852.99      | $7.92 \pm 0.40$  | 9.1          |
| $1235.760 \pm 0.006$        | 809.22      | $29.01 \pm 0.33$ | 35.4         |
| $1299.269 \pm 0.021$        | 769.66      | $8.94 \pm 0.63$  | 10.6         |
| $1736.429 \pm 0.008$        | 575.89      | $21.68 \pm 0.33$ | 26.7         |
| $2150.059 \pm 0.060$        | 465.10      | $3.004 \pm 0.70$ | 4            |
| $2154.861 \pm 0.090$        | 464.07      | $2.06 \pm 0.73$  | 4            |
| $2158.634 \pm 0.011$        | 463.26      | $17.3 \pm 0.3$   | 22.9         |
| $2358.722 \pm 0.034$        | 423.96      | $5.501 \pm 0.7$  | 8.1          |



Table 3—Continued

| Frequency<br>$\mu\text{Hz}$ | Period<br>s | Amplitude<br>mma | Signal/Noise |
|-----------------------------|-------------|------------------|--------------|
| $2362.884 \pm 0.023$        | 423.21      | $8.48 \pm 0.70$  | 12.45        |
| $2366.742 \pm 0.024$        | 422.52      | $8.11 \pm 0.7$   | 11.9         |

Table 4. List of 15 Periods Used in Fitting GD358 and Corresponding Best Fit Periods.

| $k$ | Frequency<br>( $\mu\text{Hz}$ ) | Period<br>(s) | Uncertainty<br>(s)    | Best fit periods<br>(s) |
|-----|---------------------------------|---------------|-----------------------|-------------------------|
| 8   | 2363.318                        | 423.13        | 0.04                  | 421.08                  |
| 9   | 2155.544                        | 463.92        | 0.04                  | 458.11                  |
| 10  | 2007.628                        | 497.83        | 0.2                   | 498.16                  |
| 11  | 1857.716                        | 538.30        | 0.3                   | 535.44                  |
| 12  | 1741.505                        | 574.22        | 0.1                   | 583.28                  |
| 13  | 1619.700                        | 617.40        | 0.2                   | 616.91                  |
| 14  | 1518.160                        | 658.69        | 0.5                   | 658.87                  |
| 15  | 1428.943                        | 699.82        | 0.2                   | 701.53                  |
| 16  | 1369.336                        | 730.28        | 0.8                   | 735.98                  |
| 17  | 1299.147                        | 769.74        | 0.2                   | 771.93                  |
| 18  | 1238.129                        | 807.67        | 0.2                   | 806.54                  |
| 19  | 1170.207                        | 854.55        | 0.6                   | 858.64                  |
| 20  | 1109.271                        | 901.49        | 0.8                   | 898.45                  |
| 22  | 1032.967                        | 968.09        | 1.4                   | 970.54                  |
| 24  | 941.333                         | 1062.32       | 3.1                   | 1054.75                 |
|     |                                 |               | $\sigma_{\text{rms}}$ | 4.22 s                  |

Table 5. Master Grid and Best Fit Parameters

|                     | $T_{\text{eff}}$<br>(K) | Mass<br>$M_{\odot}$ | $M_{\text{env}}$<br>$\log(M_*)$ | $M_{\text{He}}$<br>$\log(M_*)$ | $X_o$<br>$\log(M_*)$ | $q_{\text{fm}}$<br>$\log(M_*)$ |        |
|---------------------|-------------------------|---------------------|---------------------------------|--------------------------------|----------------------|--------------------------------|--------|
| Minimum             | 21,000                  | 0.500               | 2.00                            | 4.00                           | 0.10                 | 0.10                           |        |
| Maximum             | 30,000                  | 0.700               | 3.40                            | 7.00                           | 1.00                 | 0.80                           |        |
| Step size           | 200                     | 0.010               | 0.20                            | 0.20                           | 0.10                 | 0.05                           |        |
| Best fit parameters |                         |                     |                                 |                                |                      |                                |        |
| Initial fit         | 26,600                  | 0.520               | −2.40                           | −4.90                          | 0.40                 | 0.16                           | 4.45 s |
| Refined fit         | 25,560                  | 0.5438              | −2.80 (fixed)                   | −5.40                          | 0.40                 | 0.18                           |        |