

# Classification of *P*-oligomorphic permutation groups Conjectures of Cameron and Macpherson

Justine Falque  
joint work with Nicolas M. Thiéry

Laboratoire de Recherche en Informatique  
Université Paris-Sud (Orsay)

SLC, April 17h of 2019

# Profile of a permutation group, a finite example

- Permutation group  $G$

# Profile of a permutation group, a finite example

- Permutation group  $G$   $\rightarrow$  induced action on *subsets*

## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$

# Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$

## Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

# Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$

## Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

## Example

The group of symmetries of the cube  $G_3$

## Profile of a permutation group, a finite example


- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$

### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0			5	
1			6	
2			7	
3			8	
4			$> 8$	





## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1			6	
2			7	
3			8	
4			$> 8$	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1			6	
2			7	
3			8	
4			$> 8$	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2			7	
3			8	
4			$> 8$	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2			7	
3			8	
4			$> 8$	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2			7	
3			8	
4			$> 8$	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2			7	
3			8	
4			$> 8$	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2			7	
3			8	
4			$> 8$	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2	3		7	
3			8	
4			> 8	





## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2	3		7	
3			8	
4			> 8	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2	3		7	
3			8	
4			$> 8$	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2	3		7	
3			8	
4			> 8	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2	3		7	
3	3		8	
4			> 8	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	
1	1		6	
2	3		7	
3	3		8	
4	6		$> 8$	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	3
1	1		6	
2	3		7	
3	3		8	
4	6		$> 8$	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	3
1	1		6	3
2	3		7	
3	3		8	
4	6		> 8	



## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	3
1	1		6	3
2	3		7	1
3	3		8	
4	6		$> 8$	





## Profile of a permutation group, a finite example

- Permutation group  $G \rightarrow$  induced action on *subsets*
- Orbit of  $n$ -subsets = orbit of *degree*  $n$


### Profile of $G$

$$\varphi_G(n) = \# \text{ orbits of degree } n$$

### Example

The group of symmetries of the cube  $G_3$

$n$	$\varphi_G$		$n$	$\varphi_G$
0	1		5	3
1	1		6	3
2	3		7	1
3	3		8	1
4	6		$> 8$	0



## Series of the profile

$$P_{G_3}(z) = 1 + 1z + 3z^2 + 3z^3 + 6z^4 + 3z^5 + 3z^6 + 1z^7 + 1z^8$$

## Series of the profile

$$P_{G_3}(z) = 1 + 1z + 3z^2 + 3z^3 + 6z^4 + 3z^5 + 3z^6 + 1z^7 + 1z^8$$

$$G \text{ infinite} \quad \rightarrow \quad \mathcal{H}_G(z) = \sum_n \varphi_G(n) z^n$$

## Series of the profile

$$P_{G_3}(z) = 1 + 1z + 3z^2 + 3z^3 + 6z^4 + 3z^5 + 3z^6 + 1z^7 + 1z^8$$

$$G \text{ infinite} \quad \rightarrow \quad \mathcal{H}_G(z) = \sum_n \varphi_G(n) z^n$$

## Hypothesis

$G$  is  **$P$ -oligomorphic**:  $\varphi_G$  is bounded by a polynomial in  $n$

## Series of the profile

$$P_{G_3}(z) = 1 + 1z + 3z^2 + 3z^3 + 6z^4 + 3z^5 + 3z^6 + 1z^7 + 1z^8$$

$$G \text{ infinite} \quad \rightarrow \quad \mathcal{H}_G(z) = \sum_n \varphi_G(n) z^n$$

## Hypothesis

$G$  is  **$P$ -oligomorphic**:  $\varphi_G$  is bounded by a polynomial in  $n$

## Example

$$\mathcal{H}_{\mathfrak{S}_\infty}(z) = 1 + z + z^2 + \cdots = \frac{1}{1-z}$$

## Series of the profile

$$P_{G_3}(z) = 1 + 1z + 3z^2 + 3z^3 + 6z^4 + 3z^5 + 3z^6 + 1z^7 + 1z^8$$

$$G \text{ infinite} \quad \rightarrow \quad \mathcal{H}_G(z) = \sum_n \varphi_G(n) z^n$$

## Hypothesis

$G$  is  **$P$ -oligomorphic**:  $\varphi_G$  is bounded by a polynomial in  $n$

## Example

$$\mathcal{H}_{\mathfrak{S}_\infty}(z) = 1 + z + z^2 + \cdots = \frac{1}{1-z}$$

## Conjecture 1 - Cameron, 70's

$$G \text{ } P\text{-oligomorphic} \quad \Rightarrow \quad \mathcal{H}_G(z) = \frac{N(z)}{\prod_i (1-z^{d_i})} \text{ with } N(z) \in \mathbb{Z}[z]$$

# Orbit algebra

Orbit algebra (Cameron, 80's)

Structure of graded algebra  $\mathcal{A}_G = \bigoplus_n \mathcal{A}_n$  on the orbits

# Orbit algebra

## Orbit algebra (Cameron, 80's)

Structure of graded algebra  $\mathcal{A}_G = \bigoplus_n \mathcal{A}_n$  on the orbits

- combinatorial description of the product



# Orbit algebra

## Orbit algebra (Cameron, 80's)

Structure of graded algebra  $\mathcal{A}_G = \bigoplus_n \mathcal{A}_n$  on the orbits

- combinatorial description of the product
- graded according to the orbital degree:

# Orbit algebra

## Orbit algebra (Cameron, 80's)

Structure of graded algebra  $\mathcal{A}_G = \bigoplus_n \mathcal{A}_n$  on the orbits

- combinatorial description of the product
- graded according to the orbital degree:  
 $orb_1 \cdot orb_2 = \text{linear combination of orbits of degree } d_1 + d_2$

# Orbit algebra

## Orbit algebra (Cameron, 80's)

Structure of graded algebra  $\mathcal{A}_G = \bigoplus_n \mathcal{A}_n$  on the orbits

- combinatorial description of the product
- graded according to the orbital degree:  
 $orb_1 \cdot orb_2 = \text{linear combination of orbits of degree } d_1 + d_2$
- $\dim(\mathcal{A}_n) = \varphi_G(n)$

# Orbit algebra

## Orbit algebra (Cameron, 80's)

Structure of graded algebra  $\mathcal{A}_G = \bigoplus_n \mathcal{A}_n$  on the orbits

- combinatorial description of the product
- graded according to the orbital degree:  
 $orb_1 \cdot orb_2 = \text{linear combination of orbits of degree } d_1 + d_2$
- $\dim(\mathcal{A}_n) = \varphi_G(n)$ , so  $\mathcal{H}_G(z) = \sum_n \text{dim}(\mathcal{A}_n) z^n$

# Orbit algebra

## Orbit algebra (Cameron, 80's)

Structure of graded algebra  $\mathcal{A}_G = \bigoplus_n \mathcal{A}_n$  on the orbits

- combinatorial description of the product
- graded according to the orbital degree:  
 $orb_1 \cdot orb_2 = \text{linear combination of orbits of degree } d_1 + d_2$
- $\dim(\mathcal{A}_n) = \varphi_G(n)$ , so  $\mathcal{H}_G(z) = \sum_n \text{dim}(\mathcal{A}_n) z^n$

Conjecture 2 (stronger) - Macpherson, 85

$G$   $P$ -oligomorphic  $\Rightarrow \mathcal{A}_G$  is finitely generated

# Block systems

## Block system

- Equivalence relation preserved by the group

# Block systems

## Block system

- Equivalence relation preserved by the group
- Blocks = the classes

# Block systems

## Block system

- Equivalence relation preserved by the group
- Blocks = the classes
- $G$  acts by permutation on the blocks




# Block systems

## Block system

- Equivalence relation preserved by the group
- Blocks = the classes
- $G$  acts by permutation on the blocks

## Example

Block systems of  $C_4$




# Block systems

## Block system

- Equivalence relation preserved by the group
- Blocks = the classes
- $G$  acts by permutation on the blocks

## Example

Block systems of  $C_4$




# Block systems

## Block system

- Equivalence relation preserved by the group
- Blocks = the classes
- $G$  acts by permutation on the blocks

## Example

Block systems of  $C_4$




# Block systems

## Block system

- Equivalence relation preserved by the group
- Blocks = the classes
- $G$  acts by permutation on the blocks

## Example

Block systems of  $C_4$



# Block systems

## Block system

- Equivalence relation preserved by the group
- Blocks = the classes
- $G$  acts by permutation on the blocks

## Example

Block systems of  $C_4$



Not a block system  $\rightarrow$



# The complete primitive $P$ -oligomorphic groups



$G_\infty$

# The complete primitive $P$ -oligomorphic groups

Macpherson:

$G$   $P$ -oligomorphic with no non trivial blocks  $\Rightarrow \varphi_G(n) = 1$



$\mathfrak{S}_\infty$

# The complete primitive $P$ -oligomorphic groups

Macpherson:

$G$   $P$ -oligomorphic with no non trivial blocks  $\Rightarrow \varphi_G(n) = 1$



Theorem (Classification, Cameron)

Only 5 complete groups such that  $\varphi_G(n) = 1 \quad \forall n$



# The complete primitive $P$ -oligomorphic groups

Macpherson:

$G$   $P$ -oligomorphic with no non trivial blocks  $\Rightarrow \varphi_G(n) = 1$



## Theorem (Classification, Cameron)

Only 5 complete groups such that  $\varphi_G(n) = 1 \quad \forall n$

- $\text{Aut}(\mathbb{Q})$  : automorphisms of the rational chain
- $\text{Rev}(\mathbb{Q})$  : generated by  $\text{Aut}(\mathbb{Q})$  and one reflection
- $\text{Aut}(\mathbb{Q}/\mathbb{Z})$ , preserving the circular order
- $\text{Rev}(\mathbb{Q}/\mathbb{Z})$  : generated by  $\text{Aut}(\mathbb{Q}/\mathbb{Z})$  and a reflection
- $\mathfrak{S}_\infty$  : the symmetric group

# The complete primitive $P$ -oligomorphic groups

Macpherson:

$G$   $P$ -oligomorphic with no non trivial blocks  $\Rightarrow \varphi_G(n) = 1$



$\mathfrak{S}_\infty$

## Theorem (Classification, Cameron)

Only 5 complete groups such that  $\varphi_G(n) = 1 \quad \forall n$


- $\text{Aut}(\mathbb{Q})$  : automorphisms of the rational chain
- $\text{Rev}(\mathbb{Q})$  : generated by  $\text{Aut}(\mathbb{Q})$  and one reflection
- $\text{Aut}(\mathbb{Q}/\mathbb{Z})$ , preserving the circular order
- $\text{Rev}(\mathbb{Q}/\mathbb{Z})$  : generated by  $\text{Aut}(\mathbb{Q}/\mathbb{Z})$  and a reflection
- $\mathfrak{S}_\infty$  : the symmetric group

Well known, nice groups (called *highly homogeneous*).  
In particular, their orbit algebra is finitely generated.


# An infinite example: $\mathfrak{S}_\infty \wr \mathfrak{S}_3$




An infinite example:  $\mathfrak{S}_\infty \wr \mathfrak{S}_3$



# An infinite example: $\mathfrak{S}_\infty \wr \mathfrak{S}_3$




# An infinite example: $\mathfrak{S}_\infty \wr \mathfrak{S}_3$



## An infinite example: $\mathfrak{S}_\infty \wr \mathfrak{S}_3$

Wreath product


$$\mathfrak{S}_\infty \wr \mathfrak{S}_3$$



## An infinite example: $\mathfrak{S}_\infty \wr \mathfrak{S}_3$


Wreath product

$$\mathfrak{S}_\infty \wr \mathfrak{S}_3 \simeq \mathfrak{S}_\infty^3 \rtimes \mathfrak{S}_3$$





## An infinite example: $\mathfrak{S}_\infty \wr \mathfrak{S}_3$




Wreath product

$$\mathfrak{S}_\infty \wr \mathfrak{S}_3 \simeq \mathfrak{S}_\infty^3 \rtimes \mathfrak{S}_3$$

Subset of shape  $2, 3, 2 \rightarrow x_1^2 x_2^3 x_3^2$

# An infinite example: $\mathfrak{S}_\infty \wr \mathfrak{S}_3$



## Wreath product


$$\mathfrak{S}_\infty \wr \mathfrak{S}_3 \simeq \mathfrak{S}_\infty^3 \rtimes \mathfrak{S}_3$$

Subset of shape  $2, 3, 2 \rightarrow x_1^2 x_2^3 x_3^2$

Orbits of subsets

$\leftrightarrow$  symmetric polynomials in  $x_1, x_2, x_3$

# An infinite example: $\mathfrak{S}_\infty \wr \mathfrak{S}_3$



## Wreath product

$$\mathfrak{S}_\infty \wr \mathfrak{S}_3 \simeq \mathfrak{S}_\infty^3 \rtimes \mathfrak{S}_3$$


Subset of shape  $\mathbf{2}, \mathbf{3}, \mathbf{2} \rightarrow x_1^2 x_2^3 x_3^2$

Orbits of subsets

$\leftrightarrow$  symmetric polynomials in  $x_1, x_2, x_3$

$$\mathcal{A}_{\mathfrak{S}_\infty \wr \mathfrak{S}_3} \simeq \text{Sym}_3[X] = \mathbb{Q}[X]^{\mathfrak{S}_3}$$

# An infinite example: $\mathfrak{S}_\infty \wr \mathfrak{S}_3$



## Wreath product

$$\mathfrak{S}_\infty \wr \mathfrak{S}_3 \simeq \mathfrak{S}_\infty^3 \rtimes \mathfrak{S}_3$$

Subset of shape  $2, 3, 2 \rightarrow x_1^2 x_2^3 x_3^2$


Orbits of subsets

$\leftrightarrow$  symmetric polynomials in  $x_1, x_2, x_3$


$$\mathcal{A}_{\mathfrak{S}_\infty \wr \mathfrak{S}_3} \simeq \text{Sym}_3[X] = \mathbb{Q}[X]^{\mathfrak{S}_3}$$

One can obtain functions counting integer partitions, combinations,  $P$ -partitions (with optional length and/or height restrictions) as profiles of wreath products...

## Lower bound




# Lower bound



$$\Rightarrow G \leq \mathfrak{S}_M \wr \mathfrak{S}_N$$


# Lower bound



$$\Rightarrow G \leq \mathfrak{S}_M \wr \mathfrak{S}_N$$

Two cases if  $G$  is  $P$ -oligomorphic :

# Lower bound




$$\Rightarrow G \leq \mathfrak{S}_M \wr \mathfrak{S}_N$$

Two cases if  $G$  is  $P$ -oligomorphic :



## Lower bound




$$\Rightarrow G \leq \mathfrak{S}_M \wr \mathfrak{S}_N$$

Two cases if  $G$  is  $P$ -oligomorphic :

- $M < \infty$

# Lower bound




$$\Rightarrow G \leq \mathfrak{S}_M \wr \mathfrak{S}_N$$

Two cases if  $G$  is  $P$ -oligomorphic :

- $M < \infty$
- $N < \infty$


## Lower bound



Two cases if  $G$  is  $P$ -oligomorphic :

- $M < \infty \quad \longrightarrow \quad \varphi_G(n) \geq O(n^{M-1})$
- $N < \infty$


# Lower bound



Two cases if  $G$  is  $P$ -oligomorphic :

- $M < \infty \quad \longrightarrow \quad \varphi_G(n) \geq O(n^{M-1})$
- $N < \infty \quad \longrightarrow \quad \varphi_G(n) \geq O(n^{N-1})$

## Lower bound



Two cases if  $G$  is  $P$ -oligomorphic :

- $M < \infty \quad \longrightarrow \quad \varphi_G(n) \geq O(n^{M-1})$
- $N < \infty \quad \longrightarrow \quad \varphi_G(n) \geq O(n^{N-1})$


Better have **big** finite blocks and/or "small" infinite ones...

# Lattices of block systems

Lattice of partitions  $\rightarrow$  structure of *lattice* on block systems


# Lattices of block systems

Lattice of partitions  $\rightarrow$  structure of *lattice* on block systems



# Lattices of block systems

Lattice of partitions  $\rightarrow$  structure of *lattice* on block systems




Non trivial fact

- $\{\text{Systems with } < \infty \text{ blocks only}\} = \text{sublattice with maximum}$
- $\{\text{Systems with } \infty \text{ blocks only}\} = \text{sublattice with minimum}$



# Lattices of block systems

Lattice of partitions  $\rightarrow$  structure of *lattice* on block systems




Non trivial fact

- $\{\text{Systems with } < \infty \text{ blocks only}\} = \text{sublattice with maximum}$
- $\{\text{Systems with } \infty \text{ blocks only}\} = \text{sublattice with minimum}$

**Remark.** If  $G$  is  $P$ -oligomorphic, both of them are actually finite!

# The nested block system


## Idea



# The nested block system

## Idea


1. Take the *maximal* system of finite blocks



# The nested block system

## Idea


1. Take the *maximal* system of finite blocks



# The nested block system

## Idea

1. Take the *maximal* system of finite blocks




Action on the maximal finite blocks...

## The nested block system

### Idea

1. Take the *maximal* system of finite blocks




Action on the maximal finite blocks...

that has no finite blocks.

# The nested block system

## Idea

1. Take the *maximal* system of finite blocks
2. Take the *minimal* system of infinite blocks of the action of  $G$  on the maximal finite blocks




Action on the maximal finite blocks...

that has no finite blocks.

# The nested block system

## Idea

1. Take the *maximal* system of finite blocks
2. Take the *minimal* system of infinite blocks of the action of  $G$  on the maximal finite blocks  $\rightarrow$  finitely many **superblocks**




Action on the maximal finite blocks... that has no finite blocks.



# The nested block system


## Idea

1. Take the *maximal* system of finite blocks
2. Take the *minimal* system of infinite blocks of the action of  $G$  on the maximal finite blocks  $\rightarrow$  finitely many **superblocks**




Action on the maximal finite blocks... that has no finite blocks.


## One superblock: examples




## One superblock: examples




## One superblock: examples




## One superblock: examples




## One superblock: examples



## One superblock: examples




## One superblock: examples



$$G|_{B_0} = H_0$$




# One superblock: examples




$$G|_{B_0} = H_0, \text{ Fix}(B_0)$$

# One superblock: examples




$$G|_{B_0} = H_0, \quad \text{Fix}(B_0)|_{B_1} = H_1$$


## One superblock: examples




## One superblock: examples




## One superblock: examples




## One superblock: examples



## One superblock: examples




## One superblock: examples






## One superblock: examples



- $H \wr G_\infty \rightarrow H, H, H, H, H, H, \dots$


## One superblock: examples



$H_0, H_1, H_2, H_3, H_4, H_5 \dots$  **Tower of  $G$**

- $H \wr \mathfrak{S}_\infty \rightarrow H, H, H, H, H, H \dots$
- " $H_0 \times \mathfrak{S}_\infty$ "  $\rightarrow H_0, \text{Id}, \text{Id}, \text{Id}, \text{Id}, \text{Id} \dots$


## One superblock: examples



$H_0, H_1, H_2, H_3, H_4, H_5 \dots$  **Tower of  $G$**

- $H \wr \mathfrak{S}_\infty \rightarrow H, H, H, H, H, H \dots$
- " $H_0 \times \mathfrak{S}_\infty$ "  $\rightarrow H_0, \text{Id}, \text{Id}, \text{Id}, \text{Id}, \text{Id} \dots$
- $\langle "H_0 \times \mathfrak{S}_\infty", H \wr \mathfrak{S}_\infty \rangle$

## One superblock: examples




$H_0, H_1, H_2, H_3, H_4, H_5 \dots$  **Tower of  $G$**

- $H \wr \mathfrak{S}_\infty \rightarrow H, H, H, H, H, H \dots$
- " $H_0 \times \mathfrak{S}_\infty$ "  $\rightarrow H_0, \text{Id}, \text{Id}, \text{Id}, \text{Id}, \text{Id} \dots$
- $\langle "H_0 \times \mathfrak{S}_\infty", H \wr \mathfrak{S}_\infty \rangle$

$H_0 \triangleright H$  w.l.o.g

## One superblock: examples



$H_0, H_1, H_2, H_3, H_4, H_5 \dots$  **Tower of  $G$**

- $H \wr \mathfrak{S}_\infty \rightarrow H, H, H, H, H, H \dots$
- " $H_0 \times \mathfrak{S}_\infty$ "  $\rightarrow H_0, \text{Id}, \text{Id}, \text{Id}, \text{Id}, \text{Id} \dots$
- $\langle "H_0 \times \mathfrak{S}_\infty", H \wr \mathfrak{S}_\infty \rangle \rightarrow H_0, H, H, H, H, H \dots$

$H_0 \triangleright H$  w.l.o.g

Notation:  $[H_0, H_\infty]$

## One superblock: classification

- The tower determines  $G$  (uses the *subdirect product*)

## One superblock: classification

- The tower determines  $G$  (uses the *subdirect product*)
- Computer exploration on finitely many blocks

## One superblock: classification

- The tower determines  $G$  (uses the *subdirect product*)
- Computer exploration on finitely many blocks
  - Observation: always some  $H_0, H, H, H, \dots, H, H_s$



## One superblock: classification

- The tower determines  $G$  (uses the *subdirect product*)
- Computer exploration on finitely many blocks
  - Observation: always some  $H_0, H, H, H, \dots, H, H_s$
  - Proof in the infinite case: always some  $H_0, H, H, H \dots$

## One superblock: classification

- The tower determines  $G$  (uses the *subdirect product*)
- Computer exploration on finitely many blocks
  - Observation: always some  $H_0, H, H, H, \dots, H, H_s$
  - Proof in the infinite case: always some  $H_0, H, H, H \dots$

## Classification

One superblock  $\Rightarrow G = [H_0, H_\infty]$

## One superblock: classification

- The tower determines  $G$  (uses the *subdirect product*)
- Computer exploration on finitely many blocks
  - Observation: always some  $H_0, H, H, H, \dots, H, H_s$
  - Proof in the infinite case: always some  $H_0, H, H, H \dots$

### Classification

One superblock  $\Rightarrow G = [H_0, H_\infty]$

$\mathbb{Q}[(X_{orb})_{orb}]$  , where  $orb$  runs through the orbits of  $H$

## One superblock: classification

- The tower determines  $G$  (uses the *subdirect product*)
- Computer exploration on finitely many blocks
  - Observation: always some  $H_0, H, H, H, \dots, H, H_s$
  - Proof in the infinite case: always some  $H_0, H, H, H \dots$

### Classification

One superblock  $\Rightarrow G = [H_0, H_\infty]$

$\mathcal{A}_G \simeq \mathbb{Q}[(X_{orb})_{orb}]^{H_0}$ , where  $orb$  runs through the orbits of  $H$

## One superblock: classification

- The tower determines  $G$  (uses the *subdirect product*)
- Computer exploration on finitely many blocks
  - Observation: always some  $H_0, H, H, H, \dots, H, H_s$
  - Proof in the infinite case: always some  $H_0, H, H, H \dots$


### Classification

One superblock  $\Rightarrow G = [H_0, H_\infty]$

$\mathcal{A}_G \simeq \mathbb{Q}[(X_{orb})_{orb}]^{H_0}$ , where  $orb$  runs through the orbits of  $H$


In particular, both conjectures hold.

# General case: minimal subgroup of finite index



# General case: minimal subgroup of finite index


Normal subgroup  $K$  of  $G$



## General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$

- that fixes the kernel






# General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$


- that fixes the kernel



## General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$


- that fixes the kernel



# General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$


- that fixes the kernel
- that stabilizes the superblocks



# General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$


- that fixes the kernel
- that stabilizes the superblocks



## General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$


- that fixes the kernel
- that stabilizes the superblocks
- of restrictions wreath products onto the superblocks



# General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$


- that fixes the kernel
- that stabilizes the superblocks
- of restrictions wreath products onto the superblocks



# General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$


- that fixes the kernel
- that stabilizes the superblocks
- of restrictions wreath products onto the superblocks



# General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$

- that fixes the kernel
- that stabilizes the superblocks
- of restrictions wreath products onto the superblocks
- in which  $\text{Rev}(\dots)$  are reduced down to  $\text{Aut}(\dots)$






# General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$


- that fixes the kernel
- that stabilizes the superblocks
- of restrictions wreath products onto the superblocks
- in which  $\text{Rev}(\dots)$  are reduced down to  $\text{Aut}(\dots)$



# General case: minimal subgroup of finite index

Normal subgroup  $K$  of  $G$

- that fixes the kernel
- that stabilizes the superblocks
- of restrictions wreath products onto the superblocks
- in which  $\text{Rev}(\dots)$  are reduced down to  $\text{Aut}(\dots)$



## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)

## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)  
Linked to the simplicity of  $\mathfrak{S}_\infty$  and  $\text{Aut}(\dots)$ , and  
the wreath products' being "too big, too free"

## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)  
Linked to the simplicity of  $\mathfrak{S}_\infty$  and  $\text{Aut}(\dots)$ , and  
the wreath products' being "too big, too free"  
 $\Rightarrow$  direct product of the restrictions  $K^{(i)}$

## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)

Linked to the simplicity of  $\mathfrak{S}_\infty$  and  $\text{Aut}(\dots)$ , and the wreath products' being "too big, too free"

$\Rightarrow$  direct product of the restrictions  $K^{(i)}$

$\Rightarrow \mathcal{A}_K = \bigotimes_i \mathcal{A}_{K^{(i)}}$

## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)

Linked to the simplicity of  $\mathfrak{S}_\infty$  and  $\text{Aut}(\dots)$ , and the wreath products' being "too big, too free"

$\Rightarrow$  direct product of the restrictions  $K^{(i)}$

$\Rightarrow \mathcal{A}_K = \bigotimes_i \mathcal{A}_{K^{(i)}}$

- $K^{(i)} = H^{(i)} \wr \mathfrak{S}_\infty$  wreath product with finite blocks

## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)  
 Linked to the simplicity of  $\mathfrak{S}_\infty$  and  $\text{Aut}(\dots)$ , and the wreath products' being "too big, too free"  
 $\Rightarrow$  direct product of the restrictions  $K^{(i)}$   
 $\Rightarrow \mathcal{A}_K = \bigotimes_i \mathcal{A}_{K^{(i)}}$
- $K^{(i)} = H^{(i)} \wr \mathfrak{S}_\infty$  wreath product with finite blocks  
 $\Rightarrow \mathcal{A}_K \simeq \bigotimes_i \mathbb{Q}[(X_{orb_i})_{orb_i}]$   
 free algebra finitely generated by the orbits of the  $H^{(i)}$ 's



## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)  
 Linked to the simplicity of  $\mathfrak{S}_\infty$  and  $\text{Aut}(\dots)$ , and  
 the wreath products' being "too big, too free"  
 $\Rightarrow$  direct product of the restrictions  $K^{(i)}$   
 $\Rightarrow \mathcal{A}_K = \bigotimes_i \mathcal{A}_{K^{(i)}}$
- $K^{(i)} = H^{(i)} \wr \mathfrak{S}_\infty$  wreath product with finite blocks  
 $\Rightarrow \mathcal{A}_K \simeq \bigotimes_i \mathbb{Q}[(X_{orb_i})_{orb_i}]$   
 free algebra finitely generated by the orbits of the  $H^{(i)}$ 's  
 (plus some idempotents brought by the kernel)

## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)  
 Linked to the simplicity of  $\mathfrak{S}_\infty$  and  $\text{Aut}(\dots)$ , and the wreath products' being "too big, too free"  
 $\Rightarrow$  direct product of the restrictions  $K^{(i)}$   
 $\Rightarrow \mathcal{A}_K = \bigotimes_i \mathcal{A}_{K^{(i)}}$
- $K^{(i)} = H^{(i)} \wr \mathfrak{S}_\infty$  wreath product with finite blocks  
 $\Rightarrow \mathcal{A}_K \simeq \bigotimes_i \mathbb{Q}[(X_{orb_i})_{orb_i}]$   
 free algebra finitely generated by the orbits of the  $H^{(i)}$ 's  
 (plus some idempotents brought by the kernel)
- Fact:  $G$  acts by permutation on these generators

## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)  
Linked to the simplicity of  $\mathfrak{S}_\infty$  and  $\text{Aut}(\dots)$ , and the wreath products' being "too big, too free"  
 $\Rightarrow$  direct product of the restrictions  $K^{(i)}$   
 $\Rightarrow \mathcal{A}_K = \bigotimes_i \mathcal{A}_{K^{(i)}}$
- $K^{(i)} = H^{(i)} \wr \mathfrak{S}_\infty$  wreath product with finite blocks  
 $\Rightarrow \mathcal{A}_K \simeq \bigotimes_i \mathbb{Q}[(X_{orb_i})_{orb_i}]$   
free algebra finitely generated by the orbits of the  $H^{(i)}$ 's  
(plus some idempotents brought by the kernel)
- Fact:  $G$  acts by permutation on these generators  
 $\Rightarrow \mathcal{A}_G$  is the *algebra of invariants* of this finite action

## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)  
 Linked to the simplicity of  $\mathfrak{S}_\infty$  and  $\text{Aut}(\dots)$ , and the wreath products' being "too big, too free"  
 $\Rightarrow$  direct product of the restrictions  $K^{(i)}$   
 $\Rightarrow \mathcal{A}_K = \bigotimes_i \mathcal{A}_{K^{(i)}}$
- $K^{(i)} = H^{(i)} \wr \mathfrak{S}_\infty$  wreath product with finite blocks  
 $\Rightarrow \mathcal{A}_K \simeq \bigotimes_i \mathbb{Q}[(X_{orb_i})_{orb_i}]$   
 free algebra finitely generated by the orbits of the  $H^{(i)}$ 's  
 (plus some idempotents brought by the kernel)
- Fact:  $G$  acts by permutation on these generators  
 $\Rightarrow \mathcal{A}_G$  is the *algebra of invariants* of this finite action
- Hilbert's theorem:  
 $\mathcal{A}_G$  finitely generated (and even *Cohen-Macaulay*)


## Shape of the orbit algebra $\mathcal{A}_G$

- In  $K$ , totally independent superblocks (and kernel)  
 Linked to the simplicity of  $\mathfrak{S}_\infty$  and  $\text{Aut}(\dots)$ , and the wreath products' being "too big, too free"  
 $\Rightarrow$  direct product of the restrictions  $K^{(i)}$   
 $\Rightarrow \mathcal{A}_K = \bigotimes_i \mathcal{A}_{K^{(i)}}$
- $K^{(i)} = H^{(i)} \wr \mathfrak{S}_\infty$  wreath product with finite blocks  
 $\Rightarrow \mathcal{A}_K \simeq \bigotimes_i \mathbb{Q}[(X_{orb_i})_{orb_i}]$   
 free algebra finitely generated by the orbits of the  $H^{(i)}$ 's  
 (plus some idempotents brought by the kernel)
- Fact:  $G$  acts by permutation on these generators  
 $\Rightarrow \mathcal{A}_G$  is the *algebra of invariants* of this finite action
- Hilbert's theorem:  
 $\mathcal{A}_G$  finitely generated (and even *Cohen-Macaulay*)

Which end the proof of the conjectures!


# Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group



# Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group,  $\mathcal{B}_0$  a block system.



# Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group,  $\mathcal{B}_0$  a block system.

For each orbit of blocks






# Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group,  $\mathcal{B}_0$  a block system.

For each orbit of blocks, choose

1. One group of profile 1




# Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group,  $\mathcal{B}_0$  a block system.

For each orbit of blocks, choose

1. One group of profile 1
  - Has to be  $\mathfrak{S}_\infty$  if the blocks are singletons




# Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group,  $\mathcal{B}_0$  a block system.

For each orbit of blocks, choose

1. One group of profile 1

- Has to be  $\mathfrak{S}_\infty$  if the blocks are singletons
- Can alternatively be  $\text{Id}_1$  for at most one orbit of one block




# Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group,  $\mathcal{B}_0$  a block system.

For each orbit of blocks, choose

## 1. One group of profile 1

- Has to be  $\mathfrak{S}_\infty$  if the blocks are singletons
- Can alternatively be  $\text{Id}_1$  for at most one orbit of one block




# Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group,  $\mathcal{B}_0$  a block system.

For each orbit of blocks, choose

## 1. One group of profile 1

- Has to be  $\mathfrak{S}_\infty$  if the blocks are singletons
- Can alternatively be  $\text{Id}_1$  for at most one orbit of one block




## Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group,  $\mathcal{B}_0$  a block system.

For each orbit of blocks, choose

1. One **group of profile 1**

- Has to be  $\mathfrak{S}_\infty$  if the blocks are singletons
- Can alternatively be  $\text{Id}_1$  for at most one orbit of one block




# Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group,  $\mathcal{B}_0$  a block system.

For each orbit of blocks, choose

1. One **group of profile 1**
  - Has to be  $\mathfrak{S}_\infty$  if the blocks are singletons
  - Can alternatively be  $\text{Id}_1$  for at most one orbit of one block
2. One normal subgroup  $H$  of  $H_0 = G_{0|B}$  for  $B$  in the orbit


$H' \wr \mathfrak{S}_\infty$



$H \wr \mathfrak{S}_\infty$



$H \wr \mathfrak{S}_\infty$



$\text{Id} \wr \text{Aut}(\mathbb{Q})$




# Classification of $P$ -oligomorphic groups

$G_0$  a finite permutation group,  $\mathcal{B}_0$  a block system.

For each orbit of blocks, choose

1. One **group of profile 1**
  - Has to be  $\mathfrak{S}_\infty$  if the blocks are singletons
  - Can alternatively be  $\text{Id}_1$  for at most one orbit of one block
2. One normal subgroup  $H$  of  $H_0 = G_{0|B}$  for  $B$  in the orbit





# Thank you for your attention !

## Context



- $G$  permutation group of a countably infinite set  $E$
- Profile  $\varphi_G$ : counts the orbits of finite subsets of  $E$
- Hypothesis:  $\varphi_G(n)$  bounded by a polynomial
- Conjecture (Cameron): rational form of the generating series
- Conjecture (Macpherson): finite generation of the orbit algebra

## Results

- Both conjectures hold !
- Classification of  $P$ -oligomorphic permutation groups
- The orbit algebra is an algebra of invariants (up to some idempotents)

# The tower determines the group (1): "straight $\mathfrak{S}_\infty$ "

$G$  contains a set of "straight" swaps of blocks



## Subdirect product

### Subdirect product of $G_1$ and $G_2$

- Formalizes the *synchronization* between  $G_1$  and  $G_2$
- Subgroup of  $G_1 \times G_2$  (with canonical projections  $G_1$  and  $G_2$ )
- $E = E_1 \sqcup E_2$  stable  $\Rightarrow G$  subdirect product of  $G|_{E_1}$  and  $G|_{E_2}$

### Synchronization in a subdirect product

Let  $N_1 = \text{Fix}_G(E_2)$  and  $N_2 = \text{Fix}_G(E_1)$ .



$$\frac{G_1}{N_1} \simeq \frac{G}{N_1 \times N_2} \simeq \frac{G_2}{N_2}$$

A subdirect product with explicit  $N_i$ 's is explicit.

**Remark.**  $N_1$  and  $N_2$  are *normal* in  $G_1$  and  $G_2$ , so the possibilities of synchronization of a group is linked to its normal subgroups.

## The tower determines the group (2): $\text{Stab}_G(\text{blocks})$

$\text{Stab}_G(\text{blocks}) = \text{explicit subdirect product of the } H_i$




← The tower  
determines  
 $\text{Stab}_G(\text{blocks})$


$$G \simeq \text{Stab}_G(\text{blocks}) \rtimes \text{"straight } \mathfrak{S}_\infty \text{"} \rightarrow \text{Ok}$$

Example of a product in the cyclic group  $\mathcal{C}_5$


## Example of a product in the cyclic group $\mathcal{C}_5$



# Example of a product in the cyclic group $\mathcal{C}_5$




## Example of a product in the cyclic group $C_5$







## Example of a product in the cyclic group $C_5$




## Example of a product in the cyclic group $\mathcal{C}_5$




## Example of a product in the cyclic group $\mathcal{C}_5$




## Example of a product in the cyclic group $\mathcal{C}_5$



## Example of a product in the cyclic group $\mathcal{C}_5$




## Example of a product in the cyclic group $\mathcal{C}_5$




The diagram illustrates the multiplication of two elements in the Hecke algebra of a Coxeter group. The top row shows a blue element (a 5-cycle with nodes 1, 2, 3, 4, 5) multiplied by a red element (a 5-cycle with nodes 1, 2, 3, 4, 5). The middle row shows the resulting sum of five terms, each a 5-cycle with one node colored red and the others blue. The bottom row shows the final result as a sum of terms, including a zero term, a zero term, and a 5-cycle with nodes 1, 2, 3, 4, 5 where nodes 1, 2, and 3 are blue, and nodes 4 and 5 are red, followed by an ellipsis.

# Example of a product in the cyclic group $\mathcal{C}_5$





## Example of a product in the cyclic group $\mathcal{C}_5$



# Example of a product in the cyclic group $\mathcal{C}_5$

$$\begin{array}{c}
 \begin{array}{c} \text{Diagram 1} \\ \times \\ \text{Diagram 2} \end{array} \Leftrightarrow \begin{array}{c} \text{Diagram 1.1} \\ + \\ \text{Diagram 1.2} \\ + \\ \text{Diagram 1.3} \\ + \\ \text{Diagram 1.4} \\ + \\ \text{Diagram 1.5} \end{array} \\
 \\
 \begin{array}{c} \text{Diagram 2.1} \\ + \\ \text{Diagram 2.2} \\ + \\ \text{Diagram 2.3} \\ + \\ \text{Diagram 2.4} \\ + \\ \text{Diagram 2.5} \end{array} \\
 \\
 \hline
 = \begin{array}{c} 0 \\ + \\ 0 \\ + \\ \text{Diagram 3.1} \\ + \\ \text{Diagram 3.2} \\ + \\ \text{Diagram 3.3} \\ + \\ \text{Diagram 3.4} \\ + \dots \end{array} \\
 \\
 \hline
 = 2 \begin{array}{c} \text{Diagram 4.1} \end{array}
 \end{array}$$

The diagrams are 5-cycles with nodes labeled 1, 2, 3, 4, 5. The first diagram has nodes 1 and 2 colored blue. The second diagram has node 1 colored red. The first row shows the decomposition of the product of these two cycles into five 5-cycles. The second row shows the decomposition of the product of the first row's five 5-cycles into five 5-cycles. The third row shows the final result, which is twice the 5-cycle with nodes 1 and 2 colored purple.

# Example of a product in the cyclic group $\mathcal{C}_5$

$$\begin{array}{c}
 \begin{array}{c} \text{Diagram 1} \\ \times \\ \text{Diagram 2} \end{array} \Leftrightarrow \begin{array}{c} \text{Diagram 1.1} \\ + \\ \text{Diagram 1.2} \\ + \\ \text{Diagram 1.3} \\ + \\ \text{Diagram 1.4} \\ + \\ \text{Diagram 1.5} \end{array} \\
 \\
 \begin{array}{c} \text{Diagram 2.1} \\ + \\ \text{Diagram 2.2} \\ + \\ \text{Diagram 2.3} \\ + \\ \text{Diagram 2.4} \\ + \\ \text{Diagram 2.5} \end{array} \\
 \\
 \hline
 = \begin{array}{c} 0 \\ + \\ 0 \\ + \\ \text{Diagram 3.1} \\ + \\ \text{Diagram 3.2} \\ + \\ \text{Diagram 3.3} \\ + \\ \text{Diagram 3.4} \\ + \dots \end{array} \\
 \\
 \hline
 = \begin{array}{c} 2 \\ \text{Diagram 4.1} \\ + \\ 2 \\ \text{Diagram 4.2} \\ + \dots \end{array}
 \end{array}$$

The diagrams are 5-cycles with nodes labeled 1, 2, 3, 4, 5. The first diagram has nodes 1 and 2 colored blue. The second diagram has node 1 colored red. The first row shows the decomposition of the product of these two cycles into five 5-cycles. The second row shows the decomposition of the product of the first row's five 5-cycles into five 5-cycles. The third row shows the simplification of the second row's five 5-cycles into two 5-cycles, each with a coefficient of 2.


# Example of a product in the cyclic group $\mathcal{C}_5$

$$\begin{array}{c}
 \begin{array}{c} \text{Diagram 1} \\ \times \\ \text{Diagram 2} \end{array} \Leftrightarrow \begin{array}{c} \text{Diagram 1.1} \\ + \\ \text{Diagram 1.2} \\ + \\ \text{Diagram 1.3} \\ + \\ \text{Diagram 1.4} \\ + \\ \text{Diagram 1.5} \end{array} \\
 \\
 \begin{array}{c} \text{Diagram 2.1} \\ + \\ \text{Diagram 2.2} \\ + \\ \text{Diagram 2.3} \\ + \\ \text{Diagram 2.4} \\ + \\ \text{Diagram 2.5} \end{array} \\
 \\
 \hline
 = \begin{array}{c} 0 \\ + \\ 0 \\ + \\ \text{Diagram 3.1} \\ + \\ \text{Diagram 3.2} \\ + \\ \text{Diagram 3.3} \\ + \\ \text{Diagram 3.4} \\ + \dots \end{array} \\
 \\
 \hline
 = \begin{array}{c} 2 \\ \text{Diagram 4.1} \\ + \\ 2 \\ \text{Diagram 4.2} \\ + \dots + 1 \\ \text{Diagram 4.3} \\ + \dots \end{array}
 \end{array}$$

The diagrams are 5-cycles with nodes labeled 1, 2, 3, 4, 5. The first diagram has nodes 1 and 2 blue. The second diagram has node 1 red. The first row shows the decomposition of the product of these two diagrams into five terms. The second row shows the decomposition of the second term into five terms. The third row shows the decomposition of the third term into five terms. The fourth row shows the decomposition of the fourth term into five terms. The fifth row shows the decomposition of the fifth term into five terms. The final result is a sum of terms with coefficients 2, 2, ..., 1.

Example :  $G = \mathfrak{S}_\infty \wr \mathfrak{S}_\infty$


$$\varphi_G(n) = ?$$



Example :  $G = \mathfrak{S}_\infty \wr \mathfrak{S}_\infty$

$$\varphi_G(n) = ?$$


An orbit of degree  $n \longleftrightarrow$  a partition of  $n$



Example :  $G = \mathfrak{S}_\infty \wr \mathfrak{S}_\infty$

$$\varphi_G(n) = ?$$


An orbit of degree  $n \longleftrightarrow$  a partition of  $n$



Example :  $G = \mathfrak{S}_\infty \wr \mathfrak{S}_\infty$

$$\varphi_G(n) = ?$$

An orbit of degree  $n \longleftrightarrow$  a partition of  $n$






Example :  $G = \mathfrak{S}_\infty \wr \mathfrak{S}_\infty$

$\varphi_G(n) =$


An orbit of degree  $n \longleftrightarrow$  a partition of  $n$



Example :  $G = \mathfrak{S}_\infty \wr \mathfrak{S}_\infty$

$$\varphi_G(n) = p(n)$$

An orbit of degree  $n \longleftrightarrow$  a partition of  $n$



# Examples of orbit algebras (1)

## Example 1

If  $G = \mathfrak{S}_\infty$ ,  $\varphi_G(n) = 1$  for all  $n$ , and  $\mathbb{Q}\mathcal{A}(G) = \mathbb{K}[x]$ .

## Examples of orbit algebras (1)

### Example 1

If  $G = \mathfrak{S}_\infty$ ,  $\varphi_G(n) = 1$  for all  $n$ , and  $\mathbb{Q}\mathcal{A}(G) = \mathbb{K}[x]$ .

### Example 2

$G = \mathfrak{S}_\infty \wr \mathfrak{S}_3$ , recall that  $\varphi_G(n) = p_3(n)$ .

## Examples of orbit algebras (1)


### Example 1

If  $G = \mathfrak{S}_\infty$ ,  $\varphi_G(n) = 1$  for all  $n$ , and  $\mathbb{Q}\mathcal{A}(G) = \mathbb{K}[x]$ .

### Example 2

$G = \mathfrak{S}_\infty \wr \mathfrak{S}_3$ , recall that  $\varphi_G(n) = p_3(n)$ .

$A_n$  = homogeneous symmetric polynomials of degree  $n$  in  $x_1, x_2, x_3$



## Examples of orbit algebras (1)


### Example 1

If  $G = \mathfrak{S}_\infty$ ,  $\varphi_G(n) = 1$  for all  $n$ , and  $\mathbb{Q}\mathcal{A}(G) = \mathbb{K}[x]$ .

### Example 2

$G = \mathfrak{S}_\infty \wr \mathfrak{S}_3$ , recall that  $\varphi_G(n) = p_3(n)$ .

$A_n$  = homogeneous symmetric polynomials of degree  $n$  in  $x_1, x_2, x_3$




$$\rightarrow \mathbb{Q}\mathcal{A}(\mathfrak{S}_\infty \wr \mathfrak{S}_3) = \mathbb{K}[x_1, x_2, x_3]^{\mathfrak{S}_3}$$


## Examples of orbit algebras (2)

More generally, for  $H$  subgroup of  $\mathfrak{S}_m$  :

- $G = \mathfrak{S}_\infty \wr H$  :  
 $\mathbb{QA}(G) = \mathbb{K}[x_1, \dots, x_m]^H$ , the algebra of invariants of  $H$   
 $\mathbb{QA}(G)$  is finitely generated by Hilbert's theorem.




- $G = H \wr \mathfrak{S}_\infty$  :  
 $\mathbb{QA}(G)$  = the free algebra generated by the age of  $H$



## Direct product in the case of finite blocks

"Speak, friend..."






# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3




# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3




# Direct product in the case of finite blocks

"Speak, friend..."


## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



"Speak, friend..."


### Example 3

 $C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3

### Direct product in the case of finite blocks

"Speak, friend..."

### Example 3


 $C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3

# Direct product in the case of finite blocks


"Speak, friend..."

## Example 3


$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



"Speak, friend..."

 $C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3

"Speak, friend..."

 $C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3




# Direct product in the case of finite blocks


"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



"Speak, friend..."


 $C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3

# Direct product in the case of finite blocks


"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



"Speak, friend..."


 $C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3

# Direct product in the case of finite blocks

"Speak, friend..."


## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3




"Speak, friend..."

### Example 3

 $C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3

"Speak, friend..."


 $C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets


$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \quad ?$



## Direct product in the case of finite blocks

"Speak, friend..."

### Example 3


 $C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3
$$G' = C_3 \text{ acting on (non empty) subsets}$$
$$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \text{ ?}$$
$$x \begin{array}{c} \bullet \\ \bullet \\ \circ \end{array}$$
$$x \begin{array}{c} \bullet \\ \circ \\ \circ \end{array}$$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \quad ?$

$$x \begin{array}{c} \bullet \\ \bullet \\ \circ \end{array} + x \begin{array}{c} \circ \\ \bullet \\ \bullet \end{array}$$


$$x \begin{array}{c} \bullet \\ \circ \\ \circ \end{array} + x \begin{array}{c} \circ \\ \circ \\ \bullet \end{array}$$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \quad ?$$


$$\begin{array}{ccccc}
 x \begin{array}{c} \bullet \\ \bullet \\ \circ \end{array} & + & x \begin{array}{c} \circ \\ \bullet \\ \bullet \end{array} & + & x \begin{array}{c} \bullet \\ \circ \\ \bullet \end{array} \\
 x \begin{array}{c} \bullet \\ \circ \\ \circ \end{array} & + & x \begin{array}{c} \circ \\ \circ \\ \bullet \end{array} & + & x \begin{array}{c} \circ \\ \circ \\ \circ \end{array}
 \end{array}$$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \text{ ?}$

$O(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix})$


$O(x \begin{smallmatrix} \bullet \\ \circ \\ \circ \end{smallmatrix})$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \quad ?$


$$O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) \cdot O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right)$$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \quad ?$


$$O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) \cdot O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) = O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right)$$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \quad ?$


$$O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) \cdot O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) = O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right)$$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \text{ ?}$

$$O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) \cdot O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) = O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix}\right)$$




# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \text{ ?}$

$$O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right).O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) = O\left(\begin{smallmatrix} \bullet & \bullet \\ \bullet & \bullet \\ \circ & \circ \end{smallmatrix}\right) + O\left(\begin{smallmatrix} \bullet & \bullet \\ \bullet & \circ \\ \circ & \bullet \end{smallmatrix}\right) + O\left(\begin{smallmatrix} \bullet & \bullet \\ \bullet & \circ \\ \circ & \bullet \end{smallmatrix}\right)$$


$$O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right).O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right)$$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow$  Orbit algebra of  $C_3 \times \mathfrak{S}_\infty$  ?

$$O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right).O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) = O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix}\right)$$


$$O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right).O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) = O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right)$$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \text{ ?}$

$$O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right).O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) = O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right)$$


$$O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right).O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) = O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right)$$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow$  Orbit algebra of  $C_3 \times \mathfrak{S}_\infty$  ?

$$O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix}\right).O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix}\right) = O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix} x \begin{smallmatrix} \circ \\ \circ \\ \circ \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix} x \begin{smallmatrix} \circ \\ \bullet \\ \bullet \end{smallmatrix}\right)$$


$$O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix}\right).O\left(\begin{smallmatrix} \bullet \\ \circ \\ \circ \end{smallmatrix}\right) = O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix} \begin{smallmatrix} \bullet \\ \circ \\ \circ \end{smallmatrix}\right) + O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix} \begin{smallmatrix} \circ \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix} \begin{smallmatrix} \circ \\ \circ \\ \bullet \end{smallmatrix}\right)$$

# Direct product in the case of finite blocks

"Speak, friend..."

## Example 3

$C_3 \times \mathfrak{S}_\infty$  acting on blocks of size 3



$G' = C_3$  acting on (non empty) subsets

$\mathbb{K}[x]^{G'} \longleftrightarrow \text{Orbit algebra of } C_3 \times \mathfrak{S}_\infty \text{ ?}$

$$O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right).O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) = O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \circ \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(x \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} x \begin{smallmatrix} \circ \\ \circ \\ \bullet \end{smallmatrix}\right)$$

$$O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right).O\left(\begin{smallmatrix} \bullet \\ \circ \\ \circ \end{smallmatrix}\right) = O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} \begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} \begin{smallmatrix} \circ \\ \bullet \\ \circ \end{smallmatrix}\right) + O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \circ \end{smallmatrix} \begin{smallmatrix} \circ \\ \circ \\ \bullet \end{smallmatrix}\right) + 3 O\left(\begin{smallmatrix} \bullet \\ \bullet \\ \bullet \end{smallmatrix}\right)$$

The tower has shape  $H_0, H, H, H \dots$

Lemma to prove

$G$  has tower  $H_0 H_1 H_2 H_3 \Rightarrow H_1 = H_2$

Proof.

An element  $s \in G$  stabilizing the blocks  $\leftrightarrow$  a quadruple

$g \in H_1 \rightarrow \exists (1, g, h, k), \quad h, k \in H_1.$

Let  $\sigma$  be an element of  $G$  that permutes "straightforwardly" the first two blocks and fixes the other two.

Conjugation of  $x$  by  $\sigma$  in  $G \rightarrow y = (g, 1, h, k)$

Then:  $x^{-1}y = (g, g^{-1}, 1, 1)$

By arguing that the tower does not depend on the ordering of the blocks,  $g^{-1}$  and therefore  $g$  are in  $H_2$ .

In the infinite case, apply to each restriction to four consecutive blocks of the fixator of the previous ones in  $G$ .

Profile, conjectures ooo	Nested block system oooooo	One superblock oo	Classification oooo	Bonus
-----------------------------	-------------------------------	----------------------	------------------------	-------

