

This Dissertation  
entitled  
Search for lepton flavor violating decays of the Higgs boson  
typeset with  $\text{NDdiss2}_{\varepsilon}$  v3.0 (2005/07/27) on June 25, 2018 for  
Fanbo Meng

This L<sup>A</sup>T<sub>E</sub>X 2 <sub>$\varepsilon$</sub>  classfile conforms to the University of Notre Dame style guidelines established in Spring 2004. However it is still possible to generate a non-conformant document if the instructions in the class file documentation are not followed!

Be sure to refer to the published Graduate School guidelines at <http://graduateschool.nd.edu> as well. Those guidelines override everything mentioned about formatting in the documentation for this  $\text{NDdiss2}_{\varepsilon}$  class file.

It is YOUR responsibility to ensure that the Chapter titles and Table caption titles are put in CAPS LETTERS. This classfile does *NOT* do that!

*This page can be disabled by specifying the “noinfo” option to the class invocation.  
(i.e., \documentclass[... , noinfo] {nddiss2e} )*

**This page is *NOT* part of the dissertation/thesis, but  
MUST be turned in to the proofreader(s) or the  
reviewer(s)!**

$\text{NDdiss2}_{\varepsilon}$  documentation can be found at these locations:

<http://www.gsu.nd.edu>  
<http://graduateschool.nd.edu>

Search for lepton flavor violating decays of the Higgs boson

A Dissertation

Submitted to the Graduate School  
of the University of Notre Dame  
in Partial Fulfillment of the Requirements  
for the Degree of

Doctor of Philosophy

by

Fanbo Meng,

---

Colin Jessop, Director

---

, Director

Graduate Program in Physics

Notre Dame, Indiana

June 2018

© Copyright by

Fanbo Meng

2017

All Rights Reserved

Search for lepton flavor violating decays of the Higgs boson

Abstract

by

Fanbo Meng

this going to put with the abstract, a summary of the whole analysis

NEW DEDICATION NAME

To be written

## CONTENTS

FIGURES . . . . .	v
TABLES . . . . .	viii
PREFACE . . . . .	x
ACKNOWLEDGMENTS . . . . .	xi
CHAPTER 1: Introduction . . . . .	1
1.1 Pre-LHC era on lepton Flavor violation(LFV) search and results . . . . .	1
1.2 Overview of CMS Run1 and Run2 direct search on LFV . . . . .	1
CHAPTER 2: Theory . . . . .	2
2.1 Standard model . . . . .	2
2.1.1 spontaneous symmetry breaking and Higgs mechanism . . . . .	5
2.2 Lepton flavour violation in beyond stand model theories . . . . .	7
CHAPTER 3: LHC and CMS experiment . . . . .	8
3.1 LHC accelerator . . . . .	8
3.2 Compact Muon Solenoid . . . . .	10
3.2.1 Tracker . . . . .	13
3.2.2 The electromagnetic calorimeter . . . . .	15
3.2.3 The hadron calorimeter . . . . .	16
3.2.4 Muon detector . . . . .	18
3.2.5 Trigger . . . . .	19
CHAPTER 4: Datasets . . . . .	22
4.1 Datasets used in LFV analysis . . . . .	22
4.1.1 $H \rightarrow \mu\tau_h$ . . . . .	22
4.1.2 $H \rightarrow e\tau_h$ . . . . .	23
4.2 Event reconstruction . . . . .	23

4.2.1	Particle flow algorithm . . . . .	27
4.2.1.1	Tracking and Calorimeter algorithm . . . . .	27
4.2.2	Muon reconstruction and selection criteria . . . . .	32
4.2.3	Electron identification . . . . .	34
4.2.4	Tau lepton reconstruction . . . . .	37
4.2.5	Jet reconstruction . . . . .	43
4.2.6	Missing transverse momentum . . . . .	46
4.3	Event simulation . . . . .	47
CHAPTER 5: LFV event selection . . . . .		49
5.1	$H \rightarrow \mu\tau_h$ . . . . .	49
5.1.1	Loose selection . . . . .	49
5.1.2	Cut-based analysis . . . . .	51
5.1.3	Multivariate analysis . . . . .	54
5.2	$H \rightarrow e\tau_h$ . . . . .	55
5.2.1	Loose selection . . . . .	55
5.2.2	Cut-based analysis . . . . .	60
CHAPTER 6: LFV analysis background estimation . . . . .		62
6.0.1	Background estimates for $H \rightarrow \mu\tau_h$ . . . . .	62
6.0.1.1	Misidentified Leptons . . . . .	62
6.0.1.2	Other backgrounds . . . . .	67
6.0.2	Background estimates for $H \rightarrow e\tau_h$ . . . . .	68
6.0.2.1	Misidentified Leptons . . . . .	68
6.0.2.2	$Z \rightarrow \tau\tau$ . . . . .	70
6.0.2.3	Other backgrounds . . . . .	72
CHAPTER 7: Signal extraction and systematics . . . . .		73
7.1	Boosted decision trees method . . . . .	73
7.2	Statistical methods . . . . .	76
7.3	Systematics . . . . .	82
7.3.1	Systematics used in $H \rightarrow \mu\tau_h$ . . . . .	82
7.3.2	Systematic uncertainties in $H \rightarrow e\tau_h$ . . . . .	86
CHAPTER 8: LFV Higgs decay searching results . . . . .		90
8.0.1	Results in $H \rightarrow \mu\tau_h$ search . . . . .	90
8.0.2	Results in $H \rightarrow e\tau_h$ search . . . . .	98
CHAPTER 9: Conclusion . . . . .		103
BIBLIOGRAPHY . . . . .		104

## FIGURES

2.1	Standard model particle group[6]	3
3.1	A sketch view LHC injection chain and main experiments associated [53].	11
3.2	A sketch view of CMS detector [18]	12
3.3	The structure of tracker in CMS [18]	14
3.4	ECAL geometrical configuration [35]	16
3.5	A longitudinal view of CMS HCAL sub-detector [18]	17
3.6	A overview of muon chamber configuration in CMS [19]	20
4.1	A sketch view of CMS detecter. Examples are given to show how the particles interact with different sub-detectors.	31
4.2	Electron BDT-based ID shows good discriminating power against background in both EB and EE [50]	36
4.3	PF electron isolation shows better performance in both EB and EE with respect to detector based isolation variable [50]	37
5.1	Expected limits based on an Asimov dataset as a function of $M_T(\tau, MET)$ for the different categories.	52
5.2	Expected limits based on an Asimov dataset as a function of $M_{jj}$ for the 2 jet categories.	54
5.3	Distributions of the input variables to the BDT for the $H \rightarrow \mu\tau_h$ channel.	56
5.4	Expected limits based on an Asimov dataset as a function of $M_T(\tau, MET)$ for the different categories.	57
5.5	Overtraining checking for the BDT training in the TMVA package.	57

5.6	With loose selection conditions, the comparison of the observed collinear mass distributions with background from prediction. The shaded grey bands indicate the total background uncertainty. The open histograms correspond to the expected signal distributions for $\mathcal{B}(H \rightarrow e\tau_h) = 100\%$ in the 0-jet, 1-jet and 2-jet categories, respectively. . . . .	59
6.1	$f_\tau$ ratio for the $\tau$ fake rate calculation shows in term of tau decay modes, $p_T$ , ECAL barrel, ECAL endcap, and $ \eta $ . . . . .	65
6.2	$f_\tau$ ratio for the $\mu$ fake rate calculation shows in term of muon $p_T$ and $ \eta $ . . . . .	66
6.3	The validation of full data driven method for the misidentified lepton background. 30% of uncertainty is assumed in both cases. . . . .	67
6.4	$f_\tau$ and $f_\tau$ ratio for the fake rate calculation shows in term of $p_T$ and $ \eta $ . . . . .	71
6.5	Data driven method validation in Region II with e and $\tau$ leptons in same sign of charge . . . . .	72
7.1	Tree structure example in BDT . . . . .	75
7.2	Test statistics distribution of signal+background and background only PDFs and the observed valued shown in arrow. . . . .	79
7.3	Signal strength modifier $\mu$ at 95% $CL_s$ distribution from the MC pseudo-data. The right plot is the cumulative distribution of $\mu^{95\%}$ with $\pm 1\sigma$ and $\pm 2\sigma$ bands. . . . .	80
8.1	$M_{col}$ distribution after the selection. The signal and backgrounds in the plots have been normalized to the best fit values. The gray bands shows the total uncertainties in each bin and the signal is plotted with the branching ratio of 5% of the higgs decay branching ratio for visualization purpose. . . . .	92
8.2	BDT discriminator distribution after the selection. The signal and backgrounds in the plots has been normalized to the best fit values. The gray bands shows the total uncertainties in each bin and the signal is plotted with the branching ratio of 5% of the higgs decay branching ratio for visualization purpose. . . . .	93
8.3	The expected and observed 95% CL limits on branching ratio of total $H \rightarrow \mu\tau$ which shows the results from both $H \rightarrow \mu\tau_h$ and $H \rightarrow \mu\tau_e$ in the $M_{col}$ fit analysis and BDT fit analysis. . . . .	96

8.4	Upper limit of flavour violating Yukawa coupling $ Y_{\mu\tau} ,  Y_{\tau\mu} $ from the BDT fit analysis. The red solid line represents the expected limit and the black solid line is the observed limit. 68% and 95% range of containing the observed limit is shown in the green and yellow band. The result from $\tau \rightarrow 3\mu$ is shown in dark green [45, 46, 59] and the one from $\tau \rightarrow \mu\gamma$ is shown in lighter green [45, 59]. The theoretical naturalness limit $Y_{ij}Y_{ji} \leq m_i m_j / v^2$ is shown by purple diagonal line [45]. . . . .	97
8.5	$M_{col}$ distribution after the selection. The signal and backgrounds in the plots have been normalized to the best fit values. The gray bands shows the total uncertainties in each bin and the signal is plotted with the branching ratio of 0.69% of the higgs decay branching ratio for visualization purpose. . . . .	99
8.6	95% CL upper limit on branching ratio on $H \rightarrow e\tau$ with both the results from $H \rightarrow e\tau_h$ and $H \rightarrow e\tau_\mu$ and the combined. . . . .	100
8.7	Upper limit of flavour violating Yukawa coupling $ Y_{e\tau} ,  Y_{\tau e} $ from the combined $H \rightarrow e\tau$ result. The red solid line represents the expected limit and the black solid line is the observed limit. 68% and 95% range of containing the observed limit is shown in the green and yellow band. The result from $\tau \rightarrow 3e$ is shown in gray, the one from $\tau \rightarrow e\gamma$ is shown in dark green and presented analysis is in light blue. The theoretical naturalness limit $Y_{ij}Y_{ji} \leq m_i m_j / v^2$ is shown by purple diagonal line [45] . . . . .	102

## TABLES

2.1	Mediators in standard model [44] . . . . .	3
4.1	Monte Carlo samples used in the search, together with their respective cross sections. . . . .	24
4.2	Continue with MC samples used in the analysis. . . . .	25
4.3	Signal and background MC samples . . . . .	26
4.4	Iterative tracking steps take in CMS . . . . .	29
4.5	Muon ID used in the analysis, for the LHC data 2016, running period BCDEF. . . . .	33
4.6	Muon ID used in the analysis, for the LHC data 2016, running period G and H, also the monte Carlo samples. . . . .	34
4.7	Dominant hadronic $\tau$ lepton decays branching fractions and the associated intermediate resonance. The h stands for both $\pi$ and K. The table is symmetric under charge conjugation. . . . .	40
4.8	$\tau$ hadronic decay mode hypothesis signatures compatibility tests. $m_\tau$ is required to be in the mass window . . . . .	42
5.1	Selection criteria in each category with the optimization of the $H \rightarrow \mu\tau_h$ analysis . . . . .	53
5.2	Selection criteria for each event category after cut optimization, for the $H \rightarrow e\tau_h$ channel . . . . .	61
6.1	Definition of the samples used to estimate the misidentified lepton background. . . . .	63
6.2	Scale factor to the events in which a muon is misidentified as a tau	68

7.1	Part one of the systematic uncertainties considered in $H \rightarrow \mu\tau_h$ analysis. All of the uncertainties listed in the systematic tables are correlated between categories besides the ones after the sign $\oplus$ . These are the uncertainties correlated in within each category but independent between categories. The theoretical uncertainties related to the acceptance and migration of events are listed in a range. The negative or positive values indicate a anticorrelated or correlated between categories . . . . .	84
7.2	Part two of the systematic uncertainties considered in $H \rightarrow \mu\tau_h$ analysis . . . . .	85
7.3	b tagging veto systematic uncertainty in each category . . . . .	85
7.4	The normalization systematic uncertainties considered in $H \rightarrow e\tau_h$ analysis. The uncertainties are correlated between categories besides the ones after $\oplus$ . These uncertainties are uncorrelated between categories. . . . .	88
7.5	The systematic uncertainties that affect the shape of $M_{col}$ distribution. . . . .	88
7.6	Theoretical uncertainties that affects the Higgs boson production cross section. These uncertainties are correlated or anticorrelated(with minus sign superscript) between all of the categories. . . . .	89
8.1	$M_{col}$ fit analysis expected and observed upper limits at 95% CL and the best fit branching fractions in each of the categories. . . . .	94
8.2	BDT fit analysis expected and observed upper limits at 95% CL and the best fit branching fractions in each of the categories. . . . .	95
8.3	Events yields of the signal and backgrounds are shown in the mass range $100 \text{ GeV} < M_{col} < 150 \text{ GeV}$ . The branching ratio of $H \rightarrow e\tau_h$ is assumed to be 0.69% of the SM higgs production cross section. The numbers in the table are normalized to the integrated luminosity of $19.7 \text{ fb}^{-1}$ . . . . .	100
8.4	$M_{col}$ fit analysis expected and observed upper limits at 95% CL and the best fit branching fractions in each of the categories of both $H \rightarrow e\tau_h$ and $H \rightarrow e\tau_\mu$ analyses. The asymmetric one standard-deviation uncertainties around the expected limits are shown in parentheses. . . . .	101

## PREFACE

probably I will write some preface, but it depends  $H \rightarrow \mu\tau_h$

## ACKNOWLEDGMENTS

this part is going to be filled with acknowledge

## CHAPTER 1

### Introduction

1.1 Pre-LHC era on lepton Flavor violation(LFV) search and results

1.2 Overview of CMS Run1 and Run2 direct search on LFV

## CHAPTER 2

### Theory

#### 2.1 Standard model

Standard model(SM) describes the properties of the fundamental particles currently known in the universe and the interactions amount them. The discovery of Higgs boson in 2012 implement the last missing piece in SM. The particles in SM is shown in Figure. 2.1. Besides Higgs boson, there are three generations of leptons and quarks and four gauge bosons. Leptons and quarks are fermions that compose the matters currently known in the universe, while the gauge bosons serve as the mediators of interactions.

SM is a gauge theory, which describes three natural force, the strong, electromagnetic and weak force by symmetry group  $SU(3)_C \times SU(2)_L \times U(1)_Y$ . Electroweak section is described by  $SU(2)_L \times U(1)_Y$ . The L in  $SU(2)_L$  stands for the left chiral signature of the weak interaction and the Y in  $U(1)_Y$  stands for the weak hypercharge. Strong interaction is described by  $SU(3)_C$ , in which the C stands for the colors of quarks. Photons, gluons and  $W^\pm$ , Z bosons are the mediators of electromagnetic, strong and weak force respectively. Gravity force is not included in SM. Each particles in Figure. 2.1 has its anti-particle or the anti-particle is the particle itself. The force and mediators characters are summarized in Table. 2.1

Fermion fields in SM can be further categorized. In weak interaction, left-

# Standard Model of Elementary Particles




Figure 2.1. Standard model particle group[6]

TABLE 2.1

Mediators in standard model [44]

Force	Mediator	Charge	Mass(GeV)	Range(m)
Strong	g(8 gluons)	0	0	$10^{-15}$
Electromagnetic	$\gamma$ (photon)	0	0	$\infty$
Weak	$W^\pm$	$\pm 1$	80.379	$10^{-18}$
	Z	0	91.1876	$10^{-18}$

handed fermions are isodoublets, while right-handed fermions are isosinglets. Only the left-handed fermions or right-handed anti-fermions participate in the weak interaction. Weak hypercharge is a combined quantity, which is defined by the electric charge  $Q$  the third component of the weak isospin  $I_f^3$ ,  $Y = Q - I_f^3$ . Weak interaction, together with Electromagnetic interaction are best described and understood by the electro-weak theory, which is be further discussed in the next section. Quarks are described by  $SU(3)$ , which are color triplets, while leptons are color singlets.

Mediators in the interaction are represented by the gauge boson fields.  $B_\mu$  is associated with symmetry  $U(1)_Y$  and the corresponds generator is  $Y$ .  $W_\mu^{1,2,3}$  are associated to  $SU(2)_L$  symmetry with the generator  $T^a$ (a=1,2,3).  $T^a$  are the  $2 \times 2$  Pauli matrices.  $G_\mu^{1...8}$  are associated with the  $SU(3)_c$  symmetry and the corresponding generators are the Gell-Mann matrices. The field strengths are expressed as

$$\begin{aligned} G_{\mu\nu}^a &= \partial_\mu G_v^a - \partial_v G_\mu^a + g_s f^{abc} G_\mu^b G_v^c \\ W_{\mu\nu}^a &= \partial_\mu W_v^a - \partial_v W_\mu^a + g_2 \epsilon^{abc} W_\mu^b W_v^c \\ B_{\mu\nu} &= \partial_\mu B_v - \partial_v B_\mu \end{aligned} \quad (2.1)$$

The  $g_s$  and  $g_2$  are the coupling constant of  $SU(3)_C$  and  $SU(2)_L$  respectively. With the requirement of local gauge invariable, covariant derivatives are widely used. An example of covariant derivative acting on left-handed quarks in Lagrangian is expressed as

$$D_\mu \psi = (\partial_\mu - ig_s T_a G_\mu^a - ig_2 T_a W_\mu^a - ig_1 \frac{Y_q}{2} B_\mu) \quad (2.2)$$

The masses of the gauge bosons in weak interaction and fermions are generated by spontaneous symmetry breaking. Fermions specifically are generated by Higgs mechanism. If mass terms are directly added in, the local  $SU(2) \times U(1)$  will be

destroyed [38]. Local symmetry referring to the transformations on the fields are the functions that involve space-time. In SM, the interaction between fermion fields and scalar fields are through Yukawa's interaction. The SM in  $SU(3)_C \times SU(2)_L \times U(1)_Y$  symmetry without the mass terms and Yukawa's interactions is given by

$$L_{SM} = -\frac{1}{4}G_{\mu\nu}^a G_a^{\mu\nu} - \frac{1}{4}W_{\mu\nu}^a W_a^{\mu\nu} - \frac{1}{4}B_{\mu\nu} B^{\mu\nu} + \bar{L}_i iD_\mu \gamma^\mu L_i + \bar{e}_{R_i} iD_\mu \gamma^\mu e_{R_i} + \bar{Q}_i iD_\mu \gamma^\mu Q_i + \bar{\mu}_{R_i} iD_\mu \gamma^\mu \mu_{R_i} + \bar{R}_i iD_\mu \gamma^\mu d_{R_i} \quad (2.3)$$

### 2.1.1 spontaneous symmetry breaking and Higgs mechanism

In SM, the electroweak sector follows the  $SU(2)_L \times U(1)_Y$ , which is spontaneously broken in the  $SU(2)_L$  part. This mechanism plays an important role in giving mass to the mediator bosons in weak interaction and introducing the Higgs boson(Higgs). Higgs is responsible for generating the mass of the fermions in SM, while SM remains renormalizable in local symmetry. A good example of showing the concept of spontaneous symmetry breaking with the  $\phi^4$  theory can be found in [60].

In the real case of SM, spontaneous symmetry breaking is introduced through a scalar doublet:

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_2 - i\phi_1 \\ \phi_4 - i\phi_3 \end{pmatrix}$$

The corresponding terms of the scalar doublet in the SM Lagrangian are expressed as

$$L_s = (D^\mu \phi)^\dagger (D_\mu \phi) - \mu^2 \phi^\dagger \phi - \lambda (\phi^\dagger \phi)^2 \quad (2.4)$$

If  $\mu^2 < 0$  and  $\lambda > 0$ , the doublet field  $\phi$  will have a none zero minimum value,

which is called the vacuum expectation value(vev). In SM, the vev is shown in the following Equation and measured to be 246 GeV.

$$\langle 0|\phi|0\rangle = \begin{pmatrix} 0 \\ \frac{v}{\sqrt{2}} \end{pmatrix} \quad \text{with} \quad v = \left( -\frac{\mu^2}{\lambda} \right)^{1/2} \quad (2.5)$$

When the SU(2) symmetry is spontaneously broken, the scalar doublet field  $\phi$  can expand around vev at first order together with the Higgs field:

$$\phi = \begin{pmatrix} \theta_2 + i\theta_1 \\ \frac{1}{\sqrt{2}}(v + H) - i\theta_3 \end{pmatrix} = e^{i\theta_a(x)T^a(x)/v} \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}}(v + H) \end{pmatrix} \quad (2.6)$$

Taking the unitary gauge, the scalar field transforms  $\phi \rightarrow e^{-i\theta_a(x)T^a(x)/v}\phi$  and takes into  $(D^\mu\phi)^\dagger(D_\mu\phi)$ :

$$\begin{aligned} (D^\mu\phi)^\dagger(D_\mu\phi) &= |(\partial_\mu - ig_2 \frac{T_a}{2}W_\mu^a - \frac{i}{2}g_1 B_\mu)\phi|^2 \\ &= \frac{1}{2}(\partial_\mu H)^2 + \frac{1}{8}g_2^2(v + H)^2|W_\mu^1 + iW_\mu^2|^2 \\ &\quad + \frac{1}{8}(v + H)^2|g_2 W_\mu^3 - g_1 B_\mu|^2 \end{aligned} \quad (2.7)$$

To meets the experimental observations, re-groups the gauge fields to have the  $W^\pm$  and Z bosons:

$$W^\pm = \frac{1}{\sqrt{2}}(W_1 \mp iW_2), Z_\mu = \frac{g_2 W_\mu^3 - g_1 B_\mu}{\sqrt{g_2^2 + g_1^2}}, A_\mu = \frac{g_2 W_\mu^3 + g_1 B_\mu}{\sqrt{g_2^2 + g_1^2}}, \quad (2.8)$$

From the terms in Equation 2.7 that involving the mass of  $W^\pm$  and Z boson, these

vector bosons acquire the mass as

$$M_W = \frac{1}{2}vg_2 \quad \text{and} \quad M_Z = \frac{1}{2}v\sqrt{g_2^2 + g_1^2} \quad (2.9)$$

while the photon  $A_\mu$  remains massless.

$$\begin{pmatrix} \gamma \\ Z^0 \end{pmatrix} = \begin{pmatrix} \cos\theta_W & \sin\theta_W \\ -\sin\theta_W & \cos\theta_W \end{pmatrix} \begin{pmatrix} B \\ W_3 \end{pmatrix}$$

$$M_Z = \frac{M_W}{\cos\theta_W} \quad \text{and} \quad \cos\theta_W = \frac{g_2}{\sqrt{g_2^2 + g_1^2}} \quad (2.10)$$

## 2.2 Lepton flavour violation in beyond stand model theories

Three questions need to answer: 1. Higgs in standard model 2. Why LFV is not possible in SM 3. LFV in beyond standard models

why LFV is not possible in SM

## CHAPTER 3

### LHC and CMS experiment

LHC [40] is the most powerful hadron accelerator and collider ever built. The circumference of this superconducting collider is 26.7 km and the design full operation energy is 14 TeV. In 2012 LHC operates on 8 TeV and in 2016, it boosts up to 13 TeV. There are four experiments holds on the LHC ring, ALICE, ATLAS, CMS and LHCb. ATLAS and CMS are two general purpose detector, aiming for the high luminosity in the order  $L = 10^{34} cm^2 s$ . LHCb focuses on the b physics study, while ALICE studies the lead-lead collision.

#### 3.1 LHC accelerator

A sketch view of the proton accelerating process is shown in Figure. 3.1. LHC is the last and most powerful accelerator and CMS experiment is located in one of the collision cites, however, before the beams are injected into LHC, a series of steps are needed and necessary to meet the requirement on the beams. For the proton-proton(p-p) collision, protons are from the source Duoplasmatron, which uses electric field to break down hydrogen gas into proton and electrons and protons are accelerated by a 90 kV supply. Leaving the source, protons are focused and accelerated to 750 keV by radio frequency quadrupole. Then the protons are further accelerated by linear accelerator Linac2 to 50 MeV. Proton synchrotron

booster further accelerate protons from 50 MeV to 1.4 GeV and injects protons into Proton synchrotron(PS). PS accelerates protons to 25 GeV and later accelerator super proton synchrotron boost the protons to 450 GeV. LHC is the last ring in the whole accelerating process and accelerate the protons to its final energy 6.5 GeV(or 7 GeV). In a normal fill with protons, LHC holds 2808 bunches with an approximation of  $10^{11}$  protons.

There are thousands of superconducting magnets along the LHC ring to bend and focus the beam. Among the magnets, there are 1232 main dipoles, which are used to bend the beam with an operating field above 8 T. Other type of magnets, for example, quadrupoles magnets can tight the beam either vertically or horizontally while sextupole, octupole and decapole can help fine tuning the the field. Radiofrequency cavities(RF) in the LHC are used to accelerate the protons from 450 GeV to 6.5 GeV(or 7 GeV), keep the bunches in the beam pipe compact and store the energy loose from synchrotron radiation. Eight RFs per beam, providing 16 MV longitudinal oscillating voltage with 400 MHz superconducting cavity system.

Machine luminosity is an important parameter of the collider. For a process under study, the number of events created per-second  $N_{\text{event}}$  is shown in Equation. 3.1, in which  $L$  is the machine luminosity and  $\sigma_{\text{event}}$  is the cross section of that process.

$$N_{\text{event}} = L\sigma_{\text{event}} \quad (3.1)$$

Machine luminosity is determined by a number of factors as shown in Equation. 3.2.  $n_b$  and  $N_b$  are the number of bunches per-beam and the number of pro-

tons per-bunch.  $f_{rev}$  and  $\gamma_\gamma$  are the revolution frequency and relativistic gamma factor respectively.  $\beta^*$  is the amplitude function of the beam at the collision point while function F describes the reduction of luminosity because of the crossing angle. This machine luminosity is also called instantaneous luminosity which is the luminosity at a unite time. The integrated luminosity which later referred as luminosity is instantaneous luminosity integrating over time 3.1.

$$L = \frac{N_b^2 n_b f_{rev} \gamma_\gamma}{4\pi\epsilon\beta^*} F \quad (3.2)$$

### 3.2 Compact Muon Solenoid

Compact muon solenoid(CMS) is general purpose detector, which institutes in one of LHC collision point. It is a high performance detector that covers a broad range of physics researches, like Standard model physics, supersymmetry and dark matter. CMS detector is designed to have good muon momentum and position resolution over large range of energy and angles, good charged particle momentum resolution and high identification efficiency within inner tracker, good electromagnetic energy and position resolution and high photon and lepton isolation efficiency in high luminosity condition, good miss transverse momentum and jet energy resolution.

An general view of CMS detector is shown in Figure. 3.2. The detector is composed of a set of sub-detectors from the inside and out in a ring structure. The main sub-detectors are the following:

- Tracker consists two parts, the pixel detector and silicon tracks which are used




Figure 3.1. A sketch view LHC injection chain and main experiments associated [53].




Figure 3.2. A sketch view of CMS detector [18]

measure the momentum and tracks of the charged particles.

- Electromagnetic calorimeter mainly measures the energy of the position of electrons and photons, also other particles will leave some percentage of energy inside.
- Hadron calorimeter measures the energy of hadrons
- Muon detector measures the tracks and momentum of the muons.

Another outstanding feature of CMS is the superconducting magnet system, which provide a 3.8 T magnetic field. The configuration of the magnet system drives the design and layout of the detector. Besides the sub-detectors listed above, trigger system is also crucial for the success of the whole program. Trigger system consists two part, the hardware based Level one trigger and the software

based High level trigger. Trigger system does the initial selection of interesting events from the huge flux of events per-collision, which makes it possible for data-acquisition and recording later. The details of the sub-detectors and other systems mentioned will be introduced later.

In general CMS detector has the dimension, 21.6 m long, 14.6 in diameter and weighted in total 12500 tonne. The coordinate system adopted by CMS sets the center at the collision point. The x-axis points towards the center of LHC and z-axis points along the beam direction. so y-axis is on the vertical direction. The azimuthal angle  $\phi$  is used to measure the angle from x-axis in the x-y plane and polar angle  $\theta$  measures the angle from z-axis and r is the radial coordinate. Another variable pseudorapidity  $\eta$ , defined as  $\eta = -\ln(\theta/2)$ . In the case  $E \gg m$ , pseudorapidity can be approximated as  $\eta = -\ln\left(\frac{E+p_z}{E-p_z}\right)$ , where  $p_z$  is the longitudinal component of momentum [18].

### 3.2.1 Tracker

The main purpose of inner tracking system of CMS is measuring the trajectories of charged particles from the LHC collision. The efficient and precise measurement is crucial for the reconstruction and identification of particles. LHC operates with the instantaneous luminosity in the order of  $10^{34} cm^{-2}s^{-1}$ , in average more than 20 p-p interactions and 1000 particles per-bunch crossing. High granularity and fast response is primary important for the tracker system. To measure the trajectories precisely, low interaction of tracker materials with incoming particles, like multi-scattering, photon conversion and nuclear interaction is also important. In the long operation period, radiation hardness of the tracker material is needed. All of these factors drive the design of CMS tracker system.

The tracker system of CMS surrounds the collision point with the dimension 5.8 m in length and 2.5 m in diameter, with a coverage up to pseudorapidity  $|\eta| < 2.5$ . A overview of the tracker system layout is shown in Figure. 3.3. Three layers of silicon pixel detector modules surround the interaction points and two additional disks of pixel modules on each side, in all 66 million pixels of the size  $100 \times 150\mu\text{m}$  each. Following the pixel detector is the silicon strip tracker. There are four components of strip tracker, tracker Inner barrel(TIB), tracker inner Discs(TID), tracker outer barrel(TOB) and tracker end caps(TEC). The strip tracker components arrangement as shown in Figure. 3.3, consisting of about 10 million strips and 10 layers.




Figure 3.3. The structure of tracker in CMS [18]

### 3.2.2 The electromagnetic calorimeter

The CMS Electromagnetic calorimeter(ECAL) is a hermetic homogeneous lead tungstate( $PbWO_4$ ) calorimeter. The whole sub-detector is composed of two parts, the central barrel(EB) covering the range  $|\eta| < 1.479$  and the endcap disks(EE) covering the range  $1.479 < |\eta| < 3.0$ . EB is made up of 61200  $PbWO_4$  crystals with  $22 \times 22 \text{ mm}^2$  in the front face, 23 cm in length(25.8 in radiation lengths). EE is made up of 7324 crystals per disk with front face  $29 \times 29 \text{ mm}^2$ , 22 in length(24.7 in radiation lengths) and a preshower detector(ES) [35]. The geometrical configuration of ECAL is shown in Figure. 3.4.

The  $PbWO_4$  crystals used in ECAL have high density( $8.28 \text{ g/cm}^3$ ), short radiation length(0.89 cm) and small Molière radius(2.2 cm), together with the specific geometrical parameters used, rendering ECAL good energy resolution, fast response, fine granularity and radiation resistant. Photodetectors are placed at the end of the crystals. In EB, avalanche photodiodes(APDs) are used, while vacuum phototriodes(VPTs) are used in EE. Both types of the photodiode show good performance in the environment with hard radiation and 4-T magnetic field. ES is in front of EE in the pseudorapidity range  $1.653 < |\eta| < 2.6$ . ES is a sampling detector with silicon strip sensors placed behind the lead radiator to measure the energy and position of the incoming particles.

Each half of EB is composed of 18 supermodules that one contains 1700 crystals. The  $PbWO_4$  crystal relative energy resolution is measured with ECAL supermdules directly exposed to electron beam without considering the materials in front as the case in reality. The relative energy resolution as a function of electron




Figure 3.4. ECAL geometrical configuration [35]

energy can be expressed as

$$\frac{\sigma}{E} = \frac{2.8\%}{\sqrt{E/\text{GeV}}} \oplus \frac{12\%}{E/\text{GeV}} \oplus 0.3\%$$

The first term stands for the contribution from stochastic factors, like the number fluctuation in production of the secondary particles. The second term is the noise contribution coming from electronics and digitization, while the last is a constant term that covers the other factors [10].

### 3.2.3 The hadron calorimeter

CMS hadron calorimeter(HCAL) is a hermetic sampling calorimeter, which is important for the measurement of the energy and momentum of hadrons, also the missing energy that caused by the non-interacting particles. HCAL is composed mainly by HCAL barrel(HB), HCAL endcaps(HE) and forward calorimeter(HF).

The mechanical structure of HCAL is shown in Figure. 3.5.

HB is mostly composed of several layer of brass absorber plates, between which




Figure 3.5. A longitudinal view of CMS HCAL sub-detector [18]

are the plastic scintillator tiles. The inner most and out most layer plates are made of stainless steel to gain structural strength. When hadrons hit the absorber, secondary particles are produced in showers. As the showers develops, the alternating layers of scintillators are activated and emit blue-violet light. The lights are collected as signals. HB covers the central pseudorapidity range  $|\eta| < 1.3$  with a individual read out unit  $\Delta\eta \times \Delta\phi = 0.087 \times 0.087$ . To have enough sampling depth in HCAL central region because of the limited spaces between ECAL and muon detector, an extra outer calorimeter(HO) is installed. HO utilizes the outside solenoid coil as additional absorber to insert plastic scintillators. Similar to HB, HE is also made of brass absorber and plastic scintillator tiles layers, which covers the range  $1.3 < |\eta| < 3.0$ . The read out units has the geometry  $\Delta\eta \times \Delta\phi = 0.17 \times 0.17$  in HE [18]. An combined ECAL and HCAL energy resolution [67] measured in test beam with pions is

$$\frac{\sigma}{E} = \frac{110\%}{\sqrt{E/\text{GeV}}} \oplus 0.9\%$$

HF situates  $\pm 11$  m from the interaction point to complement the large pseudorapidity measurement of HE in the range  $3.0 < |\eta| < 5.0$ . HF is made of grooved steel plates with quartz fibers. Charged shower particles generate Cherenkov light, which is collected by the quartz fibers as signals. Radiation hardness is critical for the operation of HF.

### 3.2.4 Muon detector

CMS muon detector is designed to measure the momentum and charge of muons. Three types of gas detectors are used in CMS, the barrel drift tube(DT) chambers, the cathode strip chambers(CSC) in the endcaps and resistive plate chambers(RPC) in both barrel and endcap regions.

In muon detector barrel(MB), DT chambers and RPCs are used which covers the pseudorapidity region  $|\eta| < 1.2$ . MB is composed of 250 chambers. The chambers are located in 4 stations inside the magnet return yoke. The yoke is further divided into 5 wheels, each of which is composed of 12 sectors. As shown in Figure. 3.6, the stations are named MB1 to MB4, which are composed one DT chamber and varied number of RPCs that depend on the exact location. DT chamber measures the position the incoming muon which knocks off electrons of the gas atoms in the chamber and collected by a large numbers of charged wires inside.

In muon detector endcaps(ME), 468 CSCs which covers the range  $0.9 < |\eta| < 2.4$  are used. ME also composed of 4 stations of chambers and in each of the discs.

CSC is in a wedge shape and composed of 6 gas gaps. The gap is filled in with a cathodes strip and anode wires what running perpendicularly to the strip. When a muon comes, the knocked off electrons creating avalanche and received by the positive charged wires.

Both DT chambers and CSCs measure the position and can also trigger on the  $p_T$  of muons. To better deal with the high luminosity and improving the trigger muon  $p_T$  resolution, the dedicated trigger system RPCs are added to both MB and ME. RPC is a double-gap bakelite chamber which operates with the avalanches that caused by the muons. RPC can provide additional fast triggering and sharp  $p_T$  threshold. Four RPCs layers are used in the MB first two stations(two each) and another two layers(one each) are used in the last two stations. In the ME, one RPC layers in each of the first three stations.

### 3.2.5 Trigger

LHC is a high luminosity collider in the order of  $10^{34}\text{cm}^2\text{s}^{-1}$ . In the p-p collision, the bunch crossing time is 25 ns and the corresponding frequency is 40 MHz. In each of the 25 ns, there are approximately 20 collisions. This high rate makes it impossible to transmit and record all of the events, also it is not necessary, since most of the events are not of current physics research interests. CMS use a two level trigger system to select events and reduce the event recording rate, Level-1(L1) trigger and High-level trigger(HLT) systems.

L1 trigger is based on custom designed and programmable electronics, situated inside the detector. It utilizes the information from calorimeters and muon system, performing simplified but effective reconstructions, corrections and selections, which reduces the event rate from 40 MHz to 100 kHz. In LHC Run II, the




Figure 3.6. A overview of muon chamber configuration in CMS [19]

L1 trigger system has been upgraded to deal with the increased luminosity and improve the performance. The L1 calorimeter trigger accesses the information from the whole ECAL and HCAL in the granularity of trigger tower level, which approximately correspond to a region  $0.087 \times 0.087$  in  $\eta$  and  $\phi$ . The trigger reconstructs the  $e/\gamma$ , jets,  $\tau$  and the sum energy of the candidates with the algorithms implemented in the time multiplexed trigger architecture [12]. The algorithms implemented at hardware trigger level, with dynamic clustering of trigger towers, pile-up migration and innovated tau and jet reconstruction with various look-up table for the calibrations and corrections [69]. The L1 muon trigger system fully utilizes three muon detectors in the track reconstruction. In general, the track reconstruction of muons are divided into three regions, barrel, overlap and endcap, based on the geometry. Through dedicated construction algorithm, the tracks are

built, so as computing the muon qualities. There informations are used in the triggering processes [62].

HLT in CMS further reduced the event rate from 100 kHz(after L1 trigger selection) down to 1 kHz that is a possible rate for event storage. HLT is software based trigger system and utilized the streamlined version of the CMS soft-ware for event reconstruction on the large computer farm [7]. Maximizing the trigger efficiency and keeping acceptable CPU-time is crucial. HLC accesses to the full granularity and sub-detectors of CMS, including the tracker. The dedicated algorithms used in HLT is the very closed to the ones used in the off-line reconstruction and analysis, besides the for some different parameter configurations [43].

## CHAPTER 4

### Datasets

#### 4.1 Datasets used in LFV analysis

##### 4.1.1 $H \rightarrow \mu\tau_h$

The data sample used in the analysis is from LHC 2016 Runs, recorded by CMS detector. Total integrated luminosity of the analyzed data is  $35.9 \text{ fb}^{-1}$  with center-of-mass energy of  $\sqrt{s} = 13 \text{ TeV}$ . The data sample labeled in the CMS experiment as SingleMuon\_Run2016B,C,D,E,F,G,H. The name SingleMuon in the data sample represents the trigger used in the analysis, which selects the events with at least a muon passed the trigger selection. Gluon gluon fusion Higgs(ggH) [42] production and vector boson fusion Higgs(VBF) [17] are the main Higgs production channels considered in the analysis. For the background samples, besides the misidentified background which is talked about in detail in Chapter6, the other background samples are all generated with Monte Carlo(MC) simulation. POWHEG [58] or MadGraph [11] generator is used for the generation and all of the MC samples, including the signal samples, the parton showering, fragmentation, and decays are performed by Pythia8 [66]. Pileup effect is taking into account in the generator by generate minimum bias events simultaneously and a correction is applied by comparing the data sample. The average number of pileup interaction per bunch crossing is 27. CMS detector environment is simulated by GEANT4 [33]. The

details of the MC samples used in the analysis are summarized in Table. 4.1 and Table. 4.2.

#### 4.1.2 $H \rightarrow e\tau_h$

The search of lepton flavour violation Higgs decay  $H \rightarrow e\tau_h$  is performed with CMS 2012 RunI dataset at center-of-mass energy  $\sqrt{s} = 8$  TeV. The dataset is named as SingleElectron\_Run2012A,B,C,D with an integrated luminosity of  $19.7 \text{ fb}^{-1}$ . Single electron HLT trigger is used which is talked more in details in Chapter 5. A detail list of simulation samples used in the analysis is listed in Table. 4.3. For the signal samples, ggH and VBF Higgs production channels are the main channels considered. For background samples, besides  $Z \rightarrow \tau\tau$  which is produced with embedding technique and misidentified background which is estimated with data-driven method are produced with MC simulation. Various simulation packages are used. Signal samples are produced with PYTHIA8, which uses sophisticated  $\tau$ -lepton decay machinery. Tauola [57] is also used for the simulation of  $\tau$  lepton decay in some of samples. CMS detector environment is simulated by GEANT4.

## 4.2 Event reconstruction

In this section, a general presentation of the event reconstruction algorithm used in CMS, particle-flow(PF) reconstruction is shown. A more detail information about the reconstruction of tracks and the objects related to the lepton flavour violation higgs decay are in the following section.

TABLE 4.1

Monte Carlo samples used in the search, together with their respective cross sections.

Processes	Generator	Cross section [pb]
DYJets $\rightarrow \ell\ell, m_{\ell\ell} > 50$ GeV	MadGraph+pythia8	4954.0
DY1Jets $\rightarrow \ell\ell, m_{\ell\ell} > 50$ GeV	MadGraph+pythia8	1012.5
DY2Jets $\rightarrow \ell\ell, m_{\ell\ell} > 50$ GeV	MadGraph+pythia8	332.8
DY3Jets $\rightarrow \ell\ell, m_{\ell\ell} > 50$ GeV	MadGraph+pythia8	101.8
DY4Jets $\rightarrow \ell\ell, m_{\ell\ell} > 50$ GeV	MadGraph+pythia8	54.8
DYJets $\rightarrow \ell\ell, m_{\ell\ell} < 50$ GeV	MadGraph+pythia8	1861.0
$t\bar{t}$	powheg+PYTHIA	831.76
$t\backslash\bar{t} \rightarrow tw$	powheg+PYTHIA8	35.85
$WZ \rightarrow \ell 3v$	MadGraph+PYTHIA8	3.05
$WZ \rightarrow \ell v 2q$	MadGraph+PYTHIA8	10.71
$WZ \rightarrow 2\ell 2q$	MadGraph+PYTHIA8	5.595
$t \rightarrow 4f$	POWHEG+PYTHIA8	136.02
$\bar{t} \rightarrow 4f$	POWHEG+PYTHIA8	80.95
$WW \rightarrow \ell v 2q$	MadGraph+PYTHIA8	1.212
$ZZ \rightarrow 2\ell 2q$	MadGraph+PYTHIA8	3.22
$VV \rightarrow 2\ell 2q$	MadGraph+PYTHIA8	11.95

TABLE 4.2

Continue with MC samples used in the analysis.

MC simulations	Generator	Cross section [pb]
$VV \rightarrow 2\ell 2q$	MadGraph+PYTHIA8	11.95
$ggH \rightarrow \tau\tau$	POWHEG+PYTHIA8	3.046
$VBFH \rightarrow \tau\tau$	POWHEG+PYTHIA8	0.237
$ggH \rightarrow WW \rightarrow 2\ell 2v$	POWHEG+PYTHIA8	1.103
$VBFH \rightarrow WW \rightarrow 2\ell 2v$	POWHEG+PYTHIA8	0.086
$ZH \rightarrow \tau\tau$	POWHEG+PYTHIA8	0.055
$W^- \backslash W^+ H \rightarrow \tau\tau$	POWHEG+PYTHIA8	0.086
$t\bar{t}H Jet \rightarrow \tau\tau$	MadGraph+PYTHIA8	0.32

TABLE 4.3  
Signal and background MC samples

Processes	Generator	Cross section [pb]
$ggH \rightarrow e\tau$	PYTHIA8	19.27
$VBH \rightarrow e\tau$	PYTHIA8	1.58
$ggH \rightarrow \tau\tau$	POWHEG+PYTHIA6	19.27
$VBF \rightarrow \tau\tau$	POWHEG+PYTHIA6	1.58
$t\bar{t} + \text{jets full leptonic}$	MadGraph+Tauola	26.20
$t\bar{t} + \text{jets Semi leptonic}$	MadGraph+Tauola	109.28
$t \rightarrow tw$	POWHEG+Tauola	56.4
$\bar{t} \rightarrow tw$	POWHEG+Tauola	30.7
$t, \bar{t}(\text{T channel})$	POWHEG+Tauola	11.1 (11.1)
$WW \rightarrow 2l2\nu + \text{jets}$	PYTHIA6+Tauola	5.824
$ZZ \rightarrow 4l$	MadGraph+Tauola	0.18
$ZZ \rightarrow 2l2Q$	MadGraph+Tauola	2.502
$ZZ \rightarrow 2l2\nu$	MadGraph+Tauola	0.716
$WZ \rightarrow 2l2Q$	MadGraph+Tauola	2.21
$WZ \rightarrow 3l\nu$	MadGraph+Tauola	1.06

### 4.2.1 Particle flow algorithm

Particle flow event reconstruction algorithm is the main algorithm used in CMS. PF performs a global event reconstruction [63], which aim to utilize the information from the whole detector to identify individual particles in each event.

#### 4.2.1.1 Tracking and Calorimeter algorithm

In the PF algorithm, the reconstruction of charged particles in the inner tracker is a crucial part. In this section, the reconstruction of the trajectories of charged particles, especially the electron and muon reconstruction is discussed.

Inner tracker is aimed at measuring the track of energetic charged particle. The track finder is based on the Kalman Filtering(KF) [9]. This reconstruction comes in a couple steps. First an initial seed is generated from a couple hits that compatible with a track in the tracker, then a trajectory is builded with the seed and other hits from the tracker along this track. At last a fit is perform on the track to determine the properties of this particle candidate, like the momentum, the charge and the direction. The qualities that can affect the performance of the reconstruction are like the number of hits in the pixel detector, the total number of hits in the tracker, the distance from the cylinder and the energy of this charged particle. This is discussed in more details later.

The performance of the track reconstruction is measured in reconstruction efficiency and misreconstruction rate. The reconstruction efficiency is defined as the ratio of track reconstructed with more than 50% hits from simulated hits and the total simulated tracks. The misreconstruction rate is defined as the fraction of the tracks can not be associated with simulated tracks of the whole simulated tracks. If a charged hadron is not identified by the tracking algorithm, then the hadron

is take as a neutral hadron and measured by the calorimeters. This will affect the jet energy and position resolution. Improving the track reconstruction efficiency while keeping the misreconstructed rate low is critical for PF reconstruction.

In CMS, an iterative tracking is perform, in which the reconstruction of tracks is done is a couple steps. Each step is aimed for a moderate efficiency but with a high purity. After one step, the hits that used to form tracks are masked and a next step is perform. This iterative tracking is down in ten steps if necessary and the detail information is shown in Table. 4.4 and more can be found in [63]. In the table, the name column points out the processes that that step of iteration is aiming at. The seeding column shows the requirement on the seeding, while the targeted tracks column shows the characters of the tracks.

Electron is one of the main particle understudy in  $H \rightarrow e\tau_h$  search. In CMS, electron track reconstruction is taken as a merge of ECAL based and tracker based strategy. The tracker based seeding strategy is as described above, besides the case when energetic photons are radiated, then a preselection based on the number of hits and  $\chi^2$  of the fit is set with the Gaussian-sum filter [8]. The ECAL based electron seeding strategy building the superclusters(SC) to gathering the bremsstrahlung photons. The energy of the SC is taken as the energy sum of the cell crystals inside and the position is evaluated with energy weight. The electron trajectory in the first layer of the the tracker is estimated and the seed from the track is selected. This works in the case when there is not much bremsstrahlung photons, then most of the energy is deposited in ECAL. In the case when soft photons are radiated mostly, ECAL based seeding may still performs well. But when the electrons are in the jets or low energy, either electron contributions are overlapping with other particle or the radiation and bending is too much, it is hard

TABLE 4.4  
Iterative tracking steps take in CMS

Iteration	Name	Seeding	Targeted Tracks
1	InitialStep	pixel triplets	prompt, high $p_t$
2	DetachedTriplet	pixel triplets	from b hadron decays, $R \lesssim 5$ cm
3	LowPtTriplet	pixel triplets	prompt, low $p_t$
4	PixelPair	pixel pairs	recover high $p_t$
5	MixedTriplet	pixel+strip triplets	displaced, $R \lesssim 7$ cm
6	PixelLess	strip triplets / pairs	very displaced, $R \lesssim 25$ cm
7	TobTec	strip triplets / pairs	very displaced, $R \lesssim 60$ cm
8	JetCoreRegional	pixel+strip pairs	inside high $p_t$ jets
9	MuonSeededInOut	muon-tagged tracks	muons
10	MuonSeededOutIn	muon detectors	muons

to recover these electrons with ECAL based algorithm only. The ECAL based and tracker based electron seeding is merged into one collection, which significantly improves the reconstruction efficiency.

Calorimeters are crucial components for the PF algorithm in CMS. The clustering algorithm in calorimeters is used to identify neutral stable particles like phone and neutral hadron, together with tracker in the identification of charged particles, reconstructing the energy of electrons and the possible associated bremsstrahlung photons and measuring the energy of charge particles that are missed by the

tracker.

The clustering algorithm is perform separately in each sub-detector system beside the HF in which each cell directly raise a cluster. The algorithm starts by finding a cluster seed. Then a topological walking around the neighbouring cells is performed. Both seed cells and neighbouring cells are required to pass certain thresholds to construct high quality candidates and suppress the contribution from noise. In ECAL endcaps, additional requirements on  $E_T$  because of the high noise level. In each of the topological clusters, the energy for the neighbouring cells are assumed from the seed cells. A Gaussian-mixture model is used in the construction of the topological clusters to evaluate the contribution of each cells. The finally parameters of the model is obtained by analytical fitting to the Gaussian model which give the maximum expectation.

To accurately measure the energy of particles like photons and neutral hadrons, the calibration of calorimeters is indispensable. The calibration of calorimeter also affects identification efficiency and misidentification rate of the particles measured in the calorimeters. The calibration of ECAL is done with a couple source, like the test beam, the radioactive source and the cosmic ray measurements and refined with the collision data. There are thresholds used in the formation of topological clusters, which results in the energy measured is smaller than the incoming particle energy. A residual energy calibration is applied to all ECAL clusters to account for these affects with simulation samples. The correction is applied as a function cluster energy and position,  $f(E, \eta) = g(E)h(\eta)$ . In the endcaps, the energy is taken as a liner combination of ECAL and preshower energy, the parameters in the combination is optimized by the  $\chi^2$  method. Hadrons generally leave energy in both ECAL and HCAL. The calibration of ECAL mentioned above is for the

photon, electron calibration. The behavior of hadron is different and a consequent calibration involving both ECAL and HCAL is needed. Simulated single neutral hadrons is used. The relationship involving both the energy measured by ECAL and HCAL and the calibrated energy  $E_{calib}$  is expressed as

$$E_{calib} = a + b(E)f(\eta)E_{ECAL} + c(E)g(\eta)E_{HCAL}$$

In the equation,  $a$  is independent of  $E$  and accounts for the effects of thresholds in the clustering algorithm. The other coefficients are determined by the  $\chi^2$  optimization with the  $E_{calib}$  and true energy  $E$  from simulation.

A detail description of particle flow algorithm can be found in [64].




Figure 4.1. A sketch view of CMS detector. Examples are given to show how the particles interact with different sub-detectors.

#### 4.2.2 Muon reconstruction and selection criteria

PF muon is used in the  $H \rightarrow \mu\tau_h$  analysis. In CMS, the event reconstruction starts from building the tracks in tracker(tracker track) and muon system(standalone-muon track) separately. Global muon reconstruction and tracker muon reconstruction are based on these tracks. [23]. Global muon reconstruction starts from the standalone-muon tracks and requires at least two muon station in the muon system. For each of the standalone-muon track, a propagating is done to find a matching tracker track on the common surface. Kalman-filter technique [41] is used in the fitting to combine the hits in standalone-muon track and tracker track. The tracker muon starts from the tracks with  $p_T > 0.5\text{GeV}$  and total momentum  $p > 2.5\text{ GeV}$ . The tracks are then extrapolated to match the tracks in the muon system with at least one muon segment. Within the geometry acceptance of muon system, the muon reconstruction efficiency is high, especially high momentum muons, constructed as either global muon or tracker muon or both has the efficiency about 99%. PF reconstruction utilizes the whole detector information to identify and reconstruct all of the particle. The muon PF algorithm applies a series of selections to the global muon and tracker muon to select out the PF muon. The selection is optimized to identify the muons in the jets with high efficiency and low misidentify rate. The details of the selection is in [1].

With the PF muon as the input, muons used in the analysis are further categorized into different identification, isolation categories. Muon ID suggested in CMS RunII, for the LHC data 2016, running period BCDEF, ICHEP medium muon ID is applied(table 4.5), for running period G and H, also the monte Carlo samples, standard medium muon ID(table 4.6)are applied to achieve the best performance for muon identification.

TABLE 4.5

Muon ID used in the analysis, for the LHC data 2016, running period BCDEF.

ICHEP mediumID description	Technical description
Loose muon ID	PFLoose Muon
Fraction of valid tracker hits	$> 0.49$
	Global muon
	Normalized global-track $\chi^2 < 3$
1.Good Global muon	Tracker-Standalone position match $< 12$
	kick finder $< 20$
	Segment compatibility $> 0.303$
2. Tight segment compatibility	Segment compatibility $> 0.451$

TABLE 4.6

Muon ID used in the analysis, for the LHC data 2016, running period G and H, also the monte Carlo samples.

Standard mediumID description	Technical description
Loose muon ID	PFLoose Muon
Fraction of valid tracker hits	$> 0.8$
	Global muon
	Normalized global-track $\chi^2 < 3$
1. Good Global muon	Tracker-Standalone position match $< 12$
	kick finder $< 20$
	Segment compatibility $> 0.303$
2. Tight segment compatibility	Segment compatibility $> 0.451$

#### 4.2.3 Electron identification

Electrons in CMS is constructed with the information from tracker and calorimeters. One of the main difficulties is the bremsstrahlung emitted by the electron during the traveling among the detector materials. The conversion of the photons from the bremsstrahlung affect the reconstruction of tracks in the track and these photons also cause significance energy loose in the electron reconstruction. The construction of Electron is discussed in section 4.2.1.1 and more details can be found in [50].

Electron ID is constructed to separate Prompt isolated electron(signal) from the background processes. The background can be the electrons from photon

conversion, from quark semi-leptonic decay and from misidentification of other particles or jets. The variables used in the identification are related to tracker and ECAL. There are mainly three types of variables. The variable related only to calorimeters. For example, the cluster shape of real electron in ECAL is usually narrower than the shape from hardronic showers and electrons leave most of the energy in ECAL and the energy ratio between ECAL and HCAL is large. The variables related to the matching of measured energy and geometry between tracker and ECAL. The variables related to tracker fitting to explore the difference between electrons and hadrons. These related variables can be used to construct cut-based selection sequence to selection electron. To achieve better performance, MVA based ID with boosted decision tree is also trained. Compared with the cut-based selection, more variables in the three categories mentioned above are used in the training [50]. An example of the BDT electron ID is shown in Figure. 4.2

Electron isolation is used to reject background events in addition to the ID variables, also inverting the requirement in the isolation can be used to setup enriched background control regions. Large numbers of background events that can possible enter the signal selection are misidentified jets or the jets in which there are real electrons, for example the jets from b quark semi-leptonic decay. For these background events, one key different character with respect to signals is more energy flowing around the electron(or misidentified electron) trajectory. The isolation requirement used in HLT level is summing over the energy depositions either in ECAL or HCAL in the certain core, for example  $\Delta R = 0.3, 0.4$ ,  $\Delta R = \sqrt{\Delta\phi^2 + \Delta\eta^2}$ . The contribution from the particle candidate is removed. In the offline algorithm, particles can better identified with the PF algorithm. Similar in concept to the isolation using the energy flow, the PF isolation sum-




Figure 4.2. Electron BDT-based ID shows good discriminating power against background in both EB and EE [50]

ming over the  $p_T$  of the particles in the direction of the reconstructed candidate trajectory momentum. PF algorithm utilizes the whole detector information and the isolation is defined as

$$\text{Iso}_{\text{PF}} = \sum p_T^{\text{charged}} + \max \left[ 0, \sum p_T^{\text{neutral had}} + \sum p_T^\gamma - p_T^{\text{PU}} \right]$$

The isolation variable sums over the contribution from charged PF candidates, neutral particles and photons in a certain selected  $\Delta R$  region around the signal. The contribution from pileup is estimated with  $p_T^{\text{PU}}$  with the FASTJET technique [16]. The energy of PF objects are better calibrated and the double counting problem which shows up the energy flow isolation described above is solved. PF isolation variable performs better than the isolation variable used in HLT. The performance comparison is shown in Figure. 4.3




Figure 4.3. PF electron isolation shows better performance in both EB and EE with respect to detector based isolation variable [50]

This should be used for B-tagging citation [65]

#### 4.2.4 Tau lepton reconstruction

In Run I CMS experiment, tau lepton are constructed with hadrons plus strips(HPS) algorithm. In general, HPS starts with PF jets which are reconstructed with  $anti - k_T$ , as the initial seeds.  $\pi_0$  components from the  $\tau$  hadronic decays are first constructed and combined with the charge hadrons parts, to identify different  $\tau$  decay modes and calculate  $\tau$  four-momentum and other quantities [27].

Photon conversions and the bremsstrahlung of electron/positron when traveling inside the CMS detector are well treated by the HPS algorithm. These phenomena broaden the signature of the tau decay. With PF jets as input, the algorithm constructs strips out of electromagnetic particles and starts by taking the strip in which contains the most energetic electromagnetic particle as the cen-

ter one. With the center strip, a window of the size  $\Delta\eta = 0.05$  and  $\Delta\phi = 0.2$  is taken. Within this window, if other charged particles are found, they are associated with the strip. The position of the strip is taken and four momentum of the strip is calculated. This procedure is repeated, until no strips can be constructed. The selected strips are required to have  $P_T^{strip} > 1GeV$ . The following decay topologies are taking into account by HPS:

- one charged particle without any strip,  $h^\pm$  and the case when  $\pi^0$  is not energetic enough to form a strip
- one charged particle plus one strip
- one charged particle plus two strips
- three charged partibles.

All of the charged hadrons and strips are required to be contained in the  $\Delta R = 2.8/P_T^{\tau_h}$  core, where the  $P_T^{\tau_h}$  is the reconstructed  $\tau_h$  transverse momentum and  $\Delta R$  is defined as  $\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$ . The  $\tau_h$  candidate is also required to match the direction of the seed PF jet within  $\Delta R = 0.1$ . Assuming all of the charged hadrons to be pions and taking in the associated strips, the HPS algorithm requires that different decay topologies meet the intermediate meson mass as listed in Table. 4.7.

The cut based  $\tau_h$  isolation discriminant required that the PF charged particles and photons to be considered in the isolation variable have  $p_T > 0.5$  GeV and within an isolation cone  $\Delta R = 0.5$  in  $\tau_h$  direction. The particles that constituent  $\tau_h$  are excluded from the summation. The effect of charged particle from pileup is eliminated by considering on the charged particle oriented from the  $\tau_h$  production vertex with in  $D_z = 0.2$  cm and  $\Delta r = 0.03$  cm. The effect of pileup on the isolation

of the photons on the strips is estimated by summing the charged particles that are not oriented from  $\tau_h$  decay primary vertex, within  $\Delta R = 0.8$  cm in the direction of and have the impact parameter  $D_z > 0.2$  cm. Then a factor  $\Delta\beta$  is multiplied to the  $p_T$  sum. The isolation variable is defined as in Equation. 4.1.

$$I_\tau = \sum p_T^{\text{charged}}(d_z < 0.2 \text{ cm}) + \max(0, \sum p_T^\gamma - \Delta\beta \sum p_T^{\text{charged}}(d_z > 0.2 \text{ cm})) \quad (4.1)$$

Tight, loose, medium working points(WP) for the tau isolation discriminants. The exact of the energy selection is suggested by the study of QCD dijet events, by requiring the  $I_\tau$  in Equation. eq:taucutis01 to have different values. The loose cut brings in approximate 1% of fake  $\tau$  from jets.

In CMS Run II, Tau reconstruction algorithm HPS has been improved [29]. The major improvement lies in Dynamic strip instead of fix size strip. Tau decay products can also affect the isolation. Charged pions in tau decay products experience nuclear interaction with tracker materials, which can results in low  $P_T$  secondary particles. Photons from the neutral pion decay can also go through pair production into  $e^+e^-$ , which further spread because of bremsstrahlung and the magnetic field. Broadening the strip is need in these cases in order to better cover the tau decay production. On the other hand, if the tau is boosted, high  $P_T$  decay products tends to be more concentrate and smaller strip size will be better. Similar to RunI tau reconstruction, the algorithm starts with hightest  $p_T$  charged particle as seeds for the strip. Starting from the seed strip, a window in  $\eta$  and  $\phi$  direction is set.

TABLE 4.7

Dominant hadronic  $\tau$  lepton decays branching fractions and the associated intermediate resonance. The h stands for both  $\pi$  and K. The table is symmetric under charge conjugation.

Decay mode	Resonance	Mass ( $MeV/c^2$ )	Branching fraction(%)
$\tau^- \rightarrow h^- v_\tau$			11.6%
$\tau^- \rightarrow h^- \pi^0 v_\tau$	$\rho^-$	770	26.0%
$\tau^- \rightarrow h^- \pi^0 \pi^0 v_\tau$	$\alpha_1^-$	1200	9.5%
$\tau^- \rightarrow h^- h^+ h^- v_\tau$	$\alpha_1^-$	1200	9.8%
$\tau^- \rightarrow h^- h^+ h^- \pi^0 v_\tau$			4.8%

$$\delta\eta = f(P_T^\gamma) + f(P_T^{strip}) \quad f(P_T) = 0.2 \cdot P_T^{-0.66}$$

$$\delta\phi = g(P_T^\gamma) + g(P_T^{strip}) \quad g(P_T) = 0.35 \cdot P_T^{-0.71}$$

The window is determined from single  $\tau$  gun MC simulation. 95% of the decay product will be covered in that range. The upward and downward limits for  $\eta$  is 0.15 and 0.05, for  $\phi$  the range is 0.3 and 0.05. The position of strip is set as  $p_T$  weighted average against all of the objects.

$$\eta_{strip} = \frac{1}{P_T^{strip}} \cdot \sum P_T^\gamma \cdot \eta_\gamma$$

$$\phi_{strip} = \frac{1}{P_T^{strip}} \cdot \sum P_T^\gamma \cdot \phi_\gamma$$

Construct the strip until no seed strip can be found. After the construction of the  $\tau$  lepton, for different decay mode,  $m_\tau$  is required to lie in different mass windows [30]. The conditions of different hadronic decay mode mass window are listed in the Table. 4.8. With respect to RunI, the difference in mass window is  $\delta m$ , which originates from dynamic clustering.  $\delta m$  is calculated as:

$$\delta m = \sqrt{\left(\frac{\partial m_\tau}{\partial \eta_{strip}} \cdot f(p_T^{strip})\right)^2 + \left(\frac{\partial m_\tau}{\partial \phi_{strip}} \cdot g(p_T^{strip})\right)^2}$$

with:

$$\frac{\partial m_\tau}{\partial \eta_{strip}} = \frac{P_z^{strip} \cdot E_\tau - E_{strip} \cdot P_z^\tau}{m_\tau}$$

$$\frac{\partial m_\tau}{\partial \phi_{strip}} = \frac{-(P_y^\tau - P_y^{strip}) \cdot P_x^{strip} + (P_x^\tau - P_x^{strip}) \cdot P_y^{strip}}{m_\tau}$$

In current algorithm,  $\tau^- \rightarrow h^- h^+ h^- v_\tau$  is not included, because of the jets contamination. This hadronic  $\tau$  decay mode composed of 4.8% of total branching fraction. The  $h^- \pi^0$  and  $h^- \pi^0 \pi^0$  are analyzed together, which is referred as  $h^- \pi^0$ .

TABLE 4.8

$\tau$  hadronic decay mode hypothesis signatures compatibility tests.  $m_\tau$  is required to be in the mass window

Decay mode	Mass window
$\tau^- \rightarrow h^-\pi^0 v_\tau$	$0.3 - \delta m_\tau < m_\tau < 1.3 \cdot \sqrt{p_T/100} + \delta m_\tau$
$\tau^- \rightarrow h^-\pi^0\pi^0 v_\tau$	$0.4 - \delta m_\tau < m_\tau < 1.2 \cdot \sqrt{p_T/100} + \delta m_\tau$
$\tau^- \rightarrow h^-h^+h^- v_\tau$	$0.8 - \delta m_\tau < m_\tau < 1.5 + \delta m_\tau$

The analysis with 2016 datasets, MVA based  $\tau$  isolation criteria is used, which keeps high identification efficiency while maintains relatively low fake rate compared with cut based discriminator. A Boosted Decision Tree(BDT) has been used in the training of the isolation variable. With BDT, the isolation variable shows a good distinguishing power against jets. Various variables have been used as BDT inputs. The variables are isolation variable( $I_\tau$ ), impact parameter from highest  $p_T$  track of  $\tau_h$  candidate,  $\tau_h$  decay mode information, shape variables like  $\Delta R$ ,  $\Delta\eta$ ,  $\tau$ -lifetime information and photon electron multiplicity, more of the exact variables used are discussed in [29, 30]. BDT uses these variables to distinguishing  $\tau_h$  dacay( $H \rightarrow \tau\tau$ ) from jets, which can be the decay products from quarks and gluons(QCD MC).

BDT method also used in the tau discriminating again electron training. The algorithm utilizes the variables that sensitive to the energy deposit in Ecal and Hcal, the electron bremsstrahlung, overall particle multiplicity and difference in electromagnetic and hadronic showers. Detail list of variable can be find in [29, 30].

Tau signals can be faked by muons, especially in  $\tau_h$  decay mode  $h^\pm$ . Tau against muon cut based muon discriminant is set by checking if there are signals in the muon system within  $\Delta R = 0.3$  of the  $\tau_h$  direction or if the energy sum from Ecal and Hcal is less than 20% of the total  $\tau$  energy. If less than two hits are found in the muon system, then it passes the loose working point. If no hits are found in the muon system, then this is the tight working point.

#### 4.2.5 Jet reconstruction

Jets are produced in the hadronic processes involving quarks or gluons. In p-p collision, jets involving in most of the processes. Understanding and constructing jets is crucial and difficult.

Anti- $k_t$  jet clustering algorithm [15] is used in CMS for the jet reconstruction. The algorithm starts by introducing the distance  $d_{ij}$  and  $d_{iB}$  as following,

$$d_{ij} = \min(k_{ti}^{2p}, k_{tj}^{2p}) \frac{\Delta_{ij}^2}{R^2}$$

$$d_{iB} = k_{ti}^{2p}$$

$d_{ij}$  is the distance between entities, the particle and the pseudojet.  $\Delta_{ij}$  is the difference of rapidity and azimuth between entry i and entry j.  $\Delta_{ij}^2 = (\eta_i - \eta_j)^2 + (\phi_i - \phi_j)^2$ . R is a radius parameter.  $k_{ti}$  and  $k_{tj}$  stands for the momentum of the entries respectively. p is a parameter used to specify jet construction algorithms. For Anti- $k_t$ , p=-1.  $d_{iB}$  is the distance between entry i and the beam. If the parameter p is set for other values, for example p=1 or 0, then the algorithm is the  $k_t$  [39] or Cambridge/Aachen jet reconstruction algorithm [68]. With the Anti- $k_t$ , assuming in one event, there are a couple hard particle with high momentum,

$k_{t1}$ ,  $k_{t2}$  and so on, also large numbers of soft ones around. Starting from  $k_{t1}$  as an example, if  $d_{1j}$  is smaller, then entry j combined with entry 1, if  $d_{1B}$  is smaller, then entry 1 is set as a jet and removed from the list. The soft entries tend to ground around the high momentum entries to the range  $\Delta_{ij} = R$ , if there is no other hard entry around. If two hard entries are within the distance R, then the two entries are clustered into a single entry. In this case, if  $k_1 \gg k_2$ , then the center is more closed to jet 1. If  $k_1 \sim k_2$ , the boundary between two jets is defined by  $\Delta R_{1b}/k_{t1} = \Delta_{2b}/k_{t2}$ . Jet clustering continues until all of the jets are clustered in the events.

Jet energy is corrected to have the correct energy scale. The correction goes through a couple of steps [52]. The major corrections are derived from simulated samples and the residual corrections from the different response between MC and data are from data-driven methods.

Pileup events can increase the measured jet energy, especially in LHC Run II, the number of pileup per event doubled. Two types of pileup affect the performance the most, in-time pileup(IT PU) and out-of time pileup(OOT PU). The IT PU refers to the additional events produced by the proton-proton collision within the same bunch-crossing as the primary hard collision. The OOT PU refers to the events that produced in previous bunch crossing or subsequent one that affect current bunch. OOT PU can be mitigated by explore the timing window, pulse shape of the calorimeter. The IT PU mitigation is mainly discussed. Charged hadrons from IT PU in CMS are removed with Charged-hadron subtraction(CHS) algorithm in CMS. Tracks with the vertexes that are identified from PU with charged particles inside are removed. CHS removes around 50% of the IT PU within the tracker coverage region. There are also soft jets from pileup

interaction. These jets are usually in low energy range and affects the JES by overlapping with the hard jets. Multivariate analysis(MVA) with inputs from jet shape and jet constitution information can more than 90% of pileup jets. This MVA ID is referred as PUJetID [3]. After rejecting the charged particle jets and soft jets from pileup, a jet area method [16] is used to further eliminate the effects from PU. In this method, an estimated energy density bringing in by the PU and effective area of jet is used to calculate the offset energy. After the PU correction, simulated response corrections are performed with simulated QCD multijet sample. A matching between particle-level jets and reconstructed jets is performed. The correction is performed with the anti- $k_T$  jets with distance parameter R=0.5 in  $p_T$  and  $\eta$  distribution and checked with jets corresponding to R in range 0.3 to 1.0. Following the simulated response corrections, the remaining residual correction is small derived with data-based methods. A comparison between data and MC samples and corrects data only.

B jets is a main component in  $t\bar{t}$  process, so the identification of b jets is important to the analysis involves  $t\bar{t}$ . In CMS, combined secondary vertex version 2(CMSv2) currently is used, which uses multivariable techniques and the character of b jets, to identify b jets [65]. B jets is from the hadronization of b quark. B hadron is relatively heavy, around 5-6 GeV, with high decay multiplicity, relatively high momentum decay products and relatively wide decay core. B hardon has long lifetime( $ct \approx 450\text{um}$ ) which give a traveling distant  $\approx 5\text{ mm}$  at the energy of 70 GeV. Exploring the property of secondary vertex is used in the identification algorithm. In each jet, at least two tracks presenting in an angular distance  $\Delta R < 0.3$  is required. The jets with combined mass of two tracks compatible with  $k_s^0$  is rejected to reduce the contamination. A number of variables [65] are used as

input into the MVA training in different secondary vertices presenting situation. Combining the input variables into a discriminant variable to identify b jets.

#### 4.2.6 Missing transverse momentum

Missing transverse momentum(MET) is defined as the sum  $p_T$  of un-detected process in an event. In CMS, particle flow MET is widely used and the MET can be expressed as  $\vec{E}_T = -\sum \vec{p}_T$ , in which the summing is over the observed PF particles. The magnitude of MET can be affected by various resources, for example the inefficiency of tracker, the thresholds on calorimeter and the asymmetry in detector response. The propagation of JEC to MET can reduce the bias. The corrected MET can be expressed as

$$\vec{E}_T^{corr} = \vec{E}_T - \sum_{jets} (\vec{p}_{T,jet}^{corr} - \vec{p}_{T,jet})$$

The sum is summing over the jets with electromagnetic fraction below 0.9 and corrected  $p_T > 10$  GeV. Further correction aiming for estimation is the effects of PU on MET is derived with charged particles associated with PU vertexes,  $\vec{v} = \sum_{charged} \vec{p}_T$ . The correction is  $f(\vec{v})\vec{v}$ , in which factor  $f(\vec{v}) = c_1(1.0 + \text{erf}(-c_2|\vec{c}|^{c_3}))$ . The coefficients in  $f(\vec{v})$  are estimated from fitting the MC minimum bias events. The correction on MET for the effect of PU is

$$\vec{E}_T^{corr} = \vec{E}_T - \sum_{PU} f(\vec{v})\vec{v}$$

The observed  $\vec{E}_T$  asymmetry in  $\phi$  shows approximately linear relationship with the number of reconstructed vertices. The correction is performed with  $N_{vtx}$

separately in x and y axis.

$M_{col}$  is the most important variable in LFV Higgs decay analysis, which is closely related to MET. Higgs is more massive than the leptons( $e, \mu, \tau$ ), so the decay products are boosted. In  $H \rightarrow \mu\tau_h$  and  $H \rightarrow e\tau_h$  analysis,  $\tau$  decays hadronically. The neutrinos in  $\tau$  decay products are highly Lorentz boosted and are considered collinear with  $\tau$ . The direction of  $\tau$  and the projection of MET amplitude on  $\vec{\tau}$  direction are used to approximate the direction and energy of the neutrinos. With this approximation and the visible mass which is the mass formed with  $e-\tau$  or  $\mu-\tau$  without the approximation,  $M_{col}$  can be derived as

$$M_{col} = M_{vis} / \sqrt{x_\tau^{vis}}$$

$$x_\tau^{vis} = p_T^{\vec{\tau}^{vis}} / (p_T^{\vec{\tau}^{vis}} + p_T^{\nu,est})$$

### 4.3 Event simulation

Monte Carlo simulation is used widely from the detector related processes, event reconstruction to physics analysis etc. The generation of MC sample in the generator involving the following steps.

In LHC, p-p collision, QCD hard subprocesses involving large momentum transferring, multiply final states and complex surrounding environment. For a particular process, MC generator starts by calculating the related QCD cross section. This involving calculating the matrix-element, choosing parton distribution functions(PDFs) and possibly performing high order correction. Followed by the parton showers. In a hard process, parton showers uses the basic building block like  $q \rightarrow qg$ ,  $q \rightarrow gq$ ,  $g \rightarrow gg$  and  $g \rightarrow q\bar{q}$ , iteratively showering from high energy

to low energy, typically, for quark and gluon in the order of 1 GeV, when they can not be treated as free particles anymore. In addition to the hard subprocesses and the associated parton showers, a real event consists more additional contributions. Typically more than one pair of partons interacts in one collision. These multiple interactions are dealt with different models and affects the total scattering energy, the number of the particle in the hadronization stage, etc. Because of the color confinement, quarks and gluons forms hadrons after completing the parton shower, this is referred as hadronization. Similarly to the QCD processes, electromagnetic radiation is also taken into account with the evolution according to the electric charge instead of color. With the generated events through the MC generate, the events are put into GEANT4 [33]

## CHAPTER 5

### LFV event selection

In both 8 TeV  $H \rightarrow e\tau_h$  and 13 TeV  $H \rightarrow \mu\tau_h$  analyses, events are selected in several steps. A loose selection selects on the different IDs, energy, geometry parameters of the analysis related objects and followed by a tighter set of selection criteria in which selection requirements are placed on the kinematics variables and fits on variable  $M_{col}$  in both  $H \rightarrow e\tau_h$  and  $H \rightarrow \mu\tau_h$  analysis. This selection sequence is referred as  $M_{col}$  fit analysis. In  $H \rightarrow \mu\tau_h$ , there is another selection sequence. After placing the loose selection, a multivariate analysis with Boosted decision tree (BDT) is exploited and fits on the BDT discriminator. This is referred as the BDT fit analysis and provides more sensitive results in  $H \rightarrow \mu\tau_h$  analysis.

## 5.1 $H \rightarrow \mu\tau_h$

### 5.1.1 Loose selection

Tau leptons from signal events decay hadronically. SM higgs is much heavier than its decay products  $\mu$  and  $\tau$ .  $\mu$  and  $\tau$  leptons are expected to have high  $P_T$ . The decay products from signal events are boosted, a cut on  $\Delta R = \sqrt{(\Delta phi)^2 + (\Delta eta)^2}$ ,  $\Delta R > 0.3$  is applied.  $\mu$  and  $\tau$  candidates are required to have opposite sign of charges as Higgs has no electric charge. Further, events with additional  $\mu$  and

$\tau$  that pass a loose selection, events with jets that are identified by the combined secondary vertex(CSVv2) b-tagging algorithm [31] as a b quark jets are discarded. The trigger HLT\_IsoMu24 or HLT\_IsoTkMu24 used in the analysis selects isolated muons that have energy higher than 24 GeV at HLT level. An further  $P_T$  cut on reconstructed  $\mu$ ,  $P_T > 26$  GeV and  $|\eta| < 2.4$  are required. Muons are required to pass the recommended Medium muon ID and tight cut based isolation  $I_{rel}^\mu < 0.15$ .

Hadronic taus are required to have  $P_T > 30$  GeV,  $|\eta| < 2.3$ , passing old tau decay mode finding, the MVA based tight tau isolation ID and tau discriminators against electrons and muons.

Events in the analysis are divided into four categories based on the number of jets. In 2-jets category, it is furthered divided into 2 categories, 2-jet gluon gluon fusion higgs production(ggH) category and 2-jet vector boson fusion(VBF) category based on the value of 2 jets invariant mass( $M_{jj}$ ) . The 0-jet category enhances ggH production mode. In 1-jet category, the dominant signal production mode is also ggH, but with a boosted jet associated with the production and VBF higgs production also contributes this category. In the 2-jet ggH category, signal events mainly come from ggH, while in 2-jet VBF, VBF higgs is the dominant production channel. The following is a more detailed list of the selection condition in each categories.

**0-jet:** No events have jets pass the loose PF ID and with jet  $P_T > 30$  GeV,  $|\eta| < 4.7$ .

**1-jet:** Events with one jet passes losse PF ID and jet  $P_T > 30$  GeV,  $|\eta| < 4.7$ .

**2-jets ggH:** Events have two jets passing loose PF ID,  $P_T > 30$  GeV,  $|\eta| < 4.7$  and a requirement on the invariant mass of the two jets,  $M_{jj} < 550$  GeV.

**2 jets VBF:** Events with two jets pass loose PF ID. Jets  $P_T > 30$  GeV,  $|\eta| < 4.7$  and

$M_{jj} > 550$  GeV are required.

The threshold on  $M_{jj}$  has been optimized to give the best expected exclusion limits.

### 5.1.2 Cut-based analysis

After the loose selection and the categorization, a further cut-based selection is applied. Variables that can help distinguish signal from background are  $P_T^\mu$ ,  $P_T^\tau$  and  $M_T(\tau_h)$ . The lepton  $P_T$  variables are very powerful background discriminant variables, but it will also cause the problem that signal picks under backgrounds. Leptons from signal process incline to have higher  $P_T$  values, by cutting tighter on the lepton  $P_T$ , more background events can be removed. However this will also reshape some of the backgrounds, making them peak closer under the signal so that signal processes will be affected more by the background statistics fluctuation. In the  $H \rightarrow e\tau_h$  analysis, the effect of cutting tighter on lepton  $P_T$  will be shown. In  $H \rightarrow \mu\tau_h$  search, lepton  $P_T$  variables are kept at loose values and optimizes on other variables to achieve tighter expected limits.

Optimization is proceeded with only Monte Carlo samples so as not to double use data. In  $H \rightarrow \mu\tau_h$  channel, the variables tuned are  $M_{jj}$  and  $M_T(\tau_h)$ . Cuts have been optimized to have the most stringent expected limits with the Asimov dataset. If looser values of the cuts give the same expected limits as the tighter ones, then the looser cut values are chosen to have more statistics. Examples of optimization by checking the limits are shown in Figure. 5.1 and 5.2, in which the optimization of  $M_T(\tau_h)$  and  $M_{jj}$  are shown. A full summary of the cuts optimized for  $H \rightarrow \mu\tau_h$  is shown in Table. 5.1




Figure 5.1. Expected limits based on an Asimov dataset as a function of  $M_T(\tau, \text{MET})$  for the different categories.

TABLE 5.1

Selection criteria in each category with the optimization of the  $H \rightarrow \mu\tau_h$  analysis

<b>0-jet category</b>
<ul style="list-style-type: none"> <li>• <math>p_T^\mu &gt; 26</math> GeV, <math>p_T^\tau &gt; 30</math> GeV</li> <li>• <math>M_T(\tau) &lt; 105</math> GeV</li> <li>• No jets with <math>p_T^{jet} &gt; 30</math> GeV, <math> \eta  &lt; 4.7</math>, LooseID</li> </ul>
<b>1-jet category</b>
<ul style="list-style-type: none"> <li>• <math>p_T^\mu &gt; 26</math> GeV, <math>p_T^\tau &gt; 30</math> GeV</li> <li>• <math>M_T(\tau) &lt; 105</math> GeV</li> <li>• One jet with <math>p_T^{jet} &gt; 30</math> GeV, <math> \eta  &lt; 4.7</math>, LooseID</li> </ul>
<b>2-jet, gg-enriched category</b>
<ul style="list-style-type: none"> <li>• <math>p_T^\mu &gt; 26</math> GeV, <math>p_T^\tau &gt; 30</math> GeV</li> <li>• <math>M_T(\tau) &lt; 105</math> GeV</li> <li>• <math>p_T^{jet1} &gt; 30</math> GeV, <math>p_T^{jet2} &gt; 30</math> GeV <math> \eta_{jet1}  &lt; 4.7,  \eta_{jet2}  &lt; 4.7</math>, LooseID</li> <li>• <math>M_{jj} &lt; 550</math> GeV</li> <li>• Two jets with <math>p_T^{jet} &gt; 30</math> GeV, <math> \eta  &lt; 4.7</math>, LooseID</li> </ul>
<b>2-jet, vbf-enriched category</b>
<ul style="list-style-type: none"> <li>• <math>p_T^\mu &gt; 26</math> GeV, <math>p_T^\tau &gt; 30</math> GeV</li> <li>• <math>M_T(\tau) &lt; 85</math> GeV</li> <li>• <math>p_T^{jet1} &gt; 30</math> GeV, <math>p_T^{jet2} &gt; 30</math> GeV <math> \eta_{jet1}  &lt; 4.7,  \eta_{jet2}  &lt; 4.7</math>, LooseID</li> <li>• <math>M_{jj} &gt; 550</math> GeV</li> <li>• Two jets with <math>p_T^{jet} &gt; 30</math> GeV, <math> \eta  &lt; 4.7</math>, LooseID</li> </ul>




Figure 5.2. Expected limits based on an Asimov dataset as a function of  $M_{jj}$  for the 2 jet categories.

### 5.1.3 Multivariate analysis

A Boost decision trees(BDT) method is used as the multivariate analysis method in  $H \rightarrow \mu\tau_h$  search which is more sensitive than the  $M_{col}$  fit analysis. BDT algorithm used in this search is implemented in the TMVA package [47]. BDT takes in signal and background datasets with a selected set of input variables. Input variables are the ones that show distinguishing power between signal and backgrounds. The training output is a weight file, which contains a list of weights to indicate how likely an event is signal like with a give set of input variables. A more detail description of the BDT method is available in section 7.1. In this analysis, signal and background events are required to pass the loose selection criteria. All of the categories are combined. The signal events from ggH and VBF higgs production mode are mixed by the weight with respect to their production cross section. The background sample used in the training is misidentified lepton background from the like sign region(Region II as in Table. 6.1). The list of BDT input variables is in the following list and the distribution of the variables are

shown in Fig. 5.3.

- Transverse mass between the  $\tau_h$  and  $E_T^{miss}$ ,  $M_T(\tau_h)$ .
- Missing transverse energy,  $E_T^{miss}$ .
- Pseudorapidity difference between the  $\mu$  and the  $\tau_h$  candidate,  $\Delta\eta(\mu, \tau_h)$ .
- Azimuthal angle between the  $\mu$  and the  $\tau_h$ ,  $\Delta\phi(\mu, \tau_h)$ .
- Azimuthal angle between the  $\tau_h$  and the  $E_T^{miss}$ ,  $\Delta\phi(\tau_h, E_T^{miss})$ .
- Collinear mass,  $M_{col}$ .
- Muon  $p_T$ .
- $\tau_h p_T$ .

The chosen input variables show low correlation in both samples as shown in Figure. 5.4. The TMVA overtraining checks is performed by checking with the testing samples to see if similar distinguishing power can be achieved with the output weight file from the training samples. Training and testing samples show good agreement and the training process exempts from overtraining as indicated in Figure. 5.5.

## 5.2 $H \rightarrow e\tau_h$

### 5.2.1 Loose selection

In  $H \rightarrow e\tau_h$  channel, trigger *HLT\_Ele27\_WP80* is used, which applies an electron  $p_T$  cut at 27 GeV at HLT level. A further cut on electron  $P_T > 30$  GeV is applied. Electrons are also required to have  $|\eta_e| < 2.3$  and  $D_z < 0.2cm$ .




Figure 5.3. Distributions of the input variables to the BDT for the  $H \rightarrow \mu\tau_h$  channel.




Figure 5.4. Expected limits based on an Asimov dataset as a function of  $M_T(\tau, MET)$  for the different categories.




Figure 5.5. Overtraining checking for the BDT training in the TMVA package.

$D_z$  is the longitudinal impact parameter that shows the displacement between primary vertex and track path. Electrons are required to pass the MVA based tight ID and cut based PF tight isolation  $I_{rel}^e < 0.1$ . Tau candidates are required to have  $P_T > 30$  GeV, pseudorapidity  $|\eta^\tau| < 2.3$  and the longitudinal impact parameter  $D_z < 0.2$  cm. Tau isolation used is the cut based tight tau isolation. In addition, tau candidates are required to pass the tau Decay mode finding and tau discriminator against electrons and muons. The analysis also requires no extra isolated electrons with  $p_T > 10$  GeV and extra taus with  $p_T > 20$  GeV. Tau and electron candidates are required to have opposite sign of charges and separate with  $\Delta R > 0.4$  from any jets in the events with  $p_T > 30$  GeV. All of the requirements contribute to the selections of good qualities candidates. The datasets are binned into three categories according to the number of jets in the events:

**0-jet:** No events have jets pass the loose PF ID and with jet  $P_T > 30$  GeV,  $|\eta| < 4.7$ .

This category enhances the gluon-gluon fusion contribution.

**1-jet:** Events with one jet passes loose PF ID and jet  $P_T > 30$  GeV ,  $|\eta| < 4.7$ .

This category enhances the gluon-gluon fusion production with initial state radiation.

**2 jets:** Events with two jets pass loose PF ID and with jet  $P_T > 30$  GeV and  $|\eta| < 4.7$ , This category contains both Higgs production mode and with an enhancement in VBF production mode.

With the preselection and binning of the jets numbers, the  $M_{col}$  distribution of  $H \rightarrow e\tau_h$  in different categories are shown in Figure. 5.6




Figure 5.6. With loose selection conditions, the comparison of the observed collinear mass distributions with background from prediction. The shaded grey bands indicate the total background uncertainty. The open histograms correspond to the expected signal distributions for  $\mathcal{B}(H \rightarrow e\tau_h) = 100\%$  in the 0-jet, 1-jet and 2-jet categories, respectively.

### 5.2.2 Cut-based analysis

A set of kinematics variables are used to further select signal events. In  $H \rightarrow e\tau_h$  channel, similar to the  $H \rightarrow \mu\tau_h$ , muon and tau leptons from the signal events are highly boosted so as  $p_T$  variables plays an important role in distinguishing from background events and the separation in  $\phi$  direction is bigger between  $\mu$  and  $\tau$  of signal events. The variable  $M_T(\tau_h)$  which is the transverse mass form by the reconstructed  $\tau$  and MET is also used in the selection. In two jets category, the invariant mass of two jets  $M_{jj}$  is used as a cut variable. The cuts have been optimized to have the most stringent expected limits with the Asimov dataset. The detailed cuts used for  $H \rightarrow e\tau_h$  is shown in table. 5.2.

TABLE 5.2

Selection criteria for each event category after cut optimization, for the  
 $H \rightarrow e\tau_h$  channel

---

<b>0-jet category</b>
<ul style="list-style-type: none"> <li>• <math>p_T^e &gt; 45</math> GeV, <math>p_T^\tau &gt; 30</math> GeV, <math>\Delta\phi_{e\tau} &gt; 2.3</math></li> <li>• <math>M_T(\tau) &lt; 70</math> GeV</li> <li>• No jets with <math>p_T^{jet} &gt; 30</math> GeV, <math> \eta  &lt; 4.7</math>, LooseID</li> </ul>
<b>1-jet category</b>
<ul style="list-style-type: none"> <li>• <math>p_T^e &gt; 35</math> GeV, <math>p_T^\tau &gt; 40</math> GeV</li> <li>• One jet with <math>p_T^{jet} &gt; 30</math> GeV, <math> \eta  &lt; 4.7</math>, LooseID</li> </ul>
<b>2-jet category</b>
<ul style="list-style-type: none"> <li>• <math>p_T^e &gt; 35</math> GeV, <math>p_T^\tau &gt; 30</math> GeV</li> <li>• <math>M_T(\tau) &lt; 50</math> GeV</li> <li>• <math>p_T^{jet1} &gt; 30</math> GeV, <math>p_T^{jet2} &gt; 30</math> GeV, <math> \eta_{jet1}  &lt; 4.7,  \eta_{jet2}  &lt; 4.7</math>, LooseID</li> <li>• <math>\Delta\eta(jet1, jet2) &gt; 2.3</math></li> <li>• <math>M_{jj} &gt; 400</math> GeV</li> <li>• Two jets with <math>p_T^{jet} &gt; 30</math> GeV, <math> \eta  &lt; 4.7</math>, LooseID</li> </ul>

---

## CHAPTER 6

### LFV analysis background estimation

#### 6.0.1 Background estimates for $H \rightarrow \mu\tau_h$

##### 6.0.1.1 Misidentified Leptons

The misidentified lepton backgrounds are estimated with full data driven method from the collision data. In  $H \rightarrow \mu\tau_h$  channel, the misidentification  $\tau$  and  $\mu$  lepton rate is obtained from independent  $Z + \text{jets}$  data sets and then apply the rate to a control region that orthogonal to the signal region to estimate the misidentified lepton backgrounds. The control regions for the estimation and validation of the misidentified lepton background are shown in Table. 6.1. Region I is the signal region with full selection applied. Region III is the background enriched region, in which one or two leptons have loose isolation but not tight isolation. Region II is the same to Region I besides the requirement that  $\mu$  and  $\tau$  lepton pairs have the same sign of charge, so as Region IV to Region III. Region III is used to estimate the misidentified background in Region I. Region II and IV are used in the validation of the method.

$Z \rightarrow \mu\mu$  is used in  $Z + \text{jets}$  data sets to estimate both misidentified  $\tau$  and  $\mu$  lepton rates.  $Z$  bosons are selected in the invariant mass window  $70 < M_{ll} < 110$  GeV and trigger HLT\_IsoMu24 or HLT\_IsoTkMu24 is used in the selection. Two muons that are composed of  $Z$  bosons are required to have  $p_T > 26$  GeV,  $|\eta| < 2.4$ ,

TABLE 6.1

Definition of the samples used to estimate the misidentified lepton background.

<b>Region I</b>	<b>Region II</b>
$\ell_1^\pm$ (isolated)	$\ell_1^\pm$ (isolated)
$\ell_2^\mp$ (isolated)	$\ell_2^\pm$ (isolated)
<b>Region III</b>	<b>Region IV</b>
$\ell_1^\pm$ (isolated)	$\ell_1^\pm$ (isolated)
$\ell_2^\mp$ (not-isolated )	$\ell_2^\pm$ (not-isolated)

cut based tight muon isolation( $I_{rel}^\mu < 0.15$ ) and passing the muon medium ID. In the  $Z+jets$  samples, with the selected  $Z$  boson in an event, the remaining jets are checked if they pass  $\tau$  or  $\mu$  selection.

To estimate the misidentified  $\tau$  lepton rate, The jets in the sample  $Z + jets$  are checked if they pass the  $\tau$  isolation ID. Misidentified  $\tau$  lepton ratio  $\tau(f_\tau)$  is calculated as in Equation . 6.1, together with one related ratio  $f_\tau$ . In Equation . 6.1,  $\ell$  stands for leptons in general.

$$\ell(f_\ell) = \frac{f_\ell}{1 - f_\ell} \quad (6.1)$$

$$f_\ell = \frac{N_\ell(Z + jets \ell \text{ tight Iso})}{N_\tau(Z + jets \ell \text{ loose Iso})} \quad (6.2)$$

$f_\tau$  is the ratio between number of jets that pass tight  $\tau$  MVA isolation ID and number of jets that pass very loose  $\tau$  MVA isolation ID. The contribution from Diboson sample is subtracted with simulation, in which the third leptons are genuine leptons. With the ratio  $f_\tau$  , misidentified rate  $\tau(f_\tau)$ (Equation. 6.2)

is applied as a weight to Region III. When estimating misidentified *tau* leptons, Region III is defined in the same selection condition as the signal region Region I , but  $\tau$  leptons are required to pass the very loose MVA isolation and not pass the tight MVA isolation. The misidentified  $\tau$  lepton rate  $\tau(f_\tau)$  shows dependence on tau  $p_T$ , tau decay mode and ECAL geometry. Thus  $\tau(f_\tau)$  is applied in term of tau  $p_T$ , tau decay modes and ECAL barrel and endcap separately. The distribution of  $f_\tau$  is shown in Fig. 6.1, which shows a comparison of the estimated  $f_\tau$  between data and simulated Z+jets samples. The distribution of  $f_\tau$  vs  $|\eta|$  are also shown, which show less dependency.

The estimation of  $\mu$  misidentified rate the same as the  $\tau$  misidentified rate, but checks if the jets in Z + jets samples pass the muon isolation. As in Equation. 6.1,  $f_\mu$  is defined as the ratio between tight cut based muon isolation( $I_{rel}^\mu < 0.15$ ) and loose cut based muon isolation( $I_{rel}^\mu < 0.25$ ). The misidentified  $\mu$  lepton rate  $\mu(f_\mu)$  is calculated by Equation. 6.2.  $\mu(f_\mu)$  is a weight to apply to Region III. In the case of estimating the misidentified  $\mu$ , Region III is defined as Region I but with  $\mu$  passing the loose cut based muon isolation and not passing tight cut based muon isolation. The ratio  $f_\mu$  is shown in Figure. 6.2 in term of muon  $p_T$  and  $|\eta|$  distribution.

There are overlap between the estimated misidentified muon and tau lepton when both of the leptons are fake ones. The overlap events are estimated as in Equation. 6.4. In the equation, the N(Region III) stands for the events in Region III. In this case, Region III is defined as muon leptons passing the loose isolation, but not passing the tight isolation, while tau leptons pass the very loose isolation but not pass the tight isolation. The final misidentified lepton background obtained by adding the misidentified muon and tau background and deducts the overlap




Figure 6.1.  $f_\tau$  ratio for the  $\tau$  fake rate calculation shows in term of tau decay modes,  $p_T$ , ECAL barrel, ECAL endcap, and  $|\eta|$




Figure 6.2.  $f_\tau$  ratio for the  $\mu$  fake rate calculation shows in term of muon  $p_T$  and  $|\eta|$ .

events between these two.

$$N(\text{overlap}) = \frac{f_\tau \cdot f_\tau}{(1 - f_\tau) \cdot (1 - f_\mu)} \cdot N(\text{Region III}) \quad (6.3)$$

(6.4)

In  $H \rightarrow \mu\tau_h$  channel, the validation of the misidentified lepton background with full data driven method is performed in a like-sign sample and W+jets enriched control region. The like-sign sample validation requires the events pass the loose selection, the same as the signal region, but with inverted charge. Misidentified background is enriched with the requirement of inverting charge. In the W+jets enriched control region, events that pass the loose selection are further selected with  $M_T(\tau) > 60$  GeV and  $M_T(\mu) > 80$  GeV. In both of the control regions, misidentified background is the dominant one, as shown in Fig. 6.3. With full data driven method, applied with the  $\tau(f_\tau)$  as weight, the like-sign and W+jets

control regions show good data and MC agreement.




Figure 6.3. The validation of full data driven method for the misidentified lepton background. 30% of uncertainty is assumed in both cases.

### 6.0.1.2 Other backgrounds

Besides the misidentified background, which is composed by  $W+jets$  and QCD multi-jets background and estimated with data driven method,  $Z \rightarrow \tau\tau$  and  $Z \rightarrow \mu\mu$  are also major background in this analysis. In  $Z \rightarrow \tau\tau$ , the muons arise from a  $\tau$  decay, while in  $Z \rightarrow \mu\mu$ , one of the  $\mu$  can be misidentified as a  $\tau$ . In other smaller backgrounds, the  $\mu\tau$  pairs can be produced from the weak decays of quarks and vector bosons. These smaller backgrounds include standard model Higgs production( $H \rightarrow \tau\tau, WW, t\bar{t}$ , single top quark, di-boson(WW,WZ,ZZ)). A full list

of background used is in the Table. ???. These backgrounds are all estimated with MC simulation.

In the  $Z \rightarrow \mu\mu$  background, an additional scale factor is applied to the tau leptons that do not come from real taus. This is checked in the MC generator level. The scale factor is applied in term of  $|\eta|$  as shown in Table. 6.2.

TABLE 6.2

Scale factor to the events in which a muon is misidentified as a tau

$ \eta $ ranges	Correction values
0-0.4	1.263
0.4-0.8	1.364
0.8-1.2	0.854
1.2-1.7	1.712
1.7-2.3	2.324

## 6.0.2 Background estimates for $H \rightarrow e\tau_h$

### 6.0.2.1 Misidentified Leptons

Misidentified lepton background is a very important one in  $H \rightarrow e\tau_h$  channel. This background mainly raises from W+jets and QCD multijets. A background

event passes the signal selection, may have one or both leptons are misidentified, which are faked by other objects. In  $H \rightarrow e\tau_h$  8 TeV analysis, both misidentified electron and tau leptons are considered.

Similar to  $H \rightarrow \mu\tau_h$  channel, the misidentified background in  $H \rightarrow e\tau_h$  is estimated with full data driven method, with independent data sets Z+jets, in which Z bosons decay into two electrons. Data sets are divided into four control regions for this estimation as in Table. 6.1. Region II is used to extra misidentified background in signal region, Region I, while Region II and Region IV are used for the validation of the fake rate. In the Z+jets data set selection, trigger HLT\_Ele27\_WP80 is used. The two electrons in Z+jets are required to have  $p_T > 30$  GeV,  $|\eta| < 2.3$ , tight electron MVA ID and cut based tight PF isolation( $I_{rel}^e < 0.1$ ). The invariant mass formed by the two electrons are required to be in the Z boson mass window. Then the jets in the events are checked if they pass  $\tau$  or e selection. Tau leptons should have  $p_T > 30$  GeV,  $|\eta| < 2.3$ , passing tau decay mode finder and discriminator against elections and muons. Electrons pass the same conditions as the elections used in forming the Z boson mass peak, but without the requirement of isolation. The isolation conditions are required in different conditions in the fake ratio estimation. Tau leptons use the cut based tight or loose isolation, while loose or tight MVA based electron isolation have the value  $I_{rel}^e < 0.1$  and  $I_{rel}^e < 0.2$  respectively. After the selection, the fake ratio  $f_\tau$  is defined as the number of tau leptons passing the tight isolation divided the number of tau leptons passing the loose isolation. Similarly the fake ratio  $f_e$  is defined as the number of elections passing the loose isolation divided by the number of electron passing the tight isolation as shown in Equation. 6.2. With the fake ratio  $f_\tau$  and  $f_e$ , the electron and tau fake rate  $\tau(f_\tau)$  and  $e(f_e)$  are estimated according to Equation. 6.1. The

fake ratio for tau leptons and electrons are shown in Figure. 6.4.

The tau and electron fake ratio is applied in term of  $|\eta|$ . In the estimation, the background source of three prompt leptons from Diboson samples are deducted. The tau and electron misidentified backgrounds are estimated separately, the overlap events in which both leptons are misidentified objects are estimated with Equation. 6.4. In the equation,  $N(\text{Region III})$  in  $H \rightarrow e\tau_h$  channel satisfies the condition both leptons are loose isolated but not tight isolated. The validation of full data driven method is done with Region II and Region IV. The distribution of the  $M_{col}$  is checked in the same sign region of the two leptons, shown in Figure. 6.5, in which observed data agrees well with the events from MC and misidentified background from data driven method.

#### 6.0.2.2 $Z \rightarrow \tau\tau$

$Z \rightarrow \tau\tau$  in another significant background in  $H \rightarrow e\tau_h$  channel. In the case one of the tau leptons decay to a electron and the other hadronically decay. The background is estimated with the embedding technique[26], which starts by selecting the  $Z \rightarrow \mu\mu$  events from collision data. The muons pass loose muon selection and then are replaced by simulated  $\tau$  leptons and reconstructed with PF algorithm. The key advantage of this technique is that the major topologies like missing energy, jet multiplicity, underlying events are directly from data. Besides tau, the other event records and detector responds are from real environment. When added to the background, the embedded data is normalized to MC prediction. Compared with MC prediction, the visible mass peak from embedding sample shifts 2%. This is mainly because of the difference in final-state radiation of photons between muon and tau and is corrected for. Identification and isolation




Figure 6.4.  $f_\tau$  and  $f_e$  ratio for the fake rate calculation shows in term of  $p_T$  and  $|\eta|$




Figure 6.5. Data driven method validation in Region II with  $e$  and  $\tau$  leptons in same sign of charge

correction for this shift to embedded sample are obtained by comparing with MC simulations.

#### 6.0.2.3 Other backgrounds

$t\bar{t}$  is one of the main background. Leptonic decay W bosons can produce opposite sign lepton pair and missing energy. The shape of this background is obtained from MC samples and normalized to  $t\bar{t}$  control region sample. The events in  $t\bar{t}$  control samples are required to pass the 2-jet base selection in  $H \rightarrow e\tau_h$  channel and at least one jet is identified as a b jet.

The other backgrounds presented in the samples are Z+jets in which one of the electrons is misidentified as a  $\tau$  lepton. Di-boson(WW,WZ,ZZ) events, single Top, SM Higgs boson production( $H \rightarrow \tau\tau$ ) and  $W\gamma^{(*)}$ . These backgrounds are all estimated from MC simulation.

## CHAPTER 7

### Signal extraction and systematics

#### 7.1 Boosted decision trees method

Boosted decision trees(BDT) [47, 61] is used in the  $H \rightarrow \mu\tau_h$  analysis. With this MVA method, the final result, which is referred as BDT fit analysis, improved a factor of two with respect to the  $M_{col}$  fit analysis. There are two parts within BDT, the boosting algorithm and the decision tree. AdaBoost(adaptive boost) is the main boosting algorithm focused on.

Decision tree is a simple two dimensional tree structure. BDT takes in one signal and one background data sets, which are further divided into training and test samples. A set of selected variables are used the training. At the initial node of the tree structure, events are ordered by the value of variables. Trying with every variable, the node is split into two branches by a binary selection of the variable that gives the best separation between signal and background in each of the branch. The splitting continues until the training reaches the number of layers or the number of events in the branch is smaller than the set limit. The splitting stops and the last notes are called leaves. Depending on the main population inside each leaf, the leafs are categorized as signal leaves or background leaves. An example of the tree structure is shown in Figure. 7.1. A criterion is needed for the selection of a variable and the exact point of the cut at the node or branch.

Purity is defined as in Equation. 7.1, in which the superscript s stands of signal events, b stands of background events and W stands of event weight. With the definition of purity, a couple of criteria can be configured. Here the one called Gini, defined in Equation. 7.2, is selected also it is the default one for BDT in TMVA package. In every splitting, the variable value that maximize the criterion in Equation. 7.3 is selected.

$$P = \frac{\sum_s W_s}{\sum_s W_s + \sum_b W_b} \quad (7.1)$$

$$Gini = \left( \sum_{i=1}^n W_i \right) P(1 - P) \quad (7.2)$$

$$Criterion = Gini_{n-1} - Gini_{n \ left} - Gini_{n \ right} \quad (7.3)$$

Decision tree method is powerful, but the performance suffers from fluctuations. The selection of two variables with similar selection quality may be affected by a small change in the training sample. One of the solution is introducing the boosting algorithm. The boosting algorithm not only help stabilize decision tree training with respect to the small fluctuations inside the training data, but also enhance the classification. AdaBoost is one of the boosting algorithms. With AdaBoost, after the first decision tree training, the misclassified events get higher even weights and as input into the second tree. Typically the AdaBoost training 1000 to 2000 trees. Misclassified event weights depend on the training error of




Figure 7.1. Tree structure example in BDT

each decision tree. Training error is calculated as in Equation. 7.4, in which the superscript m, w are the tree label and event weight respectively.  $y_i$  indicates the ith event type, 1 for signal and -1 for background.  $T_m(x_i)$  indicate the what kind of leaf event i with variable x lands on, 1 for signal leaf and -1 for background leaf. Variable  $I(y_i \neq T_m(x_i))$  inside the Equation. 7.4 constructs on the variables  $y_i$  and  $T_m$  equal 1 if  $y_i \neq T_m(x_i)$  or 0 if  $y_i = T_m(x_i)$ . As shown in Equation. 7.5, with the intermediate quantity  $\alpha_m$  for the tree m, the weight for event i is updated to its new value. While  $\text{error}_m$  is required to be less than 0.5 so as the event weights are updated to the right direction. Learning rate parameter  $\beta$  can be used to adjust the step of each re-weighting. The default value of  $\beta$  is 1. Event weights in each tree is renormalized to keep the summed weights constant. The final score of each event after the boosting and training processes is obtained with Equation. 7.8. High score indicates a signal like event while low score indicates a background ground like event.

$$\text{error}_m = \frac{\sum_{i=1}^N w_i I(y_i \neq T_m(x_i))}{\sum_{i=1}^N w_i} \quad (7.4)$$

$$\alpha_m = \beta \times \ln((1 - \text{error}_m)/\text{error}_m) \quad (7.5)$$

$$w_i \rightarrow w_i \times e^{\alpha I(y_i \neq T_m(x_i))} \quad (7.6)$$

$$w_i \rightarrow w_i / \sum_{i=1}^N w_i \quad (7.7)$$

$$T(x) = \sum_{m=1}^{N_{tree}} \alpha_m T_m(x) \quad (7.8)$$

## 7.2 Statistical methods

The physics results are extracted with statistical methods. The limit of branching ratio with confident level(CL) method, significance and best fit branching fraction are mentioned in the analysis.

Probability density function(PDF) and likelihood function are the base functions used. PDF of a continuous variable  $x$  can be interpreted as the likelihood of  $x$  at its different values. Probability density function holds the property that it is normalized to unity.

$$\int f(x) dx = 1$$

Usually PDF of a variable interested accompanied with a set of parameters,

so as PDF is usually written in the form  $f(data|\alpha)$ , which means the probability density function of variables in data given the parameter set  $\alpha$ . These parameters can come from various sources, for example, from theory estimation, detector response and Monte Carlo simulation. All of the parameters in a model or a system, besides the parameters of interest, the other parameters that have an impact on the results are referred as nuisance parameters.

Likelihood function of the same model  $L(data|\alpha)$  is a function of parameter set  $\alpha$  given data. In particle physics, Likelihood function more often in the form of  $L(data|\mu, \theta)$ , in which  $\mu$  stands for signal strength modifier and  $\theta$  stands for a full suite of nuisance parameters. Data in Likelihood function can be experimental data or pseudo-data which is produced to generate sample distribution. Taking the counting experiment as an example, PDFs and likelihood function in a binned sample can be estimated with Poisson distribution as the PDF can be expressed as

$$f(data|u, \theta) = \prod_i \frac{(\mu s_i + b_i)^{n_i}}{n_i!} e^{-\mu s_i - b_i}$$

The  $s_i$  and  $b_i$  stand for the expected number of signal and background events respectively and both of them are a function of nuisance parameter  $\theta$  as  $s_i(\theta)$  and  $b_i(\theta)$ .  $n_i$  represents the number of events observed in bin  $i$ . The likelihood function in numerical form is similar to PDFs but implement the systematic error pdfs  $p(\theta|\tilde{\theta})$  if there are systematics considered in the model [2]. The form of likelihood function can be expressed as

$$\mathcal{L} = \prod_i \frac{(\mu s_i + b_i)^{n_i}}{n_i!} e^{-\mu s_i - b_i} \cdot p(\tilde{\theta}|\theta)$$

Both PDF and likelihood function rely on the knowledge of nuisance parameters. Auxiliary measurements or control regions are often used for the scale estimation of systematic uncertainties. With observations in different auxiliary measurements, a PDF  $p(\tilde{\theta}|\theta)$  which is referred as the posteriors can be measured. Together with  $\pi(\theta)$  which is referred as a prior, the systematic error pdfs can be constructed under the Bayes' theorem as

$$\rho(\theta|\tilde{\theta}) \sim p(\tilde{\theta}|\theta) \cdot \pi_\theta(\theta)$$

The systematic error pdfs can be improved and less affected by the choice of prior  $\pi(\theta)$  [54] by the auxiliary measurements.

In the search of lepton flavour violaton higgs decay, the discovery will be finding the higgs events either decay into  $\mu\tau$  pair in 13 TeV search or  $e\tau$  pair in 8 TeV search. LFV is forbidden in Standard Model(SM), thus, if take SM as a background only model, roughly speaking, a discovery can be claimed if the observation is not compatible with the background only model [34]. To test the compatibility of the observed data with respect to the background only model and further calculate limits on the signal strength modifier of the signal+background hypothesis, the quantity test statistics and the PDFs of the two model under tested are needed.

Test statistics is a function which can map a set of data into a number. According to the Neyman-Pearson lemma, likelihood ratio of two model under test gives the most powerful test, which is expressed as

$$\tilde{q}_\mu = -2\ln \frac{\mathcal{L}(data|\mu, \tilde{\theta}_\mu)}{\mathcal{L}(data|\hat{\mu}, \hat{\theta})}$$




Figure 7.2. Test statistics distribution of signal+background and background only PDFs and the observed valued shown in arrow.

$\mu$  is the given signal strength modifier and  $0 < \hat{\mu} < \mu$ . In the formula of  $\tilde{q}_\mu$ ,  $\hat{\theta}$  is referred as conditional maximum likelihood estimator, given the signal strength modifier  $\mu$ , while in the denominator,  $\hat{\mu}$  and  $\hat{\theta}$  are allowed to flow freely to maximize the likelihood. The PDFs of signal+background model and background only model can be calculated in the following way. The observed  $\theta_\mu^{obs}$  and  $\theta_0^{obs}$  for signal+background and background only model are obtained by maximizing the likelihood function respectively, also the observed  $\tilde{q}_\mu^{obs}$  are calculated give  $\mu$ . The PDFs of signal+background model  $f(\tilde{q}_\mu|\mu, \theta_\mu^{obs})$  and background only model  $f(\tilde{q}_\mu|0, \theta_0^{obs})$  are constructed with the MC pseudo-data as shown in Figure. 7.2.

Confident Level(CLs) are defined with two values  $p_\mu$  and  $1-p_b$ . In signal+background model,  $p_\mu$  is defined as

$$p_\mu = P(\tilde{q}_\mu \geq \tilde{q}_\mu^{obs} | \text{signal+background}) = \int_{\tilde{q}_\mu^{obs}}^{\infty} f(\tilde{q}_\mu|\mu, \hat{\theta}_\mu^{obs}) d\tilde{q}_\mu$$




Figure 7.3. Signal strength modifier  $\mu$  at 95%  $CL_s$  distribution from the MC pseudo-data. The right plot is the cumulative distribution of  $\mu^{95\%}$  with  $\pm 1\sigma$  and  $\pm 2\sigma$  bands.

In background only model  $1-p_b$  is defined as

$$1 - p_b = P(\tilde{q} \geq \tilde{q}_\mu^{obs} | \text{background only}) = \int_{\tilde{q}_0^{obs}}^{\infty} f(\tilde{q}_\mu | 0, \hat{\theta}_0^{obs}) d\tilde{q}_\mu$$

The  $CL_s$  then is expressed as

$$CL_s = \frac{p_\mu}{1 - p_b}$$

With a given value of signal strength modifier  $\mu$ ,  $CL_s \leq \alpha$ , then the signal+background is said excluded with  $(1 - \alpha)$   $CL_s$ . Usually  $\alpha$  is picked as 95%.

The expected limit shown in lepton flavour violation Higgs decay is the median 95%  $CL_s$  upper limit with the  $\pm 1\sigma$  and  $\pm 2\sigma$  bands in the background only model. By generating a large number of background only pseudo-data, the CLs and  $\mu^{95\%}$  of each toy are calculated. An example of  $\mu_{95\%}$  distributions is shown in Figure. 7.3. As shown in the cumulative distribution of  $\mu^{95\%}$ , the median corresponds to 50%, while  $\pm 1\sigma$ ,  $\pm 2\sigma$  correspond to range 16% to 84% and 2.5% to 97.5% respectively.

The estimator p value and significance are used to check if the data is compatible with background only model. Test statistics in the checking of background only model is expressed as

$$\tilde{q}_\mu = -2\ln \frac{\mathcal{L}(data|0, \tilde{\theta}_\mu)}{\mathcal{L}(data|\hat{\mu}, \hat{\theta})}$$

Similar procedure as mentioned above to get the PDF of background ground only model  $f(q_0|0, \hat{\theta}_0^{obs})$  and  $q_0^{obs}$ . p-value corresponding to the experimental observable is calculated as

$$p_0 = P(q_0 \geq q_0^{obs}) = \int_{q_0^{obs}}^{\infty} f(q_0|0, \hat{\theta}_0^{obs}) dq_0$$

Significance Z of the observable with respect to the background only model test can be derived from p value [2].

$$p = \int_Z^{\infty} \frac{1}{\sqrt{2\pi}} \exp(-x^2/2) dx$$

The term used in the discovery as  $5\sigma$  correspond to  $Z=5$  and  $p=2.8 \times 10^{-7}$ . The distribution of test statistics tends to a chi-squared distribution as according to the Wilk's theorem, which gives a quick way of estimating of pdfs in test statistics [34]. In the case when the expected number of events is large, expected limit with this asymptotic approximation gives a fairly well performance and save the computing time for toy samples. Asymptotic approximation can also used in the estimation

of p value as

$$p^{estimate} = \frac{1}{2} \left[ 1 - erf \left( \sqrt{q_0^{obs}/2} \right) \right]$$

### 7.3 Systematics

Systematic uncertainties originated from different sources that experimentally or theoretically affects the physics results. These uncertainties are considered by the effects on the normalization and shape of the distribution of different processes.

#### 7.3.1 Systematics used in $H \rightarrow \mu\tau_h$

The systematic uncertainties considered in this analysis are summarized in Table. 7.1 and Table. 7.2. The uncertainty on muon trigger, identification and isolation together amounts to 2% and the hadronic tau lepton efficiency amounts to 5%. The uncertainties on lepton selections that include trigger, ID, isolation efficiencies are estimated Tag and probe method with Z boson data sets [22, 28, 48, 49, 51]. The systematic uncertainty of b tagging veto is taken from the uncertainty of b tagging veto efficiency measurement, which adjusts b tagging veto performance in simulation to match data samples. The exact values used in each category of  $t\bar{t}$  and single top samples are in Table. 7.3. The uncertainties on  $Z \rightarrow \tau\tau$ , WW, WZ, ZZ,  $t\bar{t}$ , single top backgrounds mainly come from the uncertainties on the cross section measurement of each process. The normalization uncertainty of  $\mu$  fake  $\tau$  scale factor measurement and the uncertainty on  $Z \rightarrow \mu\mu$  process together amounts to 25%. An additional shape uncertainty of muon misidentifying tau is extracted from the measurement of the scale factor and treats independently of hadronic tau decay modes.  $Z \rightarrow \mu\mu$  is the main source that contributes to

$\mu$  fake  $\tau$ . The uncertainty of the misidentified lepton background is estimated from the matching of the same sign control region which is defined as region II in Table. 6.1. The 30% on normalization uncorrelated between each category and additional 10% partially correlated within each category is a conservative estimation. Only the tau shape uncertainties are considered, which are taken from the variations of the misidentified tau parameters in fitting. Muon misidentified shape uncertainty is omitted as it would be negligible compared with the bin-by-bin uncertainty applied. The shape uncertainties of jet energy scale is estimated by varying the fitting parameters of different sources that affect this scale by one  $\sigma$ . Tau energy scale is treated as shape uncertainties and each of the tau decay modes is considered as independently. Unclustered energy refers to the energy loses from the jets  $p_T < 10$  GeV and the PF candidates that are not included in the jets reconstruction. The unclustered energy uncertainty is considered independently for charged particles, photons, neutral hadrons and very forward particles. The uncertainties on Higgs boson cross section is affected by the renormalization scales, factorization, parton distribution functions(PDF) and the strong coupling constant( $\alpha_s$ ). These affections in normalization are taken from [36].The uncertainties on Higgs cross section also affect the acceptance and result in the migration of events between categories. The bin-by-bin uncertainty is applied to account for the statistics uncertainties within each bin. These uncertainties are uncorrected between different bins and categories. The uncertainty of integrated luminosity [5] amounts to 2.5%.

TABLE 7.1

Part one of the systematic uncertainties considered in  $H \rightarrow \mu\tau_h$  analysis.

All of the uncertainties listed in the systematic tables are correlated between categories besides the ones after the sign  $\oplus$ . These are the uncertainties correlated in within each category but independent between categories. The theoretical uncertainties related to the acceptance and migration of events are listed in a range. The negative or positive values indicate a anticorrelated or correlated between categories

Systematic uncertainty	$H \rightarrow \mu\tau_h$
Muon trigger/identification/isolation	2%
Hadronic tau lepton efficiency	5%
b tagging veto	2.0–4.5%
$Z \rightarrow \tau\tau + \text{jets}$ background	10% $\oplus$ 5%
WW, ZZ background	5% $\oplus$ 5%
t <bar>t background</bar>	10% $\oplus$ 5%
Single top quark background	5% $\oplus$ 5%
$\mu \rightarrow \tau_h$ background	25%
Jet $\rightarrow \tau_h, \mu$ background	30% $\oplus$ 10%
Jet energy scale	3–20%
$\tau_h$ energy scale	1.2%

TABLE 7.2

Part two of the systematic uncertainties considered in  $H \rightarrow \mu\tau_h$  analysis

Systematic uncertainty	$H \rightarrow \mu\tau_h$
$\mu \rightarrow \tau_h$ energy scale	1.5%
$\mu$ energy scale	0.2%
Unclustered energy scale	$\pm 1\sigma$
Renorm./fact. scales (g gH) [36]	3.9%
Renorm./fact. scales (VBF and VH) [36]	0.4%
PDF + $\alpha_s$ (g gH) [36]	3.2%
PDF + $\alpha_s$ (VBF and VH) [36]	2.1%
Renorm./fact. acceptance (g gH)	-3.0% – +2.0%
Renorm./fact. acceptance (VBF and VH)	-0.3% – +1.0%
PDF + $\alpha_s$ acceptance (g gH)	-1.5% – +0.5%
PDF + $\alpha_s$ acceptance (VBF and VH)	-1.5% – +1.0%
Integrated luminosity	2.5%

TABLE 7.3

b tagging veto systematic uncertainty in each category

	0 jet	1 jet	2 jets gg-enriched	2 jets VBF-enriched
$t\bar{t}$	–	2.45%	4.37%	2.59%
$t$	–	2.11%	3.07%	1.98%

### 7.3.2 Systematic uncertainties in $H \rightarrow e\tau_h$

The systematic uncertainties affect the  $H \rightarrow e\tau_h$  analysis are summarized in Table. 7.4, 7.6 and 7.5, which includes the ones affect the normalization and the ones affects the shape of  $M_{col}$  distribution. The uncertainty of the electron measurements, which include the trigger, ID and isolation come from the Tag and Probe measurement with Z boson datasets [21, 51]. The uncertainties from the  $Z \rightarrow \tau\tau$  is from the uncertainty of the cross section measurement [25] and the  $\tau$  identification in the embedded technique. The normalization uncertainty on  $Z \rightarrow \mu\mu$  amounts to 30% which is the from the measurement of cross section and statistics uncertainty in the yields. An extra 5% shape uncertainty due to the mismeasured energy of the electron reconstructed as  $\tau$  in  $Z \rightarrow ee$  background. The shift in  $M_{col}$  distribution is measured by comparison between data and MC simulation. The uncertainty in misidentified tau lepton background is 30% normalization uncertainty correlated between categories and shape uncertainties. The shape uncertainties is obtained by varying the parameters from fitting the fake ratio one standard deviation. The uncertainty from the pileup process are estimated by varying the total inelastic cross section by  $\pm 5$  percentage [24]. The uncertainty from Diboson, single top quark background is taken from the measurement of the cross section. In  $t\bar{t}$  background, the uncertainty in 0 jet, 1 jet category are taken from the cross section measurement, in 2 jets category, an extra 33% uncertainty uncorrelated between categories is added due to the statistical uncertainty. The luminosity uncertainty amounts to 2.6%. Jet energy scale(JES) and resolution is measured with  $\gamma/Z+jets$  and dijet data [20]. The uncertainty from JES is applied as a function of  $p_T$  and  $\eta$ . Jet energy resolution(JER) uncertainty is obtained by smearing jets energy as a function of  $p_T$  and  $\eta$ . Both JES and JER affect the

shape of  $M_{col}$  distribution and are taken as shape uncertainties. The energy of the jets below 10 GeV and PF objets that are not clustered are accounted as the unclustered energy scale, which is also taken as shape uncertainties. The uncertainty of  $\tau_h$  energy scale is from the comparing  $Z \rightarrow \tau\tau$  distruption between data and MC sample and applied as a shape uncertainty. The theoretical uncertainties considered in the  $H \rightarrow e\tau_h$  analysis are listed in Table. 7.6. There are several sources that contribute to the theoretical uncertainties and considered fully correlated between LFV Higgs and SM Higgs production. The uncertainty in Parton distribution function is obtained by counting the yields with different PDFs, CT10 [56], MSTW [55], NNPDF [13] from the recommendation in PDF4LHC [14]. The uncertainty in renormalization and factorization scales are from scaling up and down by a factor of two to their normal values( $\mu_R = \mu_F = M_H/2$ ). The uncertainty in underlying events and parton shower is estimated by checking with different PYTHIA tunes. The theoretical uncertainties are all correlated or anticorrelated which have a minus superscript and are the results of the events migration.

TABLE 7.4: The normalization systematic uncertainties considered in  $H \rightarrow e\tau_h$  analysis. The uncertainties are correlated between categories besides the ones after  $\oplus$ . These uncertainties are uncorrelated between categories.

Systematic uncertainty	$H \rightarrow e\tau_h$		
	0-jet	1-jet	2-jet
Electron trigger/ID/isolation	1%	1%	2%
Efficiency of $\tau_h$	6.7%	6.7%	6.7%
$Z \rightarrow \tau\tau$ background	3% $\oplus$ 5%	3% $\oplus$ 5%	3% $\oplus$ 10%
$Z \rightarrow ee$ background	30%	30%	30%
Misidentified leptons background	30%	30%	30%
Pileup	4%	4%	2%
WW, WZ, ZZ+jets background	15%	15%	15%
t $\bar{t}$ background	10%	10%	10% $\oplus$ 33%
Single top quark background	25%	25%	25%
Luminosity	2.6%	2.6%	2.6%

TABLE 7.5

The systematic uncertainties that affect the shape of  $M_{col}$  distribution.

Systematic Uncertainty	$H \rightarrow e\tau_h$
$Z \rightarrow ee$ bias	5%
Jet energy scale	3%–7%
Jet energy resolution	1%–10%
Unclustered energy scale	10%
$\tau_h$ energy scale	3%

TABLE 7.6: Theoretical uncertainties that affects the Higgs boson production cross section. These uncertainties are correlated or anticorrelated (with minus sign superscript) between all of the categories.

Systematic uncertainty	Gluon fusion			Vector boson fusion		
	0-jet	1-jet	2-jet	0-jet	1-jet	2-jet
Parton distribution function	9.7%	9.7%	9.7%	3.6%	3.6%	3.6%
Renormalization/factorization scale	8%	10%	30 <sup>-</sup> %	4%	1.5%	2%
Underlying event/parton shower	4%	5 <sup>-</sup> %	10 <sup>-</sup> %	10%	<1%	1 <sup>-</sup> %

## CHAPTER 8

### LFV Higgs decay searching results

Results can be extracted after applying the selection criteria and with the full consideration of the relevant systematics. In both  $H \rightarrow \mu\tau_h$  and  $H \rightarrow e\tau_h$  analysis, a maximum likelihood fit is performed to derive the expected and observed limits. Each category in the analysis is fitted separately and then combined. Systematic uncertainties are used as nuisance parameter in the fittings. The upper limits on signal branching fraction with  $CL_s$  criterion are set with asymptotic formula.

#### 8.0.1 Results in $H \rightarrow \mu\tau_h$ search

In Lepton flavour violation  $H \rightarrow \mu\tau_h$  search at central of mass energy of 13 TeV, the analysis is performed in two parallel searching method, the  $M_{col}$  fit analysis and BDT fit analysis. After the selection in  $M_{col}$  fit analysis and adjusting all of distribution by the fit, the distribution of the signal and background discriminant variable  $M_{col}$  is shown in Figure. 8.3. In the BDT fit analysis, in which the BDT discriminator is used as the variable to distinguish signal from background. After fitting to all of the distributions, the observed data versers the expected backgrounds is shown in Figure. 8.2. No excess over the backgrounds is observed in both of the methods. The expected and observed median upper limits at 95% CL and the best fit branching fraction in the  $H \rightarrow \mu\tau_h$  search with the  $M_{col}$  fit

analysis is shown in Table. 8.1 and the result of the BDT fit analysis is shown in Table. 8.2.

The search in  $H \rightarrow \mu\tau$  with CMS 13 TeV,  $36 \text{ fb}^{-1}$  data is motivated by the search CMS 8 TeV  $19.7 \text{ fb}^{-1}$   $H \rightarrow \mu\tau$ , in which a significance of 2.4 standard deviation excess of data with respect to the SM background-only hypothesis was observed. This thesis is written in  $H \rightarrow \mu\tau_h$  search, together with another search  $H \rightarrow \mu\tau_e$  [32], where the tau lepton decays electronically to an electron, instead of tau hadronically decay, gives the most recent and update check for the best fit branching ratio  $H \rightarrow \mu\tau$ . This combined result is shown with BDT fit analysis and  $M_{col}$  fit analysis separately in Figure. 8.3. No evidence of excess is found in this combined result. Compared with the 8 TeV result, the new upper limits on the signal branching fraction improved a factor of 5. The BDT fit analysis is more sensitive to the  $M_{col}$  fit analysis. The new and tighter limit result comes from the BDT fit analysis.

The constraints on Yukawa couplings,  $|Y_{\mu\tau}|, |Y_{\tau\mu}|$ , can be derived from the limits on branching ratio  $\mathcal{B}(H \rightarrow \mu\tau)$ , which is the combined result between  $H \rightarrow \mu\tau_h$  search and  $H \rightarrow \mu\tau_e$  search. Lepton flavour violating decay can rise from the tree level by assuming the Yukawa interactions. In  $H \rightarrow \mu\tau$  decays, the Yukawa interactions is related to the decay width  $\Gamma(H \rightarrow \ell^\alpha \ell^\beta)$ ,  $\ell^\alpha, \ell^\beta$  can be  $\mu$  or  $\tau$  and  $\ell^\alpha \neq \ell^\beta$ .  $\Gamma$  in terms of the Yukawa couplings is given by:

$$\Gamma(H \rightarrow \ell^\alpha \ell^\beta) = \frac{m_H}{8\pi} (|Y_{\ell^\beta \ell^\alpha}|^2 + |Y_{\ell^\alpha \ell^\beta}|^2), \quad (8.1)$$




Figure 8.1.  $M_{col}$  distribution after the selection. The signal and backgrounds in the plots have been normalized to the best fit values. The gray bands shows the total uncertainties in each bin and the signal is plotted with the branching ratio of 5% of the higgs decay branching ratio for visualization purpose.




Figure 8.2. BDT discriminator distribution after the selection. The signal and backgrounds in the plots has been normalized to the best fit values. The gray bands shows the total uncertainties in each bin and the signal is plotted with the branching ratio of 5% of the higgs decay branching ratio for visualization purpose.

TABLE 8.1

$M_{col}$  fit analysis expected and observed upper limits at 95% CL and the best fit branching fractions in each of the categories.

Expected limits (%)					
	0-jet	1-jet	2-jets	VBF	Combined
$\mu\tau_h$	< 1.14	< 1.26	< 2.12	< 1.41	< 0.71
Observed limits (%)					
	0-jet	1-jet	2-jets	VBF	Combined
$\mu\tau_h$	< 1.04	< 1.74	< 1.65	< 1.30	< 0.66
Best fit branching fractions (%)					
	0-jet	1-jet	2-jets	VBF	Combined
$\mu\tau_h$	-0.30 $\pm$ 0.45	0.68 $\pm$ 0.56	-1.23 $\pm$ 1.04	-0.23 $\pm$ 0.66	-0.08 $\pm$ 0.34

TABLE 8.2

BDT fit analysis expected and observed upper limits at 95% CL and the best fit branching fractions in each of the categories.

Expected limits (%)					
	0-jet	1-jet	2-jets	VBF	Combined
$\mu\tau_h$	<0.43	<0.56	<0.94	<0.58	<0.29
Observed limits (%)					
	0-jet	1-jet	2-jets	VBF	Combined
$\mu\tau_h$	<0.51	<0.53	<0.56	<0.51	<0.27
Best fit branching fractions (%)					
	0-jet	1-jet	2-jets	VBF	Combined
$\mu\tau_h$	$0.12 \pm 0.20$	$-0.05 \pm 0.25$	$-0.72 \pm 0.43$	$-0.22 \pm 0.31$	$-0.04 \pm 0.14$




Figure 8.3. The expected and observed 95% CL limits on branching ratio of total  $H \rightarrow \mu\tau$  which shows the results from both  $H \rightarrow \mu\tau_h$  and  $H \rightarrow \mu\tau_e$  in the  $M_{col}$  fit analysis and BDT fit analysis.

and the branching fraction by:

$$\mathcal{B}(H \rightarrow \ell^\alpha \ell^\beta) = \frac{\Gamma(H \rightarrow \ell^\alpha \ell^\beta)}{\Gamma(H \rightarrow \ell^\alpha \ell^\beta) + \Gamma_{\text{SM}}}. \quad (8.2)$$

The 125 GeV Standard Model Higgs decay width is taken as  $\Gamma_{\text{SM}} = 4.1 \text{ MeV}$  [37]. In BDT fit analysis, the 95% CL upper limit on the Yukawa coupling is  $\sqrt{|Y_{\mu\tau}|^2 + |Y_{\tau\mu}|^2} < 1.43 \times 10^{-3}$  and in  $M_{\text{col}}$  fit analysis,  $\sqrt{|Y_{\mu\tau}|^2 + |Y_{\tau\mu}|^2} < 2.05 \times 10^{-3}$ . The upper limit on Yukawa couplings of BDT fit analysis is shown in Figure. 8.4. The comparison with the previous experimental search and theoretical naturalness results is also shown in the plot.




Figure 8.4. Upper limit of flavour violating Yukawa coupling  $|Y_{\mu\tau}|, |Y_{\tau\mu}|$  from the BDT fit analysis. The red solid line represents the expected limit and the black solid line is the observed limit. 68% and 95% range of containing the observed limit is shown in the green and yellow band. The result from  $\tau \rightarrow 3\mu$  is shown in dark green [45, 46, 59] and the one from  $\tau \rightarrow \mu\gamma$  is shown in lighter green [45, 59]. The theoretical naturalness limit  $Y_{ij}Y_{ji} \leq m_i m_j / v^2$  is shown by purple diagonal line [45].

### 8.0.2 Results in $H \rightarrow e\tau_h$ search

The search of  $H \rightarrow e\tau_h$  is performed in  $M_{col}$  fit analysis with CMS 8 TeV 19.7  $fb^{-1}$ . The analysis is binned with 3 categories. After applying the selection cuts and adjusting the signal and background processes by the fit, the fitting variable  $M_{col}$  distribution is shown in Figure. 8.5. The number of event yields of both signal and background processes are shown in Table. 8.3. The yields are normalized to the integrated luminosity of 19.7  $fb^{-1}$  and the signal branching ratio is assumed to be 0.69% of the SM higgs production cross section. This thesis is focused and written on the analysis  $H \rightarrow e\tau_h$  which is one part of the whole search on lepton flavour violating higgs decay to  $e\tau$ . Another closely related search is  $H \rightarrow e\tau_\mu$  [4]. The results of each categories and combined ones of the two analysis  $H \rightarrow e\tau_h$  and  $H \rightarrow e\tau_\mu$  on the expected and observed upper signal branching fraction  $\mathcal{B}(H \rightarrow e\tau)$  limits is shown in Table. 8.4 and summarized in Figure. 8.6. The combined upper limit on  $\mathcal{B}(H \rightarrow e\tau) < 0.69$  is the first direct search on this channel and the tightest limit by the time the result was published. Yukawa coupling between electron and tau leptons is related to decay width  $\Gamma$  and branching fraction  $\mathcal{B}$  as shown in Equation. 8.1 and 8.2. The 95% CL upper limit on Yukawa coupling from the combined result in  $H \rightarrow e\tau$  is  $\sqrt{|Y_{e\tau}|^2 + |Y_{\tau e}|^2} < 2.4 \times 10^{-3}$ . The limit on  $e\tau$  Yukawa coupling from this search and other previous search results are shown in Figure. 8.0.2.




Figure 8.5.  $M_{col}$  distribution after the selection. The signal and backgrounds in the plots have been normalized to the best fit values. The gray bands shows the total uncertainties in each bin and the signal is plotted with the branching ratio of 0.69% of the higgs decay branching ratio for visualization purpose.

TABLE 8.3: Events yields of the signal and backgrounds are shown in the mass range  $100 \text{ GeV} < M_{\text{col}} < 150 \text{ GeV}$ . The branching ratio of  $H \rightarrow e\tau_h$  is assumed to be 0.69% of the SM higgs production cross section. The numbers in the table are normalized to the integrated luminosity of  $19.7 \text{ fb}^{-1}$ .

Jet category:	0-jet	1-jet	2-jet
Misidentified leptons	$3366 \pm 25$	$223 \pm 11$	$8.7 \pm 2.2$
$Z \rightarrow ee, \mu\mu$	$714 \pm 30$	$85 \pm 4$	$3.2 \pm 0.2$
$Z \rightarrow \tau\tau$	$270 \pm 10$	$32 \pm 3$	$1.6 \pm 0.3$
$t\bar{t}, t, \bar{t}$	$10 \pm 2$	$13 \pm 2$	$0.5 \pm 0.2$
$ZZ, WZ, WW$	$53 \pm 2$	$6 \pm 1$	$0.3 \pm 0.1$
SM H background	$12 \pm 1$	$3 \pm 1$	$1.0 \pm 0.1$
Sum of background	$4425 \pm 28$	$363 \pm 11$	$15.3 \pm 2.3$
Observed	4438	375	13
LFV H signal	$61 \pm 4$	$15 \pm 1$	$2.8 \pm 0.5$




Figure 8.6. 95% CL upper limit on branching ratio on  $H \rightarrow e\tau$  with both the results from  $H \rightarrow e\tau_h$  and  $H \rightarrow e\tau_\mu$  and the combined.

TABLE 8.4

$M_{col}$  fit analysis expected and observed upper limits at 95% CL and the best fit branching fractions in each of the categories of both  $H \rightarrow e\tau_h$  and  $H \rightarrow e\tau_\mu$  analyses. The asymmetric one standard-deviation uncertainties around the expected limits are shown in parentheses.

	0-jet	1-jet	2-jet
Expected limits at 95% CL (%)			
$e\tau_\mu$	$<1.63^{(+0.66)}_{(-0.44)}$	$<1.54^{(+0.71)}_{(-0.47)}$	$<1.59^{(+0.93)}_{(-0.55)}$
$e\tau_h$	$<2.71^{(+1.05)}_{(-0.75)}$	$<2.76^{(+1.07)}_{(-0.77)}$	$<3.55^{(+1.38)}_{(-0.99)}$
Observed limits at 95% CL (%)			
$e\tau_\mu$	$<1.83$	$<0.94$	$<1.49$
$e\tau_h$	$<3.92$	$<3.00$	$<2.88$
$e\tau$	$<0.69$		




Figure 8.7. Upper limit of flavour violating Yukawa coupling  $|Y_{e\tau}|, |Y_{\tau e}|$  from the combined  $H \rightarrow e\tau$  result. The red solid line represents the expected limit and the black solid line is the observed limit. 68% and 95% range of containing the observed limit is shown in the green and yellow band. The result from  $\tau \rightarrow 3e$  is shown in gray, the one from  $\tau \rightarrow e\gamma$  is shown in dark green and presented analysis is in light blue. The theoretical naturalness limit  $Y_{ij}Y_{ji} \leq m_i m_j / v^2$  is shown by purple diagonal line [45]

## CHAPTER 9

### Conclusion

## BIBLIOGRAPHY

1. Particle-flow commissioning with muons and electrons from J/Psi and W events at 7 TeV. Technical Report CMS-PAS-PFT-10-003, CERN, Geneva, 2010. URL <http://cds.cern.ch/record/1279347>.
2. Procedure for the LHC Higgs boson search combination in Summer 2011. Technical Report CMS-NOTE-2011-005. ATL-PHYS-PUB-2011-11, CERN, Geneva, Aug 2011. URL <https://cds.cern.ch/record/1379837>.
3. Pileup Jet Identification. Technical Report CMS-PAS-JME-13-005, CERN, Geneva, 2013. URL <http://cds.cern.ch/record/1581583>.
4. Search for lepton flavour violating decays of the higgs boson to  $e\tau$  and  $e\mu$  in proton-proton collisions at  $\sqrt{s} = 8$  TeV. *Phys. Lett. B*, 763:472, 2016. doi: 10.1016/j.physletb.2016.09.062.
5. CMS luminosity measurements for the 2016 data taking period. CMS Physics Analysis Summary CMS-PAS-LUM-17-001, 2017. URL <https://cds.cern.ch/record/2257069>.
6. 06 2018. URL [https://en.wikipedia.org/wiki/Standard\\_Model](https://en.wikipedia.org/wiki/Standard_Model).
7. D. Acosta. Cms trigger improvements towards run ii. *Nuclear and Particle Physics Proceedings*, 273-275:1008 – 1013, 2016. ISSN 2405-6014. doi: <https://doi.org/10.1016/j.nuclphysbps.2015.09.158>. URL <http://www>.

[sciencedirect.com/science/article/pii/S2405601415006471](http://sciencedirect.com/science/article/pii/S2405601415006471). 37th International Conference on High Energy Physics (ICHEP).

8. W. Adam, R. Fröhwirth, A. Strandlie, and T. Todor. Reconstruction of Electrons with the Gaussian-Sum Filter in the CMS Tracker at the LHC. 2005.
9. W. Adam, B. Mangano, T. Speer, and T. Todorov. Track Reconstruction in the CMS tracker. Technical Report CMS-NOTE-2006-041, CERN, Geneva, Dec 2006. URL <https://cds.cern.ch/record/934067>.
10. P. Adzic et al. Energy resolution of the barrel of the CMS electromagnetic calorimeter. *JINST*, 2:P04004, 2007. doi: 10.1088/1748-0221/2/04/P04004.
11. J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H.-S. Shao, T. Stelzer, P. Torrielli, and M. Zaro. The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations. *JHEP*, 07:079, 2014. doi: 10.1007/JHEP07(2014)079.
12. M. Baber, M. Blake, J. Brooke, M. C. Hermida, S. Dasu, T. Durkin, S. Fayer, E. K. Friis, T. Gorski, G. Hall, K. Harder, G. Iles, S. Ives, J. Jones, P. R. Klabbers, A. G. Levine, C. Lucas, R. Lucas, D. Newbold, J. Marrouche, S. Paramesvaran, T. M. Perry, A. Rose, D. Sankey, W. Smith, A. Tapper, A. Thea, and T. Williams. Development and testing of an upgrade to the cms level-1 calorimeter trigger. *Journal of Instrumentation*, 9(01):C01006, 2014. URL <http://stacks.iop.org/1748-0221/9/i=01/a=C01006>.
13. R. D. Ball, L. Del Debbio, S. Forte, A. Guffanti, J. I. Latorre, J. Rojo, and M. Ubiali. A first unbiased global NLO determination of parton dis-

- tributions and their uncertainties. *Nucl. Phys. B*, 838:136, 2010. doi: 10.1016/j.nuclphysb.2010.05.008.
14. M. Botje, J. Butterworth, A. Cooper-Sarkar, A. de Roeck, J. Feltesse, S. Forte, A. Glazov, J. Huston, R. McNulty, T. Sjöstrand, and R. S. Thorne. The PDF4LHC Working Group Interim Recommendations. 2011.
  15. M. Cacciari, G. P. Salam, and G. Soyez. The anti- $k_t$  jet clustering algorithm. *JHEP*, 04:063, 2008. doi: 10.1088/1126-6708/2008/04/063.
  16. M. Cacciari, G. P. Salam, and G. Soyez. FastJet user manual. *Eur. Phys. J. C*, 72:1896, 2012. doi: 10.1140/epjc/s10052-012-1896-2.
  17. R. N. Cahn, S. D. Ellis, R. Kleiss, and W. J. Stirling. Transverse-momentum signatures for heavy Higgs bosons. *Phys. Rev. D*, 35:1626, 1987. doi: 10.1103/PhysRevD.35.1626.
  18. S. Chatrchyan et al. The CMS Experiment at the CERN LHC. *JINST*, 3:S08004, 2008. doi: 10.1088/1748-0221/3/08/S08004.
  19. S. Chatrchyan et al. Aligning the CMS Muon Chambers with the Muon Alignment System during an Extended Cosmic Ray Run. *JINST*, 5:T03019, 2010. doi: 10.1088/1748-0221/5/03/T03019.
  20. S. Chatrchyan et al. Determination of jet energy calibration and transverse momentum resolution in CMS. *JINST*, 6:11002, 2011. doi: 10.1088/1748-0221/6/11/P11002.
  21. S. Chatrchyan et al. Measurement of the inclusive W and Z production cross sections in pp collisions at  $\sqrt{s} = 7$  TeV with the CMS experiment. *JHEP*, 10:132, 2011. doi: 10.1007/JHEP10(2011)132.

22. S. Chatrchyan et al. Performance of CMS muon reconstruction in pp collision events at  $\sqrt{s} = 7$  TeV. *JINST*, 7:P10002, 2012. doi: 10.1088/1748-0221/7/10/P10002.
23. S. Chatrchyan et al. Performance of CMS muon reconstruction in *pp* collision events at  $\sqrt{s} = 7$  TeV. *JINST*, 7:P10002, 2012. doi: 10.1088/1748-0221/7/10/P10002.
24. S. Chatrchyan et al. Measurement of the inelastic proton-proton cross section at  $\sqrt{s} = 7\text{TeV}$ . *Phys. Lett. B*, 722:5, 2013. doi: 10.1016/j.physletb.2013.03.024.
25. S. Chatrchyan et al. Measurement of inclusive  $W$  and  $Z$  boson production cross sections in  $pp$  collisions at  $\sqrt{s} = 8$  TeV. *Phys. Rev. Lett.*, 112:191802, 2014. doi: 10.1103/PhysRevLett.112.191802.
26. S. Chatrchyan et al. Evidence for the 125 GeV Higgs boson decaying to a pair of  $\tau$  leptons. 2014.
27. C. Collaboration". Performance of  $\tau$ -lepton reconstruction and identification in CMS. *JINST*, 7(arXiv:1109.6034. CMS-TAU-11-001. CERN-PH-EP-2011-137):P01001. 33 p, Sep 2011. URL <http://cds.cern.ch/record/1385560>.
28. C. Collaboration. Performance of reconstruction and identification of tau leptons in their decays to hadrons and tau neutrino in LHC Run-2. CMS Physics Analysis Summary CMS-PAS-TAU-16-002, 2016. URL <https://cds.cern.ch/record/2196972>.
29. C. Collaboration. Performance of reconstruction and identification of tau leptons in their decays to hadrons and tau neutrino in LHC Run-2. 2016.

30. C. Collaboration. Reconstruction and identification of tau lepton decays to hadrons and tau neutrino at CMS. *JINST* 11 (2016) P01019, 11, 2016. doi: 10.1088/1748-0221/11/01/P01019.
31. C. Collaboration. Identification of b quark jets at the CMS Experiment in the LHC Run 2. Technical Report CMS-PAS-BTV-15-001, CERN, Geneva, 2016. URL <https://cds.cern.ch/record/2138504>.
32. C. Collaboration. Search for lepton flavour violating decays of the Higgs boson to  $\tau$  and  $e$  in proton-proton collisions at  $\sqrt{s} = 13$  TeV. *JHEP*, 2017.
33. G. Collaboration. Geant4—a simulation toolkit. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 506(3):250 – 303, 2003. ISSN 0168-9002. doi: [https://doi.org/10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8). URL <http://www.sciencedirect.com/science/article/pii/S0168900203013688>.
34. K. Cranmer. Practical Statistics for the LHC. *CERN-2014-003*, pp. 267 - 308, 2015. doi: 10.5170/CERN-2015-004.301.
35. S. Dasu et al. CMS. The TriDAS project. Technical design report, vol. 1: The trigger systems. 2000.
36. D. de Florian et al. Handbook of LHC Higgs cross sections: 4. deciphering the nature of the Higgs sector. CERN Report CERN-2017-002-M, 2016.
37. A. Denner, S. Heinemeyer, I. Puljak, D. Rebuzzi, and M. Spira. Standard model Higgs-boson branching ratios with uncertainties. *Eur. Phys. J. C*, 71: 1753, 2011. doi: 10.1140/epjc/s10052-011-1753-8.

38. A. Djouadi. The anatomy of electroweak symmetry breaking: Tome i: The higgs boson in the standard model. *Physics Reports*, 457(1):1 – 216, 2008. ISSN 0370-1573. doi: <https://doi.org/10.1016/j.physrep.2007.10.004>. URL <http://www.sciencedirect.com/science/article/pii/S0370157307004334>.
39. S. D. Ellis and D. E. Soper. Successive combination jet algorithm for hadron collisions. *Phys. Rev.*, D48:3160–3166, 1993. doi: 10.1103/PhysRevD.48.3160.
40. L. Evans and P. Bryant. Lhc machine. *Journal of Instrumentation*, 3(08):S08001, 2008. URL <http://stacks.iop.org/1748-0221/3/i=08/a=S08001>.
41. R. Fruhwirth. Application of Kalman filtering to track and vertex fitting. *Nucl. Instrum. Meth.*, A262:444–450, 1987. doi: 10.1016/0168-9002(87)90887-4.
42. H. M. Georgi, S. L. Glashow, M. E. Machacek, and D. V. Nanopoulos. Higgs bosons from two gluon annihilation in proton proton collisions. *Phys. Rev. Lett.*, 40:692, 1978. doi: 10.1103/PhysRevLett.40.692.
43. V. Gori. The CMS High Level Trigger. *Int. J. Mod. Phys. Conf. Ser.*, 31:1460297, 2014. doi: 10.1142/S201019451460297X.
44. D. J. Griffiths. *Introduction to elementary particles; 2nd rev. version*. Physics textbook. Wiley, New York, NY, 2008. URL <https://cds.cern.ch/record/111880>.
45. R. Harnik, J. Kopp, and J. Zupan. Flavor violating higgs decays. *JHEP*, 03:26, 2013. doi: 10.1007/JHEP03(2013)026.
46. K. Hayasaka et al. Search for lepton flavor violating  $\tau$  decays into three

- leptons with 719 million produced  $\tau^+\tau^-$  pairs. *Phys. Lett. B*, 687:139, 2010. doi: 10.1016/j.physletb.2010.03.037.
47. A. Hoecker, P. Speckmayer, J. Stelzer, J. Therhaag, E. von Toerne, H. Voss, M. Backes, T. Carli, O. Cohen, A. Christov, D. Dannheim, K. Danielowski, S. Henrot-Versille, M. Jachowski, K. Kraszewski, A. Krasznahorkay, Jr., M. Kruk, Y. Mahalalel, R. Ospanov, X. Prudent, A. Robert, D. Schouten, F. Tegenfeldt, A. Voigt, K. Voss, M. Wolter, and A. Zemla. TMVA - Toolkit for Multivariate Data Analysis. *ArXiv Physics e-prints*, Mar. 2007.
48. V. Khachatryan et al. Measurements of inclusive W and Z cross sections in pp collisions at  $\sqrt{s} = 7$  TeV. *J. High Energy Phys.*, 01:080, 2011. doi: 10.1007/JHEP01(2011)080.
49. V. Khachatryan et al. Performance of electron reconstruction and selection with the CMS detector in proton-proton collisions at  $\sqrt{s} = 8$  TeV. *JINST*, 10:P06005, 2015. doi: 10.1088/1748-0221/10/06/P06005.
50. V. Khachatryan et al. Performance of electron reconstruction and selection with the CMS detector in proton-proton collisions at  $\sqrt{s} = 8$  TeV. *JINST*, 10:P06005, 2015. doi: 10.1088/1748-0221/10/06/P06005.
51. V. Khachatryan et al. Reconstruction and identification of tau lepton decays to hadrons and  $\nu_\tau$  at CMS. *JINST*, 11:P01019, 2016. doi: 10.1088/1748-0221/11/01/P01019.
52. V. Khachatryan et al. Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV. *JINST*, 12:P02014, 2017. doi: 10.1088/1748-0221/12/02/P02014.

53. C. Lefèvre. The CERN accelerator complex. Complexe des accélérateurs du CERN. Dec 2008. URL <https://cds.cern.ch/record/1260465>.
54. L. Lista. Practical Statistics for Particle Physicists. 2016.
55. A. D. Martin, W. J. Stirling, R. S. Thorne, and G. Watt. Parton distributions for the LHC. *Eur. Phys. J. C*, 63:189, 2009. doi: 10.1140/epjc/s10052-009-1072-5.
56. P. M. Nadolsky, H.-L. Lai, Q.-H. Cao, J. Huston, J. Pumplin, D. Stump, W.-K. Tung, and C.-P. Yuan. Implications of CTEQ global analysis for collider observables. *Phys. Rev. D*, 78:013004, 2008. doi: 10.1103/PhysRevD.78.013004.
57. G. N. Davidson et al. Universal interface of tauola technical and physics documentation. doi: 10.1016/j.cpc.2011.12.009.
58. C. Olari. The POWHEG-BOX. *Nucl.Phys.Proc.Suppl.* 205-206:36-41, 2010, 2010. doi: 10.1016/j.nuclphysbps.2010.08.016.
59. Particle Data Group, C. Patrignani, et al. Review of Particle Physics. *Chin. Phys. C*, 40:100001, 2016. doi: 10.1088/1674-1137/40/10/100001.
60. M. E. Peskin and D. V. Schroeder. *An Introduction to quantum field theory*. Addison-Wesley, Reading, USA, 1995. ISBN 9780201503975, 0201503972. URL <http://www.slac.stanford.edu/~mpeskin/QFT.html>.
61. B. P. Roe et al. Boosted Decision Trees as an Alternative to Artificial Neural Networks for Particle Identification. 2005.

62. D. Rabady, J. Ero, G. Flouris, J. Fulcher, N. Loukas, E. Paradas, T. Reis, H. Sakulin, and C.-E. Wulz. Upgrade of the CMS muon trigger system in the barrel region. *Nucl. Instrum. Meth.*, A845:616–620, 2017. doi: 10.1016/j.nima.2016.05.071.
63. A. M. Sirunyan et al. Particle-flow reconstruction and global event description with the cms detector. *JINST*, 12:P10003, 2017. doi: 10.1088/1748-0221/12/10/P10003.
64. A. M. Sirunyan et al. Particle-flow reconstruction and global event description with the cms detector. *JINST*, 12:P10003, 2017. doi: 10.1088/1748-0221/12/10/P10003.
65. A. M. Sirunyan et al. Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV. *JINST*, 13:P05011, 2018. doi: 10.1088/1748-0221/13/05/P05011.
66. T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands. An introduction to PYTHIA 8.2. *Comput. Phys. Commun.*, 191:159, 2015. doi: 10.1016/j.cpc.2015.01.024.
67. E. Yazgan and the CMS ECAL/HCAL Collaborations. The cms barrel calorimeter response to particle beams from 2 to 350 gev/c. *Journal of Physics: Conference Series*, 160(1):012056, 2009. URL <http://stacks.iop.org/1742-6596/160/i=1/a=012056>.
68. S. Yu.L.Dokshitzer, G.D.Leder and B.R.Webber. Better Jet Clustering Algorithms. *JHEP* 9708:001, 1997, 1997. doi: 10.1088/1126-6708/1997/08/001.

69. A. Zabi, F. Beaudette, L. Cadamuro, L. Mastrolorenzo, T. Romanteau, J. Sauvan, T. Strebler, J. Marrouche, N. Wardle, R. Aggleton, F. Ball, J. Brooke, D. Newbold, S. Paramesvaran, D. Smith, M. Baber, A. Bundock, M. Citron, A. Elwood, G. Hall, G. Iles, C. Laner, B. Penning, A. Rose, A. Tapper, T. Durkin, K. Harder, S. Harper, C. Shepherd-Themistocleous, A. Thea, and T. Williams. Triggering on electrons, jets and tau leptons with the cms upgraded calorimeter trigger for the lhc run ii. *Journal of Instrumentation*, 11(02):C02008, 2016. URL <http://stacks.iop.org/1748-0221/11/i=02/a=C02008>.

*This document was prepared & typeset with pdfL<sup>A</sup>T<sub>E</sub>X, and formatted with NDDiss2<sub>E</sub> classfile (v3.0[2005/07/27]) provided by Sameer Vijay.*