

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>
This is a question about risk factors for Alzheimer's , so I will consider each option's association with Alzheimer's risk . Lower education and family history of Alzheimer's are known risk factors . Being married , on the other hand , has been associated with a lower risk , possibly due to the social support it provides . Higher intake of saturated fats and smoking are both linked to cardiovascular health , which is connected to overall cognitive health and Alzheimer's . Older age , of course , is a clear risk factor . High consumption of fish , particularly fatty fish , is rich in omega - 3 fatty acids , which support brain health . Regular drinking has complex effects on health , but moderate consumption is often associated with a lower risk of dementia . Being African or Hispanic American and living in a city may have mixed associations with dementia risk , depending on socioeconomic factors . Factors A , B , E , and I are generally associated with an increased risk , so I can eliminate those . Options C , F , and J are also risk factors , so this leaves me with either G or H as a correct answer . However , G is definitely a major risk factor , so the option would be H .
</think><answer> H . Eating fish

