

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>

1 . Being African or Hispanic American and having a family history of Alzheimer 's are associated with an increased risk of developing Alzheimer 's , so they are not associated with a decreased risk .

2 . Lower levels of education and family history of Alzheimer 's are associated with a higher risk , not a decreased risk .

3 . Living in a city , older age , and having a family history of Alzheimer 's are also not associated with a decreased risk .

4 . Being a smoker and high intake of saturated fats are linked to a higher risk , not a decreased risk .

5 . Being married and eating fish might offer some protection against Alzheimer 's , and drinking alcohol regularly could either increase or decrease the risk , depending on the amount and pattern .

6 . High intake of saturated fats is linked to a higher risk , not a decreased risk .

7 . Eating fish , especially omega - 3 fatty acids , which are present in fish , is often associated with a lower risk of Alzheimer 's .

8 . Older age is associated with a higher risk , not a decreased risk .

9 . Drinking alcohol regularly , especially in excess , can increase the risk of Alzheimer 's , but moderate consumption (one drink for women , two drinks for men) is linked to a lower risk .

10 . High intake of saturated fats can increase the risk , not decrease it .

</think>

< answer >

H . Eating fish

</answer>

