

Question:

Which of the following factors is associated with a decreased risk of Alzheimer's?

Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

<think>

being African or Hispanic American , lower level of education , living in a city , being a smoker , drinking alcohol regularly , and having a family history of Alzheimer's are all risk factors for Alzheimer's .

older age is also considered a risk factor .

being married , having a family history of Alzheimer's , and eating fish are not associated with increased risk .

factors associated with a decreased risk of Alzheimer's include a higher level of education , living in a city , being married , and eating fish .

men are less likely to develop Alzheimer's than women , and since being married is associated with a lower risk , and there are more married men , it can be inferred that being married is associated with a decreased risk .

having a family history of Alzheimer's is a risk factor .

smoking and drinking alcohol regularly increase the risk of Alzheimer's .

drinking fish is a protective factor .

eating fish is associated with a decreased risk of Alzheimer's .

taking fish oil supplements is associated with a decreased risk of Alzheimer's .

fish oil is a source of omega - 3 fatty acids , which is associated with a lower risk of Alzheimer's .

omega - 3 fatty acids are found in fish such as salmon , mackerel , herring , lake trout , albacore tuna , and anchovies .

omega - 3 fatty acids may reduce the risk of Alzheimer's by improving brain cell connections and by reducing the inflammation of the brain .

omega - 3 fatty acids are also associated with a lower risk of cardiovascular disease , which may contribute to a lower risk of Alzheimer's .

taking fish oil supplements and eating fish are associated with a lower risk of Alzheimer's .

taking fish oil supplements is associated with a lower risk of Alzheimer's .

omega - 3 fatty acids are associated with a lower risk of Alzheimer's .

omega - 3 fatty acids are a type of unsaturated fat , which is not associated with an increased risk of Alzheimer's .

only option H "Eating fish" is not associated with an increased risk of Alzheimer's .

</think>

<answer>

H. Eating fish

</answer>