

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>
Decreased risk factors for Alzheimer's disease are those that are associated with a lower likelihood of developing the condition . Being married (D) and eating fish (H) have been linked to a lower risk of Alzheimer's disease . Being African or Hispanic American (A), having a lower level of education (B), living in a city (C), having a family history of Alzheimer's (E), eating a high intake of saturated fats (F), being older (G), smoking (I), and drinking alcohol regularly (J) are all associated with increased risk of Alzheimer's .

The question asks for a decreased risk factor , and among the options , the most strongly supported association with decreased risk is being married (D).

- Answer Options :
- A . Being African or Hispanic American
 - B . A lower level of education
 - C . Living in a city
 - D . Being married
 - E . Having a family history of Alzheimer's
 - F . High intake of saturated fats
 - G . Older age
 - H . Eating fish
 - I . Being a smoker
 - J . Drinking alcohol regularly

</think>
<answer>D . being married</answer>