

Question:

A 22-year-old G1P1 woman comes to the clinic asking about “the morning after pill.” She reports that she had sexual intercourse with her boyfriend last night and she thinks the condom broke. She is not using any other form of contraception. She reports her last menstrual period was 10 days ago, and they are normally regular. The patient’s medical history is significant for obesity, asthma and allergic rhinitis. Her medications include albuterol and occasional intranasal corticosteroids. She has no history of sexually transmitted diseases and is sexually active with only her current boyfriend of 5 years. The patient denies genitourinary symptoms. Her temperature is 98°F (36.7°C), blood pressure is 112/74 mmHg, pulse is 63/min, and respirations are 12/min with an oxygen saturation of 99% O2 on room air. Physical examination, including a pelvic exam, shows no abnormalities. The patient is worried because she is back in graduate school and cannot afford another child. Which of the following is the most effective emergency contraception? Answer Choices: A. Copper intrauterine device B. High-dose oral contraceptive therapy C. Levonorgesterel pill D. Ulipristal pill

Reasoning + Answer:

<think>The patient is seeking emergency contraception after potentially unprotected intercourse, and the condom broke. The timing and method of action of each emergency contraceptive option should be considered. The copper intrauterine device (IUD) is highly effective for emergency contraception but requires insertion within 5 days of unprotected sex. High-dose oral contraceptive therapy (COC) is less effective than other methods. Levonorgestrel pills are taken within 72 hours of unprotected sex and are approximately 85% effective. Ulipristal, when used within 120 hours of unprotected sex, is approximately 85% effective and is used in many countries as an alternative to levonorgestrel. Given the patient's need for immediate contraception and the recent intercourse, the levonorgestrel pill is the most effective and commonly used method in many countries. The patient can take this within 72 hours of unprotected sex to reduce the risk of pregnancy. </think> <answer> C . Levonorgesterel pill </answer>

