

Question:

Shock causes:

Answer Choices:

- A. Stagnant hypoxia
- B. Anemic hypoxia
- C. Hypoxic hypoxia
- D. Histotoxic hypoxia

Reasoning + Answer:

<think>

To determine the correct answer , let's analyze each option in relation to shock .

Shock is a systemic condition characterized by inadequate tissue perfusion , leading to inadequate oxygen delivery and potentially life-threatening organ dysfunction . The primary cause of shock can be categorized into several types , including hypoxic hypoxia , anemic hypoxia , and histotoxic hypoxia .

A. Stagnant hypoxia : This type of hypoxia occurs when there is a reduction in blood flow to the tissues , leading to inadequate oxygen delivery . This can occur in conditions such as cardiac arrest or severe cardiac failure . However , it is not the most direct cause of shock .

B. Anemic hypoxia : Anemic hypoxia occurs when there is a decrease in the amount of functional hemoglobin or red blood cells , reducing the ability of the blood to carry oxygen to tissues . This can occur in conditions such as anemia or carbon monoxide poisoning . While anemic hypoxia can contribute to shock , it is not the primary cause .

C. Hypoxic hypoxia : Hypoxic hypoxia , also known as hypoxemic hypoxia , occurs when there is a decrease in the amount of oxygen in the blood , which can be due to various causes such as respiratory failure , high altitude , or pulmonary embolism . This is a common cause of shock , as it directly affects the delivery of oxygen to tissues .

D. Histotoxic hypoxia : Histotoxic hypoxia occurs when there is a disruption in the ability of tissues to use oxygen , often due to the presence of toxins or poisons . This can occur in conditions such as cyanide poisoning . While histotoxic hypoxia can contribute to shock , it is not the primary cause .

Considering the definitions and causes of shock , hypoxic hypoxia is the most direct cause of shock , as it directly affects the delivery of oxygen to tissues .

</think>

<answer>

C. Hypoxic hypoxia

</answer>