

Question:

Shock causes:

Answer Choices:

- A. Stagnant hypoxia
- B. Anemic hypoxia
- C. Hypoxic hypoxia
- D. Histotoxic hypoxia

Reasoning + Answer:

<think>
To determine the correct answer, let's analyze the options given. Shock can be caused by various factors, including inadequate blood flow, leading to insufficient oxygen delivery to tissues. Stagnant hypoxia (A) occurs when there is a lack of blood flow due to cardiac dysfunction, resulting in inadequate oxygen delivery to tissues. This is a common cause of shock, particularly in conditions such as cardiogenic shock.

Anemic hypoxia (B) is caused by a decrease in the amount of hemoglobin or red blood cells, which reduces the capacity of the blood to carry oxygen. While this can lead to tissue hypoxia, it is not the primary cause of shock.

Hypoxic hypoxia (C) occurs when there is a decrease in the partial pressure of oxygen in the air or in the blood. This can be caused by respiratory or altitude-related issues but is not directly related to the causes of shock.

Histotoxic hypoxia (D) is caused by the presence of substances that interfere with the body's ability to use oxygen, such as certain toxins. While this can lead to tissue hypoxia, it is not a direct cause of shock.

Considering the options, stagnant hypoxia is a direct cause of shock due to its impact on blood flow and oxygen delivery to tissues.

</think>
<answer>
A. Stagnant hypoxia
</answer>

