

Question:

Shock causes:

Answer Choices:

- A. Stagnant hypoxia
- B. Anemic hypoxia
- C. Hypoxic hypoxia
- D. Histotoxic hypoxia

Reasoning + Answer:

<think>

To answer this question, we need to identify the cause of shock. Shock is a systemic condition characterized by inadequate perfusion of vital organs, leading to organ dysfunction. It can result from various causes, including inadequate oxygen delivery to the tissues (hypoxia).

A. Stagnant hypoxia refers to a situation where there is impaired blood flow, leading to inadequate oxygen delivery to tissues. This can be caused by cardiac arrest, severe cardiac dysfunction, or significant blood loss. However, this is not the primary cause of shock.

B. Anemic hypoxia occurs when the blood lacks sufficient red blood cells or hemoglobin, impairing oxygen delivery to tissues. This can be caused by anemia, severe bleeding, or carbon monoxide poisoning. While anemia can contribute to shock, it is not the most direct cause.

C. Hypoxic hypoxia refers to inadequate oxygen levels in the air breathed in (hypoxia) or a failure of the lungs to exchange oxygen properly. This can be caused by high altitudes, respiratory failure, or carbon monoxide poisoning. Hypoxic hypoxia is a direct cause of shock, as it impairs the body's ability to deliver oxygen to tissues.

D. Histotoxic hypoxia occurs when certain toxins or substances interfere with the body's ability to use oxygen. This can be caused by carbon monoxide poisoning, cyanide poisoning, or other toxins. While histotoxic hypoxia can contribute to shock, it is not the primary cause.

Considering the options, the most direct cause of shock is related to inadequate oxygen delivery to tissues, which is best described by option C. Hypoxic hypoxia is a direct cause of shock, as it impairs the body's ability to deliver oxygen to tissues.

</think>

<answer>

C. Hypoxic hypoxia

</answer>

2

1

0

-1

-2