

Question:

Shock causes:

Answer Choices:

- A. Stagnant hypoxia
- B. Anemic hypoxia
- C. Hypoxic hypoxia
- D. Histotoxic hypoxia

Reasoning + Answer:

mobx

Shock leads to inadequate tissue perfusion , resulting in reduced oxygen delivery to tissues . The primary issue is that oxygen cannot reach tissues adequately due to poor blood flow , even if oxygen content in the blood is normal . This condition is classified as hypoxic hypoxia , where the tissue is deprived of oxygen because of insufficient oxygen delivery due to impaired perfusion .

Let’s evaluate the options :

A . Stagnant hypoxia – This occurs when blood flow is reduced but oxygen content is normal , and the tissue is hypoxic due to poor blood flow . It is a form of hypoxic hypoxia and can be seen in shock , but the term "stagnant hypoxia " is more specific to reduced blood flow without oxygen content change . However , in shock , the main issue is reduced delivery , which is more precisely described as hypoxic hypoxia .

B . Anemic hypoxia – This occurs due to reduced hemoglobin (e.g ., hemorrhage), leading to decreased oxygen-carrying capacity . While hemorrhagic shock can cause anemia , the core pathophysiology of shock is not primarily anemia per se .

C . Hypoxic hypoxia – This is defined by reduced oxygen tension in the tissues due to inadequate oxygen delivery from the blood , typically due to poor perfusion . This is the hallmark of shock . The tissues are hypoxic because blood flow is insufficient , even if oxygen content is normal .

D . Histotoxic hypoxia – This occurs when cells cannot utilize oxygen (e.g ., cyanide poisoning), not due to delivery issues .

In shock , the primary mechanism is reduced perfusion leading to inadequate oxygen delivery to tissues , which is the definition of hypoxic hypoxia .

< answer >

C . Hypoxic hypoxia

</ answer >