

Question:
Which of the following is the causative organism for formation of Baholin cyst?
Answer Choices:
A. Candida
B. Anaerobes
C. Neissea
D. Trichomonas

Reasoning + Answer:

<tool_call>

Baholin cyst, also known as a cystic lesion of the genital area, is most commonly associated with infection by Neisseria gonorrhoeae. This organism is a gram-negative diplococcus and is a primary pathogen in the development of genital infections, including the formation of cysts in the genital tract. The cysts are typically seen in the male genital region and are linked to chronic or recurrent infections due to Neisseria gonorrhoeae, especially in the context of urethral or periurethral inflammation.

Candida (A) is a common cause of candidiasis, particularly in the vulvovaginal region, but not specifically associated with Baholin cysts.

Anaerobes (B) are often involved in abscesses or infections in the perianal or pelvic regions, but not classically linked to Baholin cysts.

Trichomonas (D) is a protozoan causing trichomoniasis, a common sexually transmitted infection, but not known to cause cystic lesions like Baholin cysts.

Neisseria (C) is the correct causative organism, as it is historically and clinically linked to the formation of Baholin cysts, particularly in the context of chronic genital infections.

Therefore, the causative organism for the formation of Baholin cyst is Neisseria.

<answer>

C. Neissea

</answer>

