

Question:

Shock causes:

Answer Choices:

- A. Stagnant hypoxia
- B. Anemic hypoxia
- C. Hypoxic hypoxia
- D. Histotoxic hypoxia

Reasoning + Answer:

<tool_call>

Shock leads to inadequate blood flow to tissues , resulting in reduced oxygen delivery to cells . This is primarily due to impaired perfusion , not a decrease in oxygen content in the blood or a problem with oxygen utilization .

In hypoxic hypoxia , there is a reduced oxygen partial pressure in the alveoli or blood , such as in pulmonary edema or hypoxemia . This is not the primary mechanism in shock .

Anemic hypoxia occurs when there is a reduction in the number of red blood cells or hemoglobin , such as in hemorrhage or hemolysis . While hemorrhagic shock can cause anemia , the primary hypoxia in shock is due to poor perfusion , not reduced oxygen-carrying capacity .

Histotoxic hypoxia occurs when cells cannot use oxygen due to toxins like cyanide , which inhibit cytochrome oxidase . This is not relevant to shock .

Stagnant hypoxia occurs when blood flow to tissues is reduced or absent , leading to poor delivery of oxygen to tissues . This is precisely what happens in shock âĖĖ due to impaired circulation , tissues receive insufficient oxygen despite adequate oxygen content in the blood .

Therefore , the most appropriate type of hypoxia in shock is stagnant hypoxia , as it reflects the impaired blood flow and tissue perfusion .

<answer>

A. Stagnant hypoxia

</answer>

