

Question:

Shock causes:

Answer Choices:

- A. Stagnant hypoxia
- B. Anemic hypoxia
- C. Hypoxic hypoxia
- D. Histotoxic hypoxia

Reasoning + Answer:

Shock is a life-threatening condition where the body's circulatory system fails to deliver sufficient oxygen and nutrients to the body's tissues. It can be caused by a variety of factors, including severe blood loss, sepsis, heart failure, and severe allergic reactions, among others. Shock leads to inadequate perfusion of tissues, which can result in hypoxia (lack of oxygen at the tissue level).

Let's examine the types of hypoxia mentioned in the answer choices:

1. **Stagnant Hypoxia**: This type of hypoxia occurs when blood flow is reduced to a point where it is not enough to deliver oxygen to tissues, even though the oxygen content in the blood remains normal. This can happen in conditions like shock, where circulation is severely compromised.

2. **Anemic Hypoxia**: This type of hypoxia is due to a decrease in the oxygen-carrying capacity of the blood, such as in cases of severe anemia where there are not enough red blood cells to carry oxygen.

3. **Hypoxic Hypoxia**: This type of hypoxia occurs when the partial pressure of oxygen in the inhaled air is too low to adequately oxygenate the blood, as seen at high altitudes.

4. **Histotoxic Hypoxia**: This type of hypoxia occurs when tissues are unable to use the oxygen that is available in the blood, such as in cases of cyanide poisoning.

Given that shock directly affects the circulation and reduces the delivery of oxygen to tissues, the primary form of hypoxia that results from shock is stagnant hypoxia.

<think>

Shock primarily leads to stagnant hypoxia due to the reduction in blood flow to tissues, even though the oxygen content in the blood remains normal.

</think>

<answer>

A. Stagnant hypoxia

</answer>