

Question:

Shock causes:

Answer Choices:

- A. Stagnant hypoxia
- B. Anemic hypoxia
- C. Hypoxic hypoxia
- D. Histotoxic hypoxia

Reasoning + Answer:

Shock is a life-threatening condition characterized by inadequate perfusion of tissues and organs , leading to cellular hypoxia and organ dysfunction . It is primarily categorized into four types based on the underlying mechanism : hypovolemic , distributive , cardiogenic , and obstructive shock .

1 . **Hypovolemic Shock**: This type of shock occurs due to a significant loss of blood or body fluids , leading to reduced circulating blood volume . The heart pumps less blood to the tissues , resulting in inadequate tissue perfusion .

2 . **Distributive Shock (e.g ., septic shock , anaphylactic shock)**: In this type , blood vessels become more permeable and dilate , leading to vasodilation and decreased blood pressure , despite adequate circulating blood volume . This results in inadequate perfusion despite normal cardiac output .

3 . **Cardiogenic Shock**: This occurs due to severe heart dysfunction , such as myocardial infarction , leading to decreased cardiac output and inadequate perfusion .

4 . **Obstructive Shock**: This type of shock is characterized by an obstruction to blood flow , such as in massive pulmonary embolism or aortic dissection .

In all types of shock , the primary issue is inadequate tissue perfusion , leading to cellular hypoxia . The type of hypoxia that best describes this scenario is **hypoxic hypoxia** , which is defined as a lack of oxygen reaching the tissues despite normal levels of oxygen in the blood .

- Hypoxic hypoxia can be caused by :
- Reduced arterial oxygen content
 - Impaired oxygen diffusion into tissues
 - Increased oxygen consumption by tissues
 - Impaired blood flow to tissues

In the context of shock , the primary issue is inadequate blood flow to tissues , leading to hypoxic hypoxia .

< answer >
C . Hypoxic hypoxia
< / answer >