

Question:

Shock causes:

Answer Choices:

- A. Stagnant hypoxia
- B. Anemic hypoxia
- C. Hypoxic hypoxia
- D. Histotoxic hypoxia

Reasoning + Answer:

<think>

- 1 . Shock is a clinical syndrome characterized by inadequate perfusion of tissues and organs , leading to cellular hypoxia and metabolic acidosis .
- 2 . The primary mechanism of hypoxia in shock is related to the reduced blood flow and oxygen delivery to tissues rather than a primary defect in oxygen utilization .
- 3 . Stagnant hypoxia occurs when blood flow is reduced but the oxygen content of the blood remains normal , which can happen in shock due to decreased cardiac output or vasodilation .
- 4 . Anemic hypoxia is caused by a reduced oxygen-carrying capacity of the blood , which is not the primary mechanism in shock .
- 5 . Hypoxic hypoxia is caused by a reduced oxygen supply to tissues , which matches the scenario of inadequate perfusion in shock .
- 6 . Histotoxic hypoxia is when tissues are unable to utilize oxygen despite normal oxygen supply , which is not the primary mechanism in shock .

Given these points , the most appropriate answer is stagnant hypoxia .

</think>

<answer>

A . Stagnant hypoxia

</answer>

