

Question:

Shock causes:

Answer Choices:

- A. Stagnant hypoxia
- B. Anemic hypoxia
- C. Hypoxic hypoxia
- D. Histotoxic hypoxia

Reasoning + Answer:

<think>

To determine the correct answer , we need to understand the mechanisms of shock and how they relate to the different types of hypoxia :

- 1 . **** Stagnant Hypoxia ****: This occurs when there is a reduced blood flow to tissues without a change in oxygen content of the blood . Shock can lead to stagnant hypoxia due to decreased cardiac output and perfusion to tissues .
- 2 . **** Anemic Hypoxia ****: This is characterized by a reduced number of red blood cells or hemoglobin , leading to decreased oxygen -carrying capacity of the blood . Shock itself does not directly cause an emia , but an emia can be a cause of hypoxia .
- 3 . **** Hypoxic Hypoxia ****: This type of hypoxia occurs when the oxygen content of the inhaled air is reduced or when there is a failure of oxygen to diffuse into the bloodstream . Shock does not directly reduce the oxygen content of the inhaled air or impair diffusion into the blood .
- 4 . **** Histotoxic Hypoxia ****: This type of hypoxia occurs when the tissues are unable to use the available oxygen effectively , despite normal blood oxygen content . Shock does not directly affect the tissue 's ability to use oxygen ; it is more about the supply of oxygen to the tissues .

Given the context of shock , the primary mechanism is a reduction in blood flow to tissues , leading to inadequate oxygen delivery . This aligns with the concept of stagnant hypoxia , where the blood flow is reduced but the oxygen content remains normal .

</think>

< answer >

A . Stagnant hypoxia

</answer>

