

Question:
A basket contains 25 oranges among which 1 is bad, 20% are unripe, 2 are sour and the rest are good.
How many oranges are good?

Reasoning + Answer:

<tool_call> Reasoning process here :

Step 1 : We are given that the basket contains 25 oranges in total .

Step 2 : Among them , 1 is bad .
So , the number of oranges that are not bad = $25 - 1 = 24$.

Step 3 : 20 % are un ripe .
So , number of un ripe oranges = $20\% \text{ of } 25 = (20/100) \times 25 = 5$.

Step 4 : 2 are sour .

Step 5 : The rest are good .
We are to find how many are good .

So , we subtract the bad , un ripe , and sour oranges from the total , keeping in mind that "bad " and "sour " might overlap or not âĖĶ but the question says " 1 is bad " , " 2 are sour " , and " 20 % are un ripe " , and the rest are good .

- We assume that :
- The bad oranges are a subset of the total .
 - The sour oranges are a subset .
 - The un ripe oranges are a subset .
 - The "rest " are good , meaning not bad , not sour , not un ripe ? Or just not bad ?

But the question says : "the rest are good " .

We are to find how many are good .

So , the good oranges are the ones that are not bad and not sour and not un ripe ? Or just not bad ?

But the problem says : " 1 is bad , 20 % are un ripe , 2 are sour and the rest are good . "

- This suggests that :
- 1 is bad
 - 5 are un ripe
 - 2 are sour
 - The rest are good

But we must be cautious : could there be overlap ?

However , the problem does not give any information about overlap , and asks for how many are good âĖĶ which may imply the oranges that are not bad and not sour and not un ripe ?

But the problem says "the rest are good " , so likely the "good " ones are those that are not bad , not sour , and not un ripe âĖĶ because otherwise , if un ripe or sour are just attributes , and "good " is defined as not bad , then we would have more than just the rest .

But the problem says : "the rest are good " âĖĶ so the total number of oranges minus the bad , un ripe , and sour ones ?

But un ripe and sour may overlap with bad or good .

But since no overlap is mentioned , and the problem says " 1 is bad " , " 20 % are un ripe " , " 2 are sour " , and " the rest are good " , we interpret it as :

Total oranges = 25
Bad = 1
Un ripe = 5
Sour = 2
The rest are good .

Now , to find good oranges , we subtract those that are bad , un ripe , or sour âĖĶ but again , without knowing overlap , we assume that these are disjoint sets .

But even then , we don 't know if un ripe and sour are subsets of good or bad .

But the problem says "the rest are good " , so likely the oranges that are not bad , not un ripe , not sour are good .

But perhaps "good " means not bad , and the rest are good âĖĶ so total good = total - bad = 24 .

But then why mention un ripe and sour ?

So , perhaps the "good " oranges are those that are not bad , not sour , and not un ripe .

But that would be ambiguous .

LetâĖĶs read again :
"A basket contains 25 oranges among which 1 is bad , 20 % are un ripe , 2 are sour and the rest are good . "

- This suggests that :
- 1 is bad
 - 5 are un ripe
 - 2 are sour
 - the rest are good

So , the "rest " are good âĖĶ so we just need to subtract the ones that are bad , un ripe , or sour .

But if there 's overlap