

Question:

A 5-year-old boy is brought to the physician by his mother because he does not “listen to her” anymore. The mother also reports that her son cannot concentrate on any tasks lasting longer than just a few minutes. Teachers at his preschool report that the patient is more active compared to other preschoolers, frequently interrupts or bothers other children, and is very forgetful. Last year the patient was expelled from another preschool for hitting his teacher and his classmates when he did not get what he wanted and for being disruptive during classes. He was born at term via vaginal delivery and has been healthy except for 3 episodes of acute otitis media at the age of 2 years. He has met all developmental milestones. His mother has major depressive disorder and his father has Graves' disease. He appears healthy and well nourished. Examination shows that the patient does not seem to listen when spoken to directly. The remainder of the examination shows no abnormalities. Which of the following is the most appropriate next step in treatment?

Answer Choices:

- A. Behavior therapy
- B. Methimazole
- C. Fluoxetine
- D. Hearing aids

Reasoning + Answer:

<think> I am thinking about the symptoms and potential causes of the 5-year-old boy's behavior. He is exhibiting symptoms of attention deficit hyperactivity disorder (ADHD), such as difficulty concentrating, impulsivity, and hyperactivity. He also has a history of aggression and disruptive behavior. I need to consider potential underlying medical causes for these symptoms, such as hearing impairment, which could be contributing to his difficulties with listening and concentrating. </think><answer> D. Hearing aids </answer>

