

Question:

A 5-year-old boy is brought to the physician by his mother because he does not “listen to her” anymore. The mother also reports that her son cannot concentrate on any tasks lasting longer than just a few minutes. Teachers at his preschool report that the patient is more active compared to other preschoolers, frequently interrupts or bothers other children, and is very forgetful. Last year the patient was expelled from another preschool for hitting his teacher and his classmates when he did not get what he wanted and for being disruptive during classes. He was born at term via vaginal delivery and has been healthy except for 3 episodes of acute otitis media at the age of 2 years. He has met all developmental milestones. His mother has major depressive disorder and his father has Graves' disease. He appears healthy and well nourished. Examination shows that the patient does not seem to listen when spoken to directly. The remainder of the examination shows no abnormalities. Which of the following is the most appropriate next step in treatment?

Answer Choices:

- A. Behavior therapy
- B. Methimazole
- C. Fluoxetine
- D. Hearing aids

Reasoning + Answer:

<think>

- The patient shows signs of hyperactivity , impulsivity , and difficulty concentrating , which are typical of ADHD (Attention - Deficit /Hyperactivity Disorder).

- The patient's mother has major depressive disorder and his father has Graves ' disease , which are not directly related to ADHD .

- The patient has no obvious physical abnormalities , so hearing aids are not indicated .

- Behavior therapy is often the first-line treatment for ADHD in young children .

- Fluoxetine is used to treat depression and anxiety , and methimazole is used to treat Graves ' disease , neither of which are indicated for ADHD .

- Therefore , the most appropriate next step in treatment is behavior therapy .

- The patient's symptoms are consistent with ADHD .

- The mother's and father's medical conditions are not relevant to the patient's symptoms .

- Behavior therapy is the most appropriate first-line treatment for ADHD in young children .

- Hearing aids are not indicated for this patient .

- Fluoxetine and methimazole are not indicated for ADHD .

- Behavior therapy is the best option for managing the patient's symptoms .

- The most appropriate next step in treatment is behavior therapy .

- The patient's symptoms are consistent with ADHD .

- Behavior therapy is the most appropriate first-line treatment for ADHD in young children .

- The patient's symptoms are not consistent with hearing loss or thyroid disorders .

- Behavior therapy is the best option for managing the patient's symptoms .

- The patient's symptoms are not consistent with depression or thyroid disorders .

- Behavior therapy is the best option for managing the patient's symptoms .

</think>

<answer>A . Behavior therapy </answer>