

Question:

A baby is born after the 32nd gestational week by cesarean delivery. The mother suffered from gestational diabetes; however, she had no other pregnancy-related diseases and was otherwise healthy. The baby has a blood pressure of 100/58 mm Hg, heart rate of 104/min, and oxygen saturation of 88%. The child has tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis. The cyanosis is responding well to initial administration of oxygen. The nasogastric tube was positioned without problems. Which of the following is the most likely diagnosis?

Answer Choices:

- A. Neonatal respiratory distress syndrome (NRDS)
- B. Tracheoesophageal fistula
- C. Pneumonia
- D. Congenital heart anomaly with right-to-left shunt

Reasoning + Answer:

<think>

- The baby is born at 32 weeks, indicating pre term birth.
- The mother had gestational diabetes but no other pregnancy-related diseases.
- The baby has tachypnea, retractions, cyanosis, and responds well to oxygen, which are signs of respiratory distress.
- There is no indication of a structural defect like a tracheoesophageal fistula.
- Pneumonia would not be indicated by the immediate response to oxygen.
- The most likely diagnosis among the given options is Neonatal Respiratory Distress Syndrome (NRDS), as it is common in pre term infants and can present with the described symptoms.

Answer Choices :

- A. Neonatal respiratory distress syndrome (NRDS)
- B. Tracheoesophageal fistula
- C. Pneumonia
- D. Congenital heart anomaly with right-to-left shunt

</think>

<answer>A. Neonatal respiratory distress syndrome (NRDS)</answer>