

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>

The factors that are associated with a decreased risk of Alzheimer's disease include higher education , being married , not smoking , not drinking alcohol regularly , and eating a balanced diet that includes foods rich in omega - 3 fatty acids , such as fish . Among the options provided , living in a city (C) is not a factor that is typically associated with a decreased risk of Alzheimer's . Instead , it is often associated with a higher risk due to increased exposure to air pollution and other environmental toxins . Being African or Hispanic American (A) is not typically associated with a decreased risk of Alzheimer's , as there is no clear evidence to suggest that ethnicity is a risk factor for the disease . Higher education (B) is actually associated with a lower risk of Alzheimer's , as people with higher education tend to have better access to healthcare and other resources that can help prevent and manage the disease . Regular exercise (not listed) is also associated with a lower risk of Alzheimer's . Being married (D) is also associated with a lower risk of Alzheimer's , as it is often associated with social support and other benefits that can help prevent the disease . Having a family history of Alzheimer's (E) is actually associated with a higher risk of the disease , as it suggests a genetic component . High intake of saturated fats (F) is associated with an increased risk of Alzheimer's , as it can contribute to the development of atherosclerosis and other cardiovascular diseases . Older age (G) is associated with an increased risk of Alzheimer's , as the risk of the disease increases with age . Drinking alcohol regularly (J) is associated with an increased risk of Alzheimer's , as excessive alcohol consumption can contribute to liver disease and other health problems . Having a higher intake of saturated fats (F) is associated with an increased risk of Alzheimer's , as it can contribute to the development of atherosclerosis and other cardiovascular diseases .

