

Question:

Which of the following factors is associated with a decreased risk of Alzheimer's?

Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

<think>

To determine which factor is associated with a decreased risk of Alzheimer's , let's analyze the relationship between each option and Alzheimer's disease .

- 1 . Being African or Hispanic American (A): Research suggests that African Americans and Hispanic Americans have a higher risk of Alzheimer's disease compared to non-Hispanic whites . Therefore , this option is unlikely to be associated with a decreased risk .
- 2 . A lower level of education (B): Studies have consistently shown that higher education is linked to a lower risk of Alzheimer's disease . This is often referred to as the "cognitive reserve hypothesis ." Therefore , a lower level of education would likely increase the risk of Alzheimer's .
- 3 . Living in a city (C): Urban vs . rural residence is not a well-established risk factor for Alzheimer's disease . Therefore , this option does not have a clear association with a decreased risk .
- 4 . Being married (D): Marriage has been linked to various health benefits , including a lower risk of cognitive decline and Alzheimer's disease . Social support from a spouse may contribute to this protective effect .
- 5 . Having a family history of Alzheimer's (E): A family history of Alzheimer's is a well-established risk factor for the disease . Therefore , this option is unlikely to be associated with a decreased risk .
- 6 . High intake of saturated fats (F): A diet high in saturated fats has been linked to an increased risk of cardiovascular disease , which is a risk factor for Alzheimer's disease . Therefore , this option is unlikely to be associated with a decreased risk .
- 7 . High intake of fish (H): A diet rich in omega - 3 fatty acids , particularly EPA and DHA found in fish , has been linked to a lower risk of cognitive decline and Alzheimer's disease . Therefore , this option is associated with a decreased risk .
- 8 . Being a smoker (I): Smoking is a well-established risk factor for various health conditions , including cardiovascular disease and lung disease . It is also linked to an increased risk of Alzheimer's disease . Therefore , this option is unlikely to be associated with a decreased risk .
- 9 . Drinking alcohol regularly (J): Moderate alcohol consumption has been linked to a lower risk of certain health conditions , including cardiovascular disease . However , excessive alcohol consumption is a risk factor for various health problems , including cognitive decline . Therefore , this option is not clearly associated with a decreased risk .

Based on the analysis , the factors associated with a decreased risk of Alzheimer's disease are being married (D) and high intake of fish (H). However , the question asks for a single answer , so we will choose the option that is most directly related to a decreased risk of Alzheimer's .

</think>< answer > D </answer>