

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

<think>
To determine which factor is associated with a decreased risk of Alzheimer's , we need to consider the known risk factors and protective factors for the disease .
Several studies have shown that a higher level of education is associated with a lower risk of Alzheimer's , as it may build cognitive reserve and help the brain adapt to age-related changes .
Therefore , a lower level of education , option B , is associated with an increased risk of Alzheimer's , not a decreased risk .
Being African or Hispanic American (option A) and having a family history of Alzheimer's (option E) are also associated with an increased risk of Alzheimer's . Living in a city (option C), older age (option G), high intake of saturated fats (option F), smoking (option I), and drinking alcohol regularly (option J) are also linked to an increased risk of Alzheimer's .
On the other hand , some studies suggest that social engagement , including being married (option D) , may be associated with a lower risk of cognitive decline and Alzheimer's . However , the relationship between marriage and Alzheimer's is complex , and more research is needed to fully understand this association .
Eating fish , which is rich in omega - 3 fatty acids , has been linked to a lower risk of Alzheimer's (option H). Therefore , option H is the correct answer . </think><answer> H </answer>

