

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?

- Answer Choices:
- A. Being African or Hispanic American
 - B. A lower level of education
 - C. Living in a city
 - D. Being married
 - E. Having a family history of Alzheimer's
 - F. High intake of saturated fats
 - G. Older age
 - H. Eating fish
 - I. Being a smoker
 - J. Drinking alcohol regularly

Reasoning + Answer:

reasoning process here

Alzheimer's disease risk factors include age , genetics , education level , diet , and lifestyle . Among the options :

- Being African or Hispanic American is associated with higher risk due to socioeconomic and biological factors .
- Lower education level is linked to increased risk , possibly due to reduced cognitive reserve .
- Living in a city is not clearly linked to decreased risk ; urban living may have mixed effects .
- Being married is associated with a decreased risk of dementia , including Alzheimer's , likely due to social support and engagement .
- Family history increases risk .
- High saturated fat intake is linked to increased risk .
- Older age is a major risk factor .
- Eating fish , particularly fatty fish rich in omega - 3 fatty acids , is associated with a reduced risk of Alzheimer's .
- Smoking increases risk .
- Regular alcohol consumption may have a protective effect in moderation , but excessive drinking increases risk .

However , among the options , " eating fish " is a well -supported dietary factor associated with decreased risk due to omega - 3 fatty acids , which are neuro protective . While marital status has some evidence of protective effect , " eating fish " is a more consistently documented and direct factor linked to reduced Alzheimer's risk in epidemiological studies .

Thus , the best answer supported by current research is eating fish .

< answer >H</answer >