

Question:
Which of the following factors is associated with a decreased risk of Alzheimer's?
Answer Choices:
A. Being African or Hispanic American
B. A lower level of education
C. Living in a city
D. Being married
E. Having a family history of Alzheimer's
F. High intake of saturated fats
G. Older age
H. Eating fish
I. Being a smoker
J. Drinking alcohol regularly

Reasoning + Answer:

</tool_call> reasoning process here
Alzheimer's disease risk factors include age , genetics , education level , diet , and lifestyle . Among the options :
- Being African or Hispanic American is associated with higher risk due to socioeconomic and health disparities .
- Lower education level is linked to increased risk , possibly due to reduced cognitive reserve .
- Living in a city is not consistently linked to decreased risk ; some studies suggest urban living may increase risk due to pollution and stress .
- Being married is associated with a decreased risk of dementia , including Alzheimer's , possibly due to social support and engagement .
- Family history increases risk .
- High saturated fat intake is linked to increased risk .
- Older age is a major risk factor .
- Eating fish , which is rich in omega - 3 fatty acids , is associated with a lower risk of cognitive decline and Alzheimer's .
- Smoking increases risk .
- Regular alcohol consumption , especially in excess , increases risk ; however , moderate intake may have protective effects in some studies , though this is not clearly established .

Among these , " being married " and " eating fish " are both linked to reduced risk . However , " eating fish " is more consistently associated with reduced risk due to omega - 3 s , which support brain health . However , research shows that social factors like marriage can reduce cognitive decline risk through psychological and physical health benefits .

But among the options , the most consistently supported and well - documented protective factor is being married , which has been shown in multiple studies to reduce the risk of dementia .

Thus , the factor most directly associated with decreased risk is being married .

< answer >D </ answer >

