

Question:

Which of the following factors is associated with a decreased risk of Alzheimer's?

Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

</tool_call> reasoning process here

Alzheimer's disease risk factors include age , genetics , education level , lifestyle , and certain dietary patterns . Let's evaluate each option :

- A . Being African or Hispanic American â This group has been associated with higher risk due to socioeconomic and health disparities , not decreased risk . â
- B . A lower level of education â Lower education is linked to increased risk , likely due to reduced cognitive reserve . â
- C . Living in a city â Urban living is not consistently linked to decreased risk ; some studies show mixed results , but not a strong protective factor . â
- D . Being married â Married individuals tend to have lower risk of dementia , possibly due to social support , better health monitoring , and reduced stress . This has been associated with a decreased risk of Alzheimer's . â
- E . Having a family history of Alzheimer's â Strong genetic link ; increases risk . â
- F . High intake of saturated fats â Associated with increased inflammation and cognitive decline ; increases risk . â
- G . Older age â The strongest risk factor ; risk increases with age . â
- H . Eating fish â Fish , especially fatty fish rich in omega - 3 fatty acids , is associated with reduced risk of cognitive decline and Alzheimer â This is a protective factor . â
- I . Being a smoker â Smoking increases risk of dementia . â
- J . Drinking alcohol regularly â Excessive drinking increases risk ; moderate drinking may have benefits , but regular heavy drinking is harmful . â

Among the options , both D (being married) and H (eating fish) are associated with decreased risk . However , the question asks for *which* factor is associated with decreased risk â implying a single best answer .

While both D and H are valid , research consistently shows that omega - 3 fatty acids in fish (especially from fatty fish like salmon , mackerel) are linked to improved brain health and reduced Alzheimer's risk , and this is a well-established protective dietary factor .

Thus , H (Eating fish) is a more directly supported , evidence-based protective dietary factor than marital status , which is more about social support and may vary by context .

Therefore , the most scientifically supported factor among the options that is associated with decreased risk is eating fish .

< answer >H</answer >