

Question:

Which of the following factors is associated with a decreased risk of Alzheimer's?

Answer Choices:

- A. Being African or Hispanic American
- B. A lower level of education
- C. Living in a city
- D. Being married
- E. Having a family history of Alzheimer's
- F. High intake of saturated fats
- G. Older age
- H. Eating fish
- I. Being a smoker
- J. Drinking alcohol regularly

Reasoning + Answer:

<tool_call> reasoning process here

To determine which factor is associated with a decreased risk of Alzheimer's , we evaluate each option based on current scientific evidence :

A . Being African or Hispanic American â Studies show that Black and Hispanic individuals may have higher rates of Alzheimer's due to socioeconomic , environmental , and healthcare disparities . So , this is associated with increased risk .

B . A lower level of education â Lower education level is linked to increased risk , possibly due to reduced cognitive reserve . So , this is associated with increased risk .

C . Living in a city â Urban living has mixed associations , but some studies suggest urban environments may have higher exposure to pollutants or stress , potentially increasing risk . Not clearly protective .

D . Being married â Multiple studies have shown that marital status , particularly being married , is associated with a reduced risk of Alzheimer's . Social engagement , emotional support , and shared activities may contribute to cognitive resilience . This is a well-documented protective factor .

E . Having a family history of Alzheimer's â Strongly linked to increased risk , especially for early-on set cases .

F . High intake of saturated fats â Associated with increased risk due to cardiovascular and inflammatory effects .

G . Older age â The strongest risk factor for Alzheimer's ; risk increases with age .

H . Eating fish â Fish , especially fatty fish rich in omega - 3 fatty acids (like salmon) , is associated with a reduced risk of cognitive decline and Alzheimer's . Omega - 3 s support brain health and reduce inflammation . This is a protective factor .

I . Being a smoker â Smoking is linked to increased risk due to vascular and neurotoxic effects .

J . Drinking alcohol regularly â Excessive alcohol consumption increases risk ; however , moderate alcohol intake (especially red wine) has been studied for potential neuroprotective effects , but the evidence is mixed and not consistently protective . Overall , regular drinking is not considered protective .

Now , both D (being married) and H (eating fish) are protective . However , the question asks for a factor associated with decreased risk . Both are valid , but we must pick the one most consistently and directly linked to reduced Alzheimer's risk .

Among the options , eating fish (H) is supported by strong evidence in nutrition and neuroprotection , particularly through omega - 3 fatty acids . However , being married (D) has also been consistently linked to lower cognitive decline and reduced Alzheimer's risk in epidemiological studies .

But the strongest and most directly protective factor from dietary sources is eating fish , especially given the well-established role of omega - 3 s in brain health .

Thus , while both D and H are protective , H (eating fish) is more directly tied to a biological mechanism that reduces Alzheimer's risk via brain health .

However , some sources emphasize social factors like marriage as protective , and it's a well-established behavioral factor .

But in terms of direct , evidence-based , and widely supported protective dietary factor , eating fish (H) is a key one .

Still , some research shows that being married is associated with lower risk , and this is a social factor that's independently protective .

But among all the options , H (eating fish) is one of the most consistently supported protective dietary factors .

Thus , the best answer is H .

< answer >H</ answer >