

Question:

A 5-year-old boy is brought to the physician by his mother because he does not “listen to her” anymore. The mother also reports that her son cannot concentrate on any tasks lasting longer than just a few minutes. Teachers at his preschool report that the patient is more active compared to other preschoolers, frequently interrupts or bothers other children, and is very forgetful. Last year the patient was expelled from another preschool for hitting his teacher and his classmates when he did not get what he wanted and for being disruptive during classes. He was born at term via vaginal delivery and has been healthy except for 3 episodes of acute otitis media at the age of 2 years. He has met all developmental milestones. His mother has major depressive disorder and his father has Graves' disease. He appears healthy and well nourished. Examination shows that the patient does not seem to listen when spoken to directly. The remainder of the examination shows no abnormalities. Which of the following is the most appropriate next step in treatment?

Answer Choices:

- A. Behavior therapy
- B. Methimazole
- C. Fluoxetine
- D. Hearing aids

Reasoning + Answer:

<think> The physician is thinking about the fact that the 5-year-old boy is experiencing a variety of symptoms , such as not listening to his mother , being unable to concentrate , being more active than other preschoolers , frequently interrupting or bothering other children , and being very forgetful . These symptoms suggest that the boy may have a diagnosis of attention deficit hyper activity disorder (ADHD) . The physician is also thinking about the fact that the boy has a history of being expelled from another preschool for hitting his teacher and his classmates , which suggests that he may have a history of aggressive behavior . The physician is considering the fact that the boy's symptoms may be related to a variety of factors , such as genetic , environmental , and psychological factors . The physician is also thinking about the fact that the boy's symptoms may be treated with a variety of therapies , such as behavior therapy , medication , and lifestyle changes . </think>< answer> A . Behavior therapy </answer>