

Question:

A baby is born after the 32nd gestational week by cesarean delivery. The mother suffered from gestational diabetes; however, she had no other pregnancy-related diseases and was otherwise healthy. The baby has a blood pressure of 100/58 mm Hg, heart rate of 104/min, and oxygen saturation of 88%. The child has tachypnea, subcostal and intercostal retractions, nasal flaring, and cyanosis. The cyanosis is responding well to initial administration of oxygen. The nasogastric tube was positioned without problems. Which of the following is the most likely diagnosis?

Answer Choices:

- A. Neonatal respiratory distress syndrome (NRDS)
- B. Tracheoesophageal fistula
- C. Pneumonia
- D. Congenital heart anomaly with right-to-left shunt

Reasoning + Answer:

<think>
pneumonia the key observation for neonatal patient is respiratory distress
tachypnea - more than 60 breaths per minute
subcostal retraction - when the intercostal muscles rise and fall
crisis - when not enough oxygen
hypoxia the baby has below the normal ranges
oxygen saturation 88% - it is low
the cyanosis is responding well to administration of oxygen
NRDS caused when there is an obstruction in lungs preventing the air to get in and out and is there for
called a respiratory distress
<answer>
neonatal respiratory distress syndrome (NRDS) </answer>
</think>