

Question:

A 38-year-old man comes to the physician because of progressive pain and swelling of his left knee for the past 2 days. He has been taking ibuprofen for the past 2 days without improvement. Four days ago, he scraped his left knee while playing baseball. He has a 2-month history of progressive pain and stiffness in his back. The pain starts after waking up and lasts for 20 minutes. He has type 2 diabetes mellitus. His older sister has rheumatoid arthritis. He is 170 cm (5 ft 7 in) tall and weighs 91 kg (201 lb); BMI is 31.5 kg/m². Temperature is 39°C (102.2°F), pulse is 90/min, and blood pressure is 135/85 mm Hg. Examination shows an erythematous, tender, and swollen left knee; range of motion is limited. There are abrasions over the lateral aspect of the left knee. The remainder of the examination shows no abnormalities. Laboratory studies show a leukocyte count of 13,500/mm³ and an erythrocyte sedimentation rate of 70 mm/h. Which of the following is the most appropriate next step in management?

Answer Choices:

- A. Bone scan
- B. MRI of the left knee
- C. Blood cultures
- D. Arthrocentesis of the left knee

Reasoning + Answer:

<think> The patient had progressive pain and swelling of the left knee for 2 days . He had a history of back pain and stiffness . The examination showed an ery th emat ous , tender , and swollen left knee with abras ions . Laboratory studies showed elevated leuk ocyte count and ery thro cyte sediment ation rate . The patient had type 2 diabetes and a family history of rhe umat oid arthritis . </think>< answer > MR Imaging of the left knee </answer>

